

An Introduction to Large Deviations

Vitalii Konarovskiy
vitali.konarovskiy@uni-hamburg.de

February 16, 2021

Abstract

The large deviations theory is one of the key techniques of modern probability. It concerns with the study of probabilities of rare events and its estimates is the crucial tool required to handle many questions in statistical mechanics, engineering, applied probability, statistics etc. The course is build as the first look at the theory and is oriented on master and PhD students.

1 Introduction and some examples

1.1 Introduction

We start from the considering of a coin-tossing experiment. Let us assume that we toss a fair coin. The law of large numbers says us that the frequency of occurrence of “heads” becomes close to $\frac{1}{2}$ as the number of trials increases to infinity. In other words, if X_1, X_2, \dots are independent random variables taking values 0 and 1 with probabilities $\frac{1}{2}$, i.e. $\mathbb{P}\{X_k = 0\} = \mathbb{P}\{X_k = 1\} = \frac{1}{2}$, then we know that for the empirical mean $\frac{1}{n}S_n = \frac{X_1 + \dots + X_n}{n}$

$$\mathbb{P}\left\{\left|\frac{1}{n}S_n - \frac{1}{2}\right| > \varepsilon\right\} \rightarrow 0^1, \quad n \rightarrow \infty,$$

or more strongly

$$\frac{1}{n}S_n \rightarrow \frac{1}{2} \text{ a.s.}^2, \quad n \rightarrow \infty.$$

We are going stop more precisely on the probabilities $\mathbb{P}\left\{\left|\frac{1}{n}S_n - \frac{1}{2}\right| > \varepsilon\right\}$. We see that this events becomes more unlikely for large n and their probabilities decay to 0. During the course, we will work with such kind of unlike events and will try to understand the rate of their decay to zero. The knowledge of decay of probabilities of such unlike events has many applications in insurance, information theory, statistical mechanics etc. The aim of the course is to give an introduction to one of the key technique of the modern probability which is called the **large deviation theory**.

Before to investigate the rate of decay of the probabilities $\mathbb{P}\left\{\left|\frac{1}{n}S_n - \frac{1}{2}\right| > \varepsilon\right\}$, we consider an example of other random variable where computations are much more simpler.

Let ξ_1, ξ_2, \dots be independent identically distribyted random variables. We also assume that

$$\begin{aligned} \mathbb{E} \xi_1 &= \mu \in \mathbb{R}, \\ \text{Var} \xi_1 &= \sigma^2 > 0, \end{aligned}$$

¹according to the *weak law of large numbers*

²according to the *strong law of large numbers*

and denote $S_n = \xi_1 + \dots + \xi_n$. Then the weak law of large numbers says that

$$\frac{1}{n}S_n \rightarrow \mu \quad \text{in probability, as } n \rightarrow +\infty,$$

that is, for all $\varepsilon > 0$ one has $\mathbb{P} \left\{ \left| \frac{1}{n}S_n - \mu \right| \geq \varepsilon \right\} \rightarrow 0$ as $n \rightarrow +\infty$. This convergence simply follows from Chebyshev's inequality. Indeed,

$$\mathbb{P} \left\{ \left| \frac{1}{n}S_n - \mu \right| \geq \varepsilon \right\} \leq \frac{1}{\varepsilon^2} \mathbb{E} \left(\frac{1}{n}S_n - \mu \right)^2 = \frac{1}{\varepsilon^2} \text{Var} \left(\frac{1}{n}S_n \right) = \frac{\sigma^2}{n\varepsilon^2} \rightarrow 0, \quad n \rightarrow +\infty. \quad (1)$$

Estimate (1) shows that the rate of convergence must be at least $\frac{1}{n}$, but this estimate is too rough. Later we will see that those probabilities decay exponentially fast. Let us demonstrate it on a particular example.

Example 1.1. Let ξ_1, ξ_2, \dots be independent normal distributed random variables with mean $\mu = 0$ and variance $\sigma = 1$ (shortly $\xi_k \sim N(0, 1)$). Then the random variable S_n has the normal distribution with mean 0 and variance n . This implies $\frac{1}{\sqrt{n}}S_n \sim N(0, 1)$.

Now we can consider for $x > 0$

$$\mathbb{P} \left\{ \frac{1}{n}S_n \geq x \right\} = \mathbb{P} \left\{ \frac{1}{\sqrt{n}}S_n \geq x\sqrt{n} \right\} = \frac{1}{\sqrt{2\pi}} \int_{x\sqrt{n}}^{+\infty} e^{-\frac{y^2}{2}} dy \sim \frac{1}{\sqrt{2\pi}x\sqrt{n}} e^{-\frac{nx^2}{2}}, \quad n \rightarrow +\infty,$$

by Exercise 1.1 below. Thus, we have for $x > 0$

$$\begin{aligned} \lim_{n \rightarrow \infty} \frac{1}{n} \ln \mathbb{P} \left\{ \frac{1}{n}S_n \geq x \right\} &= \lim_{n \rightarrow \infty} \frac{1}{n} \ln \frac{1}{\sqrt{2\pi}x\sqrt{n}} e^{-\frac{nx^2}{2}} \\ &= - \lim_{n \rightarrow \infty} \frac{1}{n} \ln \sqrt{2\pi}x\sqrt{n} - \lim_{n \rightarrow \infty} \frac{x^2}{2} = -\frac{x^2}{2} \end{aligned}$$

due to Exercise 1.2 3).

Remark 1.1. By symmetry, one can show that

$$\lim_{n \rightarrow \infty} \frac{1}{n} \ln \mathbb{P} \left\{ \frac{1}{n}S_n \leq x \right\} = -\frac{x^2}{2}$$

for all $x < 0$. Indeed,

$$\lim_{n \rightarrow \infty} \frac{1}{n} \ln \mathbb{P} \left\{ \frac{1}{n}S_n \leq x \right\} = \lim_{n \rightarrow \infty} \frac{1}{n} \ln \mathbb{P} \left\{ -\frac{1}{n}S_n \geq -x \right\} = \lim_{n \rightarrow \infty} \frac{1}{n} \ln \mathbb{P} \left\{ \frac{1}{n}S_n \geq -x \right\} = -\frac{(-x)^2}{2}$$

because $-\xi_k \sim N(0, 1)$ for $k \geq 1$.

Exercise 1.1. Show that

$$\int_x^{+\infty} e^{-\frac{y^2}{2}} dy \sim \frac{1}{x} e^{-\frac{x^2}{2}}, \quad x \rightarrow +\infty.$$

Exercise 1.2. Let $(a_n)_{n \geq 1}$ and $(b_n)_{n \geq 1}$ be two sequences of positive real numbers. We say that they are **logarithmically equivalent** and write $a_n \simeq b_n$ if

$$\lim_{n \rightarrow \infty} \frac{1}{n} (\ln a_n - \ln b_n) = 0.$$

1. Show that $a_n \simeq b_n$ iff $b_n = a_n e^{o(n)}$.
2. Show that $a_n \sim b_n$ implies $a_n \simeq b_n$ and that the inverse implication is not correct.
3. Show that $a_n + b_n \simeq \max\{a_n, b_n\}$.

Exercise 1.3. Let ξ_1, ξ_2, \dots be independent normal distributed random variables with mean μ and variance σ^2 . Let also $S_n = \xi_1 + \dots + \xi_n$. Compute $\lim_{n \rightarrow \infty} \frac{1}{n} \ln \mathbb{P} \left\{ \frac{1}{n} S_n \geq x \right\}$ for $x > \mu$.

1.2 Coin-tossing

In this section, we come back to the coin-tossing experiment and compute the decay of the probability $\mathbb{P} \left\{ \frac{1}{n} S_n \geq x \right\}$. Let, as before, X_1, X_2, \dots be independent random variables taking values 0 and 1 with probabilities $\frac{1}{2}$ and $S_n = X_1 + \dots + X_n$ denote their partial sum. We will show that

$$\lim_{n \rightarrow \infty} \frac{1}{n} \ln \mathbb{P} \left\{ \frac{1}{n} S_n \geq x \right\} = -I(x) \quad (2)$$

for all $x \geq \frac{1}{2}$, where I is some function of x .

We first note that for $x > 1$

$$\lim_{n \rightarrow \infty} \frac{1}{n} \ln \mathbb{P} \left\{ \frac{1}{n} S_n \geq x \right\} = -\infty.^3$$

Next, for $x \in [\frac{1}{2}, 1]$ we observe that

$$\mathbb{P} \left\{ \frac{1}{n} S_n \geq x \right\} = \mathbb{P} \{ S_n \geq xn \} = \sum_{k \geq xn} \mathbb{P} \{ S_n = k \} = \frac{1}{2^n} \sum_{k \geq xn} C_n^k,$$

where $C_n^k = \frac{n!}{k!(n-k)!}$. Then we can estimate

$$\frac{1}{2^n} \max_{k \geq xn} C_n^k \leq \mathbb{P} \{ S_n \geq xn \} \leq \frac{n+1}{2^n} \max_{k \geq xn} C_n^k. \quad (3)$$

Note that the maximum is attained at $k = \lfloor xn \rfloor$, the smallest integer $\geq xn$, because $x \geq \frac{1}{2}$. We denote $l := \lfloor xn \rfloor$. Using Stirling's formula

$$n! = n^n e^{-n} \sqrt{2\pi n} \left(1 + O\left(\frac{1}{n}\right) \right),$$

we have

$$\begin{aligned} \lim_{n \rightarrow \infty} \frac{1}{n} \ln \max_{k \geq xn} C_n^k &= \lim_{n \rightarrow \infty} \frac{1}{n} \ln C_n^l = \lim_{n \rightarrow \infty} \frac{1}{n} (\ln n! - \ln l! - \ln(n-l)!) \\ &= \lim_{n \rightarrow \infty} \left(\ln n - 1 - \frac{l}{n} \ln l + \frac{l}{n} - \frac{n-l}{n} \ln(n-l) + \frac{n-l}{n} \right) \\ &= \lim_{n \rightarrow \infty} \left(\frac{l}{n} \ln n + \frac{n-l}{n} \ln n - \frac{l}{n} \ln l - \frac{n-l}{n} \ln(n-l) \right) \\ &= \lim_{n \rightarrow \infty} \left(-\frac{l}{n} \ln \frac{l}{n} - \frac{n-l}{n} \ln \frac{n-l}{n} \right) = -x \ln x - (1-x) \ln(1-x), \end{aligned}$$

³we always assume that $\ln 0 = -\infty$


because $\frac{l}{n} = \frac{\lfloor xn \rfloor}{n} \rightarrow x$ as $n \rightarrow +\infty$. This together with estimate (3) implies

$$\lim_{n \rightarrow \infty} \frac{1}{n} \ln \mathbb{P} \left\{ \frac{1}{n} S_n \geq x \right\} = -\ln 2 - x \ln x - (1-x) \ln(1-x)$$

for all $x \in [\frac{1}{2}, 1]$.

So, we can take

$$I(x) = \begin{cases} \ln 2 + x \ln x + (1-x) \ln(1-x) & \text{if } x \in [0, 1], \\ +\infty & \text{otherwise.} \end{cases} \quad (4)$$



4

Remark 1.2. Using the symmetry, we have that

$$\lim_{n \rightarrow \infty} \frac{1}{n} \ln \mathbb{P} \left\{ \frac{1}{n} S_n \leq x \right\} = -I(x) \quad (5)$$

for all $x \leq \frac{1}{2}$. Indeed,

$$\begin{aligned} \lim_{n \rightarrow \infty} \frac{1}{n} \ln \mathbb{P} \left\{ \frac{1}{n} S_n \leq x \right\} &= \lim_{n \rightarrow \infty} \frac{1}{n} \ln \mathbb{P} \left\{ \frac{n}{n} - \frac{1}{n} S_n \geq 1 - x \right\} \\ &= \lim_{n \rightarrow \infty} \frac{1}{n} \ln \mathbb{P} \left\{ \frac{(1 - X_1) + \dots + (1 - X_n)}{n} \geq 1 - x \right\} = -I(1-x) = -I(x) \end{aligned}$$

because X_k and $1 - X_k$ have the same distribution.

Theorem 1.1. Let ξ_1, ξ_2, \dots be independent Bernoulli distributed random variables with parameter p for some $p \in (0, 1)$, that is, $\mathbb{P} \{ \xi_k = 1 \} = p$ and $\mathbb{P} \{ \xi_k = 0 \} = 1 - p$ for all $k \geq 1$. Let also $S_n = \xi_1 + \dots + \xi_n$. Then for all $x \geq p$

$$\lim_{n \rightarrow \infty} \frac{1}{n} \ln \mathbb{P} \left\{ \frac{1}{n} S_n \geq x \right\} = -I(x),$$

where

$$I(x) = \begin{cases} x \ln \frac{x}{p} + (1-x) \ln \frac{1-x}{1-p} & \text{if } x \in [0, 1], \\ +\infty & \text{otherwise.} \end{cases}$$

⁴The picture was taken from [dH00]

Exercise 1.4. Prove Theorem 1.1.

Exercise 1.5. Using (2) and (5) show that

$$\sum_{n=1}^{\infty} \mathbb{P} \left\{ \left| \frac{S_n}{n} - \frac{1}{2} \right| \geq \varepsilon \right\} < \infty,$$

for all $\varepsilon > 0$. Conclude that $\frac{S_n}{n} \rightarrow \frac{1}{2}$ a.s. as $n \rightarrow \infty$ (*strong law of large numbers*).
 (Hint: Use the Borel-Cantelly lemma to show the convergence with probability 1)

2 Cramer's theorem

2.1 Comulant generating function

The aim of this section is to obtain an analog of Theorem 1.1 for any sequence of independent identically distributed random variables. In order to understand the form of the rate function I , we will make the following computations, trying to obtain the upper bound for $\mathbb{P} \left\{ \frac{1}{n} S_n \geq x \right\}$.

Let ξ_1, ξ_2, \dots be independent identically distributed random variables with mean $\mu \in \mathbb{R}$. Let also $S_n = \xi_1 + \dots + \xi_n$. We fix $x \geq \mu$ and $\lambda > 0$ and use Chebyshev's inequality in order to estimate the following probability

$$\mathbb{P} \left\{ \frac{1}{n} S_n \geq x \right\} = \mathbb{P} \{ S_n \geq xn \} = \mathbb{P} \left\{ e^{\lambda S_n} \geq e^{\lambda xn} \right\} \leq \frac{1}{e^{\lambda xn}} \mathbb{E} e^{\lambda S_n} = \frac{1}{e^{\lambda xn}} \prod_{k=1}^n \mathbb{E} e^{\lambda \xi_k} = \frac{1}{e^{\lambda xn}} \left(\mathbb{E} e^{\lambda \xi_1} \right)^n.$$

Thus, we have

$$\overline{\lim}_{n \rightarrow \infty} \frac{1}{n} \ln \mathbb{P} \{ S_n \geq xn \} \leq \overline{\lim}_{n \rightarrow \infty} \frac{1}{n} \ln e^{-\lambda xn} + \overline{\lim}_{n \rightarrow \infty} \frac{1}{n} \ln \left(\mathbb{E} e^{\lambda \xi_1} \right)^n = -\lambda x + \varphi(\lambda), \quad (6)$$

where $\varphi(\lambda) := \ln \mathbb{E} e^{\lambda \xi_1}$. We also remark that $-\lambda x + \varphi(\lambda) \geq 0$ for all $\lambda \leq 0$, according to Exercise 2.2 below. Therefore, the inequality

$$\overline{\lim}_{n \rightarrow \infty} \frac{1}{n} \ln \mathbb{P} \{ S_n \geq xn \} \leq -\lambda x + \varphi(\lambda)$$

trivially holds for every $\lambda \leq 0$. Taking infimum over all $\lambda \in \mathbb{R}$, we obtain

$$\overline{\lim}_{n \rightarrow \infty} \frac{1}{n} \ln \mathbb{P} \{ S_n \geq xn \} \leq \inf_{\lambda \in \mathbb{R}} \{-\lambda x + \varphi(\lambda)\} = -\sup_{\lambda \in \mathbb{R}} \{\lambda x - \varphi(\lambda)\}.$$

Later we will see that the function $\sup_{\lambda \in \mathbb{R}} \{\lambda x - \varphi(\lambda)\}$ plays an important role, namely, it is exactly the rate function I .

Definition 2.1. Let ξ be a random variable on \mathbb{R} . The function

$$\varphi(\lambda) := \ln \mathbb{E} e^{\lambda \xi}, \quad \lambda \in \mathbb{R},$$

where the infinite values are allowed, is called the **logarithmic moment generating function** or **comulant generating function** associated with ξ .

Example 2.1. We compute the cumulant generating function associated with Bernoulli distributed random variables ξ with parameter $p = \frac{1}{2}$. So, since $\mathbb{P}\{\xi = 1\} = \mathbb{P}\{\xi = 0\} = \frac{1}{2}$, we obtain

$$\varphi(\lambda) = \ln \mathbb{E} e^{\lambda \xi} = \ln \left(e^{\lambda \cdot \frac{1}{2}} + e^{\lambda \cdot 0} \frac{1}{2} \right) = -\ln 2 + \ln \left(e^{\lambda} + 1 \right), \quad \lambda \in \mathbb{R}.$$

Example 2.2. In this example, we will compute the cumulant generating function associated with exponentially distributed random variable ξ with rate γ . We recall that the density of ξ is given by the following formula

$$p_{\xi}(x) = \begin{cases} \gamma e^{-\gamma x} & \text{if } x \geq 0, \\ 0 & \text{if } x < 0. \end{cases}$$

So,

$$\varphi(\lambda) = \ln \int_0^{\infty} e^{\lambda x} \gamma e^{-\gamma x} dx = \ln \int_0^{\infty} \gamma e^{-(\gamma-\lambda)x} dx = \ln \left(-\frac{\gamma}{\gamma-\lambda} e^{-(\gamma-\lambda)x} \Big|_0^{\infty} \right) = \ln \frac{\gamma}{\gamma-\lambda}$$

if $\lambda < \gamma$. For $\lambda \geq \gamma$ trivially $\varphi(\lambda) = +\infty$. Thus,

$$\varphi(\lambda) = \begin{cases} \ln \gamma - \ln(\gamma - \lambda) & \text{if } \lambda < \gamma, \\ +\infty & \text{if } \lambda \geq \gamma. \end{cases}$$

Exercise 2.1. Show that the function φ is convex⁵.

Solution. In order to show the convexity of φ , we will use Hölder's inequality.⁶ We take $\lambda_1, \lambda_2 \in \mathbb{R}$ and $t \in (0, 1)$. Then for $p = \frac{1}{t}$ and $q = \frac{1}{1-t}$

$$\begin{aligned} \varphi(t\lambda_1 + (1-t)\lambda_2) &= \ln \mathbb{E} \left[e^{t\lambda_1 \xi} e^{(1-t)\lambda_2 \xi} \right] \leq \ln \left(\left[\mathbb{E} e^{\lambda_1 \xi} \right]^t \left[\mathbb{E} e^{\lambda_2 \xi} \right]^{1-t} \right) \\ &= t \ln \mathbb{E} e^{\lambda_1 \xi} + (1-t) \ln \mathbb{E} e^{\lambda_2 \xi} = t\varphi(\lambda_1) + (1-t)\varphi(\lambda_2). \end{aligned}$$

Exercise 2.2. Assume that a random variable ξ has a finite first moment $\mathbb{E}\xi = \mu$ and let φ be the cumulant generating function associated with ξ . Show that for every $x \geq \mu$ and all $\lambda \leq 0$

$$\lambda x - \varphi(\lambda) \leq 0.$$

(Hint: Use Jensen's inequality.⁷)

Exercise 2.3. Let φ be a cumulant generating function associated with ξ . Show that the function φ is differentiable in the interior of the domain $\mathcal{D}_{\varphi} := \{x \in \mathbb{R} : \varphi(x) < \infty\}$. In particular, show that $\varphi'(0) = \mathbb{E}\xi$ if $0 \in \mathcal{D}_{\varphi}^{\circ}$.

(Hint: To show the differentiability of φ , it is enough to show that $\mathbb{E} e^{\lambda \xi}$, $\lambda \in \mathbb{R}$, is differentiable. For the differentiability of the latter function, use the definition of the limit, the dominated convergence theorem⁸ and the fact that the function $\frac{e^{\varepsilon a} - 1}{\varepsilon} = \int_0^a e^{\varepsilon x} dx$ increases in $\varepsilon > 0$ for each $a \geq 0$.)

⁵A function $f : \mathbb{R} \rightarrow (-\infty, +\infty]$ is convex if for all $x_1, x_2 \in \mathbb{R}$ and $t \in (0, 1)$, $f(tx_1 + (1-t)x_2) \leq tf(x_1) + (1-t)f(x_2)$

⁶Let $p, q \in (1, +\infty)$, $\frac{1}{p} + \frac{1}{q} = 1$ and ξ, η be random variables. Then $\mathbb{E}(\xi\eta) \leq (\mathbb{E}\xi^p)^{\frac{1}{p}} (\mathbb{E}\eta^q)^{\frac{1}{q}}$.

⁷For any random variable ξ with a finite first moment and a convex function $f : \mathbb{R} \rightarrow \mathbb{R}$, one has $f(\mathbb{E}\xi) \leq \mathbb{E}f(\xi)$.

⁸For the dominated convergence theorem see [Kal02, Theorem 1.21]


2.2 Fenchel-Legendre transform

In this section, we discuss the Fenchel-Legendre transform of a convex function that appeared in the previous section. Let $f : \mathbb{R} \rightarrow (-\infty, +\infty]$ be a convex function.

Definition 2.2. The function


$$f^*(y) := \sup_{x \in \mathbb{R}} \{yx - f(x)\}$$

is called the **Fenchel-Legendre transform** of f .



9

Fenchel-Legendre transformation: definition



10

Fenchel-Legendre transformation of a function f

Exercise 2.4. Show that the Fenchel-Legendre transform of a convex function f is also convex.

(*Hint:* Show first that the supremum of convex functions is a convex function. Then note that the function $\lambda x - \varphi(\lambda)$ is convex in the variable x)

Exercise 2.5. Compute the Fenchel-Legendre transform of the cumulant generating function associated with the Bernoulli distribution with $p = \frac{1}{2}$.

Solution. Let ξ be a Bernoulli distributed random variable with parameter $p = \frac{1}{2}$, i.e. $\mathbb{P} \{ \xi = 1 \} = \mathbb{P} \{ \xi = 0 \} = \frac{1}{2}$. We first write its cumulant generating function:

$$\varphi(\lambda) = -\ln 2 + \ln(1 + e^\lambda)$$

⁹It turns out that the Fenchel-Legendre transform of f^* coincides with f (see e.g. [Swa12, Proposition 2.3]). The picture was taken from [Swa12].

¹⁰The picture was taken from [Swa12].

(see Example 2.1). In order to compute the Fenchel-Legendre transform of φ , we have to find the supremum of the function

$$g(\lambda) := \lambda x - \varphi(\lambda) = \lambda x + \ln 2 - \ln(1 + e^\lambda), \quad \lambda \in \mathbb{R},$$

for every $x \in \mathbb{R}$. So, we fix $x \in \mathbb{R}$ and find

$$g'(\lambda) = x - \frac{e^\lambda}{1 + e^\lambda} = 0.$$

Hence

$$\lambda = \ln \frac{x}{1-x} \quad \text{if } x \in (0, 1)$$

is a local maximum. Due to the convexity of φ , this point is also the global maximum. Consequently,

$$\begin{aligned} \varphi^*(x) &= \sup_{\lambda \in \mathbb{R}} \{\lambda x - \varphi(\lambda)\} = x \ln \frac{x}{1-x} + \ln 2 - \ln \left(1 + \frac{x}{1-x}\right) \\ &= x \ln x - x \ln(1-x) + \ln 2 + \ln(1-x) = \ln 2 + x \ln x + (1-x) \ln(1-x), \quad x \in (0, 1). \end{aligned}$$

If $x < 0$ or $x > 1$ then $\varphi^*(x) = +\infty$. For $x = 0$ and $x = 1$ one can check that $\varphi^*(x) = \ln 2$.

Exercise 2.6. Show that the function φ^* from the previous exercise equals $+\infty$ for $x \in (-\infty, 0) \cup (1, +\infty)$ and $\ln 2$ for $x \in \{0, 1\}$.

Compering the Fenchel-Legendre transformation φ^* of the cumulant generating function associated with the Bernoulli distribution ξ and the rate function I given by (4), we can see that they coincide.

- Exercise 2.7.**
- Show that the Fenchel-Legendre transform of the cumulant generating function associated with $N(0, 1)$ coincides with $\frac{x^2}{2}$.
 - Show that the Fenchel-Legendre transform of the cumulant generating function associated with Bernoulli distribution with parameter $p \in (0, 1)$ coincides with the function I from Theorem 1.1.
 - Find the Fenchel-Legendre transform of the cumulant generating function associated with exponential distribution.

Exercise 2.8. Suppose that φ^* is the Fenchel-Legendre transform of the cumulant generating function of a random variable ξ with $\mathbb{E}\xi = \mu$. Show that

- $\varphi^*(x) \geq 0$ for all $x \in \mathbb{R}$. (*Hint:* Use the fact that $\varphi(0) = 0$)
- $\varphi^*(\mu) = 0$. (*Hint:* Use (i) and Jensen's inequality to show that $\varphi^*(\mu) \leq 0$)
- φ^* increases on $[\mu, \infty)$ and decreases on $(-\infty, \mu]$. (*Hint:* Use the convexity of φ^* (see Exercise 2.4) and (ii))
- $\varphi^*(x) > 0$ for all $x \neq \mu$. (*Hint:* Get a contradiction with the assumption $\varphi^*(x) = 0$ for $x > \mu$)
- Show that φ^* strictly increases on $\{x \geq \mu : \varphi^*(x) < \infty\}$ and strictly decreases on $\{x \leq \mu : \varphi^*(x) < \infty\}$. (*Hint:* Use (iv) and the convexity of φ^*)

2.3 Cramer's theorem

The goal of this section is to prove the Cramer's theorem.

Theorem 2.1 (Cramer). *Let ξ_1, ξ_2, \dots be independent identically distributed random variables with mean $\mu \in \mathbb{R}$ and cumulant generating function φ . Let also $S_n = \xi_1 + \dots + \xi_n$. Then, for every $x \geq \mu$*

$$\lim_{n \rightarrow \infty} \frac{1}{n} \ln \mathbb{P} \left\{ \frac{1}{n} S_n \geq x \right\} = -\varphi^*(x),$$

where φ^* is the Fenchel-Legendre transform of φ .

We will need the following lemma for the proof of Cramer's theorem.

Lemma 2.1. *Let φ be the cumulant generating function associated with a random variable ξ . Let also φ takes finite values on \mathbb{R} . Then φ is continuously differentiable¹¹ on \mathbb{R} , $\varphi'(0) = \mathbb{E} \xi$ and*

$$\lim_{\lambda \rightarrow +\infty} \varphi'(\lambda) = \text{ess sup } \xi. \quad (7)$$

Proof. Using the dominated convergence theorem, similarly as in the solution of Exercise 2.3, one can check that

$$\frac{d}{d\lambda} \mathbb{E} e^{\lambda \xi} = \mathbb{E} \xi e^{\lambda \xi}, \quad \lambda \in \mathbb{R}.$$

Therefore,

$$\varphi'(\lambda) = \frac{\mathbb{E} [\xi e^{\lambda \xi}]}{\mathbb{E} e^{\lambda \xi}}, \quad \lambda \in \mathbb{R}.$$

In particular, this trivially implies the equality $\varphi'(0) = \mathbb{E} \xi$.

Let $\beta := \text{ess sup } \xi$. If $\mathbb{P} \{ \xi = \beta \} > 0$, then the limit of $\varphi'(\lambda)$ can be simply computed as follows

$$\lim_{\lambda \rightarrow +\infty} \varphi'(\lambda) = \lim_{\lambda \rightarrow +\infty} \frac{\mathbb{E} [\xi e^{\lambda \xi}]}{\mathbb{E} e^{\lambda \xi}} = \lim_{\lambda \rightarrow +\infty} \frac{e^{-\lambda \beta} \mathbb{E} [\xi e^{\lambda \xi}]}{e^{-\lambda \beta} \mathbb{E} e^{\lambda \xi}} = \lim_{\lambda \rightarrow +\infty} \frac{\mathbb{E} [\xi e^{-\lambda(\beta - \xi)}]}{\mathbb{E} e^{-\lambda(\beta - \xi)}} = \frac{\mathbb{E} \xi \mathbb{I}_{\{\xi = \beta\}}}{\mathbb{E} \mathbb{I}_{\{\xi = \beta\}}} = \beta, \quad (8)$$

by the dominated convergence theorem and the fact that $e^{-\lambda(\beta - \xi)} \rightarrow \mathbb{I}_{\{\xi = \beta\}}$ a.s. as $\lambda \rightarrow +\infty$.

If $\mathbb{P} \{ \xi = \beta \} = 0$, then the proof is more technical, where one needs to estimate the limit from below by $\beta - \varepsilon$ and then make $\varepsilon \rightarrow 0+$. We leave this as an exercise. \square

Exercise 2.9. Check equality (7) in Lemma 2.1.

Exercise 2.10. Let φ^* be the Fenchel-Legendre transform of the cumulant generating function of a random variable ξ . Let also $\beta = \text{ess sup } \xi < \infty$. Show that $\varphi^*(x) = +\infty$ for all $x > \beta$.

Hint: Show that $\lim_{\lambda \rightarrow +\infty} (\lambda x - \varphi(\lambda)) = +\infty$.

Proof of Theorem 2.1. The upper bound

$$\overline{\lim}_{n \rightarrow \infty} \frac{1}{n} \ln \mathbb{P} \left\{ \frac{1}{n} S_n \geq x \right\} \leq -\varphi^*(x)$$

¹¹One can even prove that φ is infinitely differentiable function on \mathbb{R} .

¹² $\text{ess sup } \xi$ is defined as $\inf \left\{ \sup_{\omega \in A^c} \xi(\omega) : \forall A \ \mathbb{P} \{A\} = 0 \right\}$.

was proved at the beginning of Section 2.1 (see inequality (6)).

We next prove the *lower bound*. For simplicity, we will assume that the cumulant generating function φ takes finite values on \mathbb{R} . The case where $\varphi(\lambda) = +\infty$ for some $\lambda \in \mathbb{R}$ can be found in [Kal02, p. 541]. By Lemma 2.1, the function φ is continuously differentiable on \mathbb{R} with $\varphi'(0) = \mathbb{E}\xi_1 = \mu$ and $\lim_{\lambda \rightarrow +\infty} \varphi'(\lambda) = \text{ess sup } \xi =: \beta$. Since φ' is continuous, for any $y \in (\mu, \beta)$ we can choose $\lambda_0 > 0$ such that $\varphi'(\lambda_0) = y$, by the intermediate value theorem.

Let $\tilde{\xi}_1, \tilde{\xi}_2, \dots$ be independent identically distributed random variables with distribution

$$\mathbb{P} \left\{ \tilde{\xi}_i \in B \right\} = e^{-\varphi(\lambda_0)} \mathbb{E} \left[e^{\lambda_0 \xi_i} \mathbb{I}_{\{\xi_i \in B\}} \right], \quad B \in \mathcal{B}(\mathbb{R}).$$

Then the cumulant generating function $\varphi_{\tilde{\xi}_i}$ associated with $\tilde{\xi}_i$ is defined by

$$\varphi_{\tilde{\xi}_i}(\lambda) = \ln \mathbb{E} e^{\lambda \tilde{\xi}_i} = \ln \left(e^{-\varphi(\lambda_0)} \mathbb{E} \left[e^{\lambda_0 \xi_i} e^{\lambda \xi_i} \right] \right) = \ln e^{-\varphi(\lambda_0)} + \ln \mathbb{E} e^{(\lambda + \lambda_0) \xi_i} = \varphi(\lambda + \lambda_0) - \varphi(\lambda_0).$$

Therefore, $\mathbb{E} \tilde{\xi}_i = \varphi'_{\tilde{\xi}_i}(0) = \varphi'(\lambda_0) = y$. By the law of large numbers, we can conclude that for every $\varepsilon > 0$

$$\mathbb{P} \left\{ \left| \frac{1}{n} \tilde{S}_n - y \right| < \varepsilon \right\} \rightarrow 1, \quad n \rightarrow \infty,$$

where $\tilde{S}_n = \tilde{\xi}_1 + \dots + \tilde{\xi}_n$.

On the other side,

$$\begin{aligned} \mathbb{P} \left\{ \left| \frac{1}{n} \tilde{S}_n - y \right| < \varepsilon \right\} &= e^{-n\varphi(\lambda_0)} \mathbb{E} \left[e^{\lambda_0(\xi_1 + \dots + \xi_n)} \mathbb{I}_{\left\{ \left| \frac{1}{n} S_n - y \right| < \varepsilon \right\}} \right] = e^{-n\varphi(\lambda_0)} \mathbb{E} \left[e^{\lambda_0 S_n} \mathbb{I}_{\left\{ \left| \frac{1}{n} S_n - y \right| < \varepsilon \right\}} \right] \\ &\leq e^{-n\varphi(\lambda_0)} e^{\lambda_0 n(y + \varepsilon)} \mathbb{P} \left\{ \left| \frac{1}{n} S_n - y \right| < \varepsilon \right\} = e^{n(\lambda_0(y + \varepsilon) - \varphi(\lambda_0))} \mathbb{P} \left\{ \left| \frac{1}{n} S_n - y \right| < \varepsilon \right\}. \end{aligned}$$

Consequently,

$$\begin{aligned} \liminf_{n \rightarrow \infty} \frac{1}{n} \ln \mathbb{P} \left\{ \left| \frac{1}{n} S_n - y \right| < \varepsilon \right\} &\geq \liminf_{n \rightarrow \infty} \frac{1}{n} \ln \left[e^{-n(\lambda_0(y + \varepsilon) - \varphi(\lambda_0))} \mathbb{P} \left\{ \left| \frac{1}{n} \tilde{S}_n - y \right| < \varepsilon \right\} \right] \\ &= -(\lambda_0(y + \varepsilon) - \varphi(\lambda_0)) + \liminf_{n \rightarrow \infty} \frac{1}{n} \ln \mathbb{P} \left\{ \left| \frac{1}{n} \tilde{S}_n - y \right| < \varepsilon \right\} \\ &= -(\lambda_0(y + \varepsilon) - \varphi(\lambda_0)) \geq -\sup_{\lambda \in \mathbb{R}} \{\lambda(y + \varepsilon) - \varphi(\lambda)\} = -\varphi^*(y + \varepsilon). \end{aligned}$$

Now, fixing any $x \in [\mu, \beta)$ and putting $y := x + \varepsilon$, we get for small enough $\varepsilon > 0$

$$\liminf_{n \rightarrow \infty} \frac{1}{n} \ln \mathbb{P} \left\{ \frac{1}{n} S_n \geq x \right\} \geq \liminf_{n \rightarrow \infty} \frac{1}{n} \ln \mathbb{P} \left\{ \left| \frac{1}{n} S_n - y \right| < \varepsilon \right\} \geq -\varphi^*(x + 2\varepsilon).$$

Since φ^* is continuous on $[\mu, \beta)$ by convexity, we may pass to the limit as $\varepsilon \rightarrow 0+$. Therefore, we obtain the lower bound.

If $x > \beta$, then $\varphi^*(x) = +\infty$ according to Exercise 2.10. Therefore, the lower bound holds.

For the case $x = \beta < \infty$, we have that

$$\begin{aligned} \liminf_{n \rightarrow \infty} \frac{1}{n} \ln \mathbb{P} \left\{ \frac{1}{n} S_n \geq \beta \right\} &= \liminf_{n \rightarrow \infty} \frac{1}{n} \ln \mathbb{P} \left\{ \xi_1 = \beta, \dots, \xi_n = \beta \right\} \\ &= \liminf_{n \rightarrow \infty} \frac{1}{n} \ln \mathbb{P} \left\{ \xi_1 = \beta \right\}^n = \ln \mathbb{P} \left\{ \xi_1 = \beta \right\}. \end{aligned}$$

We also compute the right hand side for the lower bound:

$$\begin{aligned} \varphi^*(\beta) &= \sup_{\lambda \in \mathbb{R}} \{\lambda\beta - \varphi(\lambda)\} \stackrel{\text{Exe. 2.2}}{=} \sup_{\lambda \geq 0} \{\lambda\beta - \varphi(\lambda)\} = \sup_{\lambda \geq 0} \left\{ -\ln \mathbb{E} e^{-\lambda\beta} - \ln \mathbb{E} e^{\lambda\xi_1} \right\} \\ &= -\inf_{\lambda \geq 0} \left\{ \ln \mathbb{E} e^{-\lambda(\beta - \xi_1)} \right\} = \ln \mathbb{P} \{ \xi_1 = \beta \}. \end{aligned}$$

For the last equality in the previous computations we have used the dominated convergence theorem and the fact that $e^{-\lambda(\beta - \xi_1)} \geq \mathbb{I}_{\{\xi_1 = \beta\}}$, $\lambda > 0$, and $e^{-\lambda(\beta - \xi_1)} \rightarrow \mathbb{I}_{\{\xi_1 = \beta\}}$ a.s. as $\lambda \rightarrow +\infty$. This completes the proof of the theorem. \square

Remark 2.1. Under the assumptions of Theorem 2.1, for every $x \leq \mu$

$$\lim_{n \rightarrow \infty} \frac{1}{n} \ln \mathbb{P} \left\{ \frac{1}{n} S_n \leq x \right\} = -\varphi^*(x). \quad (9)$$

In order to obtain this equality, one needs to apply Cramer's Theorem 2.1 to the family of random variables $2\mu - \xi_1, 2\mu - \xi_2, \dots$ and show that $\varphi_{2\mu - \xi_1}^*(2\mu - x) = \varphi^*(x)$, $x \in \mathbb{R}$.

Exercise 2.11. Check equality (9).

Exercise 2.12. Let ξ_1, ξ_2, \dots be independent identically distributed random variables. Consider a non-negative Borel measurable function $f : \mathbb{R} \rightarrow [0, \infty)$ such that $\mathbb{E} f(\xi_1) \in (0, \infty)$. Define the family of independent random variables η_1, η_2, \dots with distribution

$$\mathbb{P} \{ \eta_i \in B \} = \frac{1}{C} \mathbb{E} [f(\xi_i) \mathbb{I}_{\{\xi_i \in B\}}], \quad B \in \mathcal{B}(\mathbb{R}),$$

where $C = \mathbb{E} f(\xi_i)$ is the normalizing constant.

1. Find the distribution of η_1 , if ξ_1 has the exponential distribution with parameter $\lambda > 0$, and $f(x) = e^{-\alpha x}$, $x \in \mathbb{R}$, where $\alpha > -\lambda$ is a positive constants.
2. Show that for every $n \in \mathbb{N}$ and $B_i \in \mathcal{B}(\mathbb{R})$

$$\mathbb{P} \{ \eta_1 \in B_1, \dots, \eta_n \in B_n \} = \frac{1}{C^n} \mathbb{E} [f(\xi_1) \dots f(\xi_n) \mathbb{I}_{\{\xi_1 \in B_1, \dots, \xi_n \in B_n\}}].$$

3. Show that

$$\mathbb{E} g(\eta_1, \dots, \eta_n) = \frac{1}{C^n} \mathbb{E} [f(\xi_1) \dots f(\xi_n) g(\xi_1, \dots, \xi_n)],$$

for any Borel measurable function $g : \mathbb{R}^n \rightarrow \mathbb{R}$.

3 Definition of large deviation principle

3.1 Large deviations for Gaussian vectors

We recall that, in previous sections, we have investigated the decay of the probability $\mathbb{P} \left\{ \frac{1}{n} S_n \geq x \right\}$, where $S_n = \xi_1 + \dots + \xi_n$ and ξ_k , $k \in \mathbb{N}$, were independent identically distributed random variables in \mathbb{R} . We start this section from some example of random variables in higher dimension and investigate the decay of similar probabilities. This will lead us to the general concept of large deviation principle in the

next section. We note that the case $\xi_k \sim N(0, 1)$ was very easy for computations (see Example 1.1). So similarly, we take independent \mathbb{R}^d -valued random element (or random vector) $\eta_k = (\eta_k^{(1)}, \dots, \eta_k^{(d)})$, $k \geq 1$, with standard Gaussian distributions¹³. We will study the decay of the probability

$$\mathbb{P} \left\{ \frac{1}{n} S_n \in A \right\} = \mathbb{P} \left\{ \frac{1}{\sqrt{n}} \eta \in A \right\} = \mathbb{P} \left\{ \eta \in \sqrt{n} A \right\},$$

where A is a subset of \mathbb{R}^d , $S_n = \eta_1 + \dots + \eta_n$, and η has a standard Gaussian distribution.

The upper bound. We recall that the density of $\frac{1}{\sqrt{n}} \eta$ is given by the formula

$$p_{\frac{\eta}{\sqrt{n}}}(x) = \frac{(\sqrt{n})^d}{(\sqrt{2\pi})^d} e^{-\frac{n\|x\|^2}{2}}, \quad x = (x_1, \dots, x_d) \in \mathbb{R}^d,$$

where $\|x\|^2 = x_1^2 + \dots + x_d^2$. Now, we can estimate

$$\begin{aligned} \liminf_{n \rightarrow \infty} \frac{1}{n} \ln \mathbb{P} \left\{ \frac{1}{\sqrt{n}} \eta \in A \right\} &= \liminf_{n \rightarrow \infty} \frac{1}{n} \ln \int_A p_{\frac{\eta}{\sqrt{n}}}(x) dx = \lim_{n \rightarrow \infty} \frac{1}{n} \ln \frac{(\sqrt{n})^d}{(\sqrt{2\pi})^d} \\ &+ \liminf_{n \rightarrow \infty} \frac{1}{n} \ln \int_A e^{-\frac{n\|x\|^2}{2}} dx \leq \liminf_{n \rightarrow \infty} \frac{1}{n} \ln \int_A \left(\sup_{y \in A} e^{-\frac{n\|y\|^2}{2}} \right) dx \\ &= \liminf_{n \rightarrow \infty} \frac{1}{n} \ln \left(e^{-n \inf_{x \in A} \frac{\|x\|^2}{2}} |A| \right) \leq \liminf_{n \rightarrow \infty} \frac{1}{n} \ln |A| + \liminf_{n \rightarrow \infty} \frac{1}{n} \ln e^{-n \inf_{x \in A} I(x)} \\ &= - \inf_{x \in A} I(x) \leq - \inf_{x \in \bar{A}} I(x), \end{aligned}$$

where $I(x) := \frac{\|x\|^2}{2}$, \bar{A} is the closure of A and $|A|$ denotes the Lebesgue measure of A .

The lower bound. In order to obtain the lower bound, we assume that the interior A° of A is non-empty and fix $x_0 \in A^\circ$. Let $B_r(x_0) = \{x \in \mathbb{R}^d : \|x - x_0\| < r\}$ denotes the ball in \mathbb{R}^d with center x_0 and radius r . We estimate

$$\begin{aligned} \liminf_{n \rightarrow \infty} \frac{1}{n} \ln \mathbb{P} \left\{ \frac{1}{\sqrt{n}} \eta \in A \right\} &= \liminf_{n \rightarrow \infty} \frac{1}{n} \ln \int_A p_{\frac{\eta}{\sqrt{n}}}(x) dx \geq \liminf_{n \rightarrow \infty} \frac{1}{n} \ln \frac{(\sqrt{n})^d}{(\sqrt{2\pi n})^d} \\ &+ \liminf_{n \rightarrow \infty} \frac{1}{n} \ln \int_{B_r(x_0)} e^{-\frac{n\|x\|^2}{2}} dx \geq \liminf_{n \rightarrow \infty} \frac{1}{n} \ln \int_{B_r(x_0)} e^{-\frac{n(\|x_0\| + \|x - x_0\|)^2}{2}} dx \\ &\geq \liminf_{n \rightarrow \infty} \frac{1}{n} \ln \int_{B_r(x_0)} e^{-\frac{n(\|x_0\| + r)^2}{2}} dx = \liminf_{n \rightarrow \infty} \frac{1}{n} \ln e^{-\frac{n(\|x_0\| + r)^2}{2}} |B_r(x_0)| \\ &= - \frac{(\|x_0\| + r)^2}{2}. \end{aligned}$$

Making $r \rightarrow 0+$, we have

$$\liminf_{n \rightarrow \infty} \frac{1}{n} \ln \mathbb{P} \left\{ \frac{1}{\sqrt{n}} \eta \in A \right\} \geq - \frac{\|x_0\|^2}{2}.$$

Now, maximizing the right hand side over all points x_0 from the interior A° , we obtain

$$\liminf_{n \rightarrow \infty} \frac{1}{n} \ln \mathbb{P} \left\{ \frac{1}{\sqrt{n}} \eta \in A \right\} \geq \sup_{x_0 \in A^\circ} \left(- \frac{\|x_0\|^2}{2} \right) = - \inf_{x \in A^\circ} I(x).$$

¹³ $\eta_k^{(i)} \sim N(0, 1)$, $i = 1, \dots, d$, and are independent

Thus, combining the lower and upper bounds, we have prove that for any Borel measurable set A

$$- \inf_{x \in A^\circ} I(x) \leq \underline{\lim}_{n \rightarrow \infty} \frac{1}{n} \ln \mathbb{P} \left\{ \frac{1}{\sqrt{n}} \eta \in A \right\} \leq \overline{\lim}_{n \rightarrow \infty} \frac{1}{n} \ln \mathbb{P} \left\{ \frac{1}{\sqrt{n}} \eta \in A \right\} \leq - \inf_{x \in \bar{A}} I(x). \quad (10)$$

3.2 Definition of large deviation principle

Let $(\xi_\varepsilon)_{\varepsilon > 0}$ be a family of random elements on a metric space E and I be a function from E to $[0, \infty]$.

Definition 3.1. We say that the family $(\xi_\varepsilon)_{\varepsilon > 0}$ satisfies the **large deviation principle (LDP) in E with rate function I** if for any Borel set $A \subset E$ we have

$$- \inf_{x \in A^\circ} I(x) \leq \underline{\lim}_{\varepsilon \rightarrow 0} \varepsilon \ln \mathbb{P} \{ \xi_\varepsilon \in A \} \leq \overline{\lim}_{\varepsilon \rightarrow 0} \varepsilon \ln \mathbb{P} \{ \xi_\varepsilon \in A \} \leq - \inf_{x \in \bar{A}} I(x). \quad (11)$$

We remark that in the case of a countable family of random elements $(\xi_n)_{n \geq 1}$, the large deviation principle corresponds to the statement

$$- \inf_{x \in A^\circ} I(x) \leq \underline{\lim}_{n \rightarrow \infty} a_n \ln \mathbb{P} \{ \xi_n \in A \} \leq \overline{\lim}_{n \rightarrow \infty} a_n \ln \mathbb{P} \{ \xi_n \in A \} \leq - \inf_{x \in \bar{A}} I(x)$$

for some sequence $a_n \rightarrow 0$. In fact, we have proved in the previous section that the family $(\frac{1}{n} S_n)_{n \geq 1}$ or $(\frac{\eta}{\sqrt{n}})_{n \geq 1}$ satisfies the large deviation principle in \mathbb{R}^d with rate function $I(x) = \frac{\|x\|^2}{2}$, $x \in \mathbb{R}^d$ and $a_n = \frac{1}{n}$ (see inequality (10)).

Lemma 3.1. A family $(\xi_\varepsilon)_{\varepsilon > 0}$ satisfies the large deviation principle in E with rate function I iff

$$\overline{\lim}_{\varepsilon \rightarrow 0} \varepsilon \ln \mathbb{P} \{ \xi_\varepsilon \in F \} \leq - \inf_{x \in F} I(x) \quad (12)$$

for every closed set $F \subset E$, and

$$\underline{\lim}_{\varepsilon \rightarrow 0} \varepsilon \ln \mathbb{P} \{ \xi_\varepsilon \in G \} \geq - \inf_{x \in G} I(x) \quad (13)$$

for every open set $G \subset E$.

Proof. We first remark that inequalities (12) and (13) immediately follow from the definition of LDP and the fact that $F = \bar{F}$ and $G = G^\circ$.

To prove (11), we fix a Borel measurable set $A \subset E$ and estimate

$$\begin{aligned} - \inf_{x \in A^\circ} I(x) &\stackrel{(13)}{\leq} \underline{\lim}_{\varepsilon \rightarrow 0} \varepsilon \ln \mathbb{P} \{ \xi_\varepsilon \in A^\circ \} \leq \underline{\lim}_{\varepsilon \rightarrow 0} \varepsilon \ln \mathbb{P} \{ \xi_\varepsilon \in A \} \\ &\leq \overline{\lim}_{\varepsilon \rightarrow 0} \varepsilon \ln \mathbb{P} \{ \xi_\varepsilon \in A \} \leq \overline{\lim}_{\varepsilon \rightarrow 0} \varepsilon \ln \mathbb{P} \{ \xi_\varepsilon \in \bar{A} \} \stackrel{(12)}{\leq} - \inf_{x \in \bar{A}} I(x). \end{aligned}$$

□

Remark 3.1. A similar statement to Lemma 3.1 can be done for a countable family of random elements $(\xi_n)_{n \geq 1}$.

Remark 3.2. We note that repeating the proof from the previous section, one can show that the family $(\sqrt{\varepsilon}\xi)_{\varepsilon>0}$ satisfies the LDP in \mathbb{R}^d with rate function $I(x) = \frac{1}{2}\|x\|^2$, where ξ is a standard Gaussian random vector in \mathbb{R}^d .

Proposition 3.1. *Let there exist a subset E_0 of the metric space E such that*

1) *for each $x \in E_0$*

$$\lim_{r \rightarrow 0^+} \liminf_{\varepsilon \rightarrow 0} \varepsilon \ln \mathbb{P} \{ \xi_\varepsilon \in B_r(x) \} \geq -I(x),$$

where $B_r(x)$ denotes the ball with center x and radius r ;

2) *for each x satisfying $I(x) < \infty$, there exists a sequence $x_n \in E_0$, $n \geq 1$, such that $x_n \rightarrow x$, $n \rightarrow \infty$, and $I(x_n) \rightarrow I(x)$, $n \rightarrow \infty$.*

Then lower bound (13) holds for any open set G .

Proof. First we note that it is enough to prove the lower bound for all open $G \subseteq E$ satisfying $\inf_{x \in G} I(x) < \infty$.

Let δ be an arbitrary positive number. Then there exists $x_0 \in G$ such that

$$I(x_0) < \inf_{x \in G} I(x) + \delta.$$

Hence, by 2) and the openness of G we can find $\tilde{x} \in G \cap E_0$ that satisfies

$$I(\tilde{x}) < I(x_0) + \delta.$$

Next, using 1) and the openness of G , there exists $r > 0$ such that $B_r(\tilde{x}) \subseteq G$ and

$$\liminf_{\varepsilon \rightarrow 0} \varepsilon \ln \mathbb{P} \{ \xi_\varepsilon \in B_r(\tilde{x}) \} \geq -I(\tilde{x}) - \delta.$$

Consequently, we can now estimate

$$\begin{aligned} \liminf_{\varepsilon \rightarrow 0} \varepsilon \ln \mathbb{P} \{ \xi_\varepsilon \in G \} &\geq \liminf_{\varepsilon \rightarrow 0} \varepsilon \ln \mathbb{P} \{ \xi_\varepsilon \in B_r(\tilde{x}) \} \\ &\geq -I(\tilde{x}) - \delta > -I(x_0) - 2\delta > -\inf_{\varphi \in G} I(\varphi) - 3\delta. \end{aligned}$$

Making $\delta \rightarrow 0$, we obtain the lower bound (13). □

Proposition 3.1 shows the local nature of the lower bound. Similarly one can prove a similar result for a countable family of random elements $(\xi_n)_{n \geq 1}$.

Exercise 3.1. Let $(\xi_\varepsilon)_{\varepsilon>0}$ satisfies the LDP in E with rate function I . Show that

a) if A is such that $\inf_{x \in A^\circ} I(x) = \inf_{x \in A} I(x)$, then

$$\lim_{\varepsilon \rightarrow 0} \varepsilon \ln \mathbb{P} \{ \xi_\varepsilon \in A \} = -\inf_{x \in A} I(x);$$

b) $\inf_{x \in E} I(x) = 0$.

Exercise 3.2. Let $E = \mathbb{R}$ and $\xi \sim N(0, 1)$. Show that the family $(\varepsilon\xi)_{\varepsilon>0}$ satisfies the LDP with rate function

$$I(x) = \begin{cases} +\infty & \text{if } x \neq 0, \\ 0 & \text{if } x = 0. \end{cases}$$

Compare this claim with the result of Remark 3.2.

Exercise 3.3. Let $a_n, b_n, n \geq 1$, be positive real numbers. Show that

$$\overline{\lim}_{n \rightarrow \infty} \frac{1}{n} \ln(a_n + b_n) = \overline{\lim}_{n \rightarrow \infty} \frac{1}{n} \ln a_n \vee \overline{\lim}_{n \rightarrow \infty} \frac{1}{n} \ln b_n,$$

where $a \vee b$ denotes the maximum of the set $\{a, b\}$.

Solution. We note that for every $n \geq 1$

$$0 \leq \ln(a_n + b_n) - \ln a_n \vee \ln b_n \leq \ln 2.$$

Dividing by n and taking $\overline{\lim}$, we obtain

$$\overline{\lim}_{n \rightarrow \infty} \frac{1}{n} \ln(a_n + b_n) = \overline{\lim}_{n \rightarrow \infty} \frac{1}{n} (\ln a_n \vee \ln b_n) = \overline{\lim}_{n \rightarrow \infty} \frac{1}{n} \ln a_n \vee \overline{\lim}_{n \rightarrow \infty} \frac{1}{n} \ln b_n.$$

Exercise 3.4. Let $\eta_1, \eta_2 \sim N(0, 1)$. Let also for every $\varepsilon > 0$ a random variable ξ_ε have the distribution defined as follows

$$\mathbb{P} \{ \xi_\varepsilon \in A \} = \frac{1}{2} \mathbb{P} \{ -1 + \sqrt{\varepsilon} \eta_1 \in A \} + \frac{1}{2} \mathbb{P} \{ 1 + \sqrt{\varepsilon} \eta_2 \in A \}$$

for all Borel sets A . Show that the family $(\xi_\varepsilon)_{\varepsilon>0}$ satisfies the LDP with rate function $I(x) = \frac{1}{2} \min \{ (x-1)^2, (x+1)^2 \}$, $x \in \mathbb{R}$.

(*Hint:* Show first that both families $(\sqrt{\varepsilon}\eta_1)_{\varepsilon>0}$ and $(\sqrt{\varepsilon}\eta_2)_{\varepsilon>0}$ satisfy LDP and find the corresponding rate functions. Then use Exercise 3.3)

4 LDP for empirical means

4.1 LDP for empirical means

In this section, we will assume that ξ_1, ξ_2, \dots be a sequence of independent identically distributed random variables in \mathbb{R} with $\mathbb{E} \xi_1 = \mu$. Similarly to Section 2, we consider the partial sums $S_n = \xi_1 + \dots + \xi_n$, $n \geq 1$, and show that the family of empirical means $(\frac{1}{n} S_n)_{n \geq 1}$ satisfies the LDP, using Cramer's theorem. As before we denote the cumulant generating function associated with ξ_1 by φ , and its Fenchel-Legendre transform by φ^* .

Proposition 4.1. *Under the assumption of Theorem 2.1, the family $(\frac{1}{n} S_n)_{n \geq 1}$ satisfies the large deviation principle in \mathbb{R} with rate function φ^* , that is, for every $A \in \mathcal{B}(\mathbb{R})$*

$$-\inf_{x \in A^\circ} \varphi^*(x) \leq \overline{\lim}_{n \rightarrow \infty} \frac{1}{n} \ln \mathbb{P} \left\{ \frac{1}{n} S_n \in A \right\} \leq \overline{\lim}_{n \rightarrow \infty} \frac{1}{n} \ln \mathbb{P} \left\{ \frac{1}{n} S_n \in A \right\} \leq -\inf_{x \in \bar{A}} \varphi^*(x).$$

Proof. In order to proof the proposition, we will show that inequalities (12) and (13) in Lemma 3.1 are satisfied.

Let F be a closed subset of \mathbb{R} . We first assume that $F \subseteq [\mu, +\infty)$. Since φ^* is convex and increasing on $[\mu, +\infty)$ (see Exercise 2.8), it is easily to seen that $\inf_{x \in F} \varphi^*(x) = \varphi^*(x_0)$, where $x_0 = \min F$. Then we can estimate

$$\overline{\lim}_{n \rightarrow \infty} \frac{1}{n} \ln \mathbb{P} \left\{ \frac{1}{n} S_n \in F \right\} \leq \overline{\lim}_{n \rightarrow \infty} \frac{1}{n} \ln \mathbb{P} \left\{ \frac{1}{n} S_n \geq x_0 \right\} = -\varphi^*(x_0) = -\inf_{x \in F} \varphi^*(x),$$

by Cramer Theorem 2.1. Similarly, we can prove the same result for $F \subseteq (-\infty, \mu]$. If $F \not\subseteq [\mu, +\infty)$ and $F \not\subseteq (-\infty, \mu]$, then we consider two closed sets $F_1 := F \cap [\mu, +\infty)$, $F_2 := F \cap (-\infty, \mu]$ and estimate

$$\begin{aligned} \overline{\lim}_{n \rightarrow \infty} \frac{1}{n} \ln \mathbb{P} \left\{ \frac{1}{n} S_n \in F \right\} &\leq \overline{\lim}_{n \rightarrow \infty} \frac{1}{n} \ln \left[\mathbb{P} \left\{ \frac{1}{n} S_n \in F_1 \right\} + \mathbb{P} \left\{ \frac{1}{n} S_n \in F_2 \right\} \right] \\ &= \overline{\lim}_{n \rightarrow \infty} \frac{1}{n} \ln \mathbb{P} \left\{ \frac{1}{n} S_n \in F_1 \right\} \vee \overline{\lim}_{n \rightarrow \infty} \frac{1}{n} \ln \mathbb{P} \left\{ \frac{1}{n} S_n \in F_2 \right\} \\ &\leq \left(-\inf_{x \in F_1} \varphi^*(x) \right) \vee \left(-\inf_{x \in F_2} \varphi^*(x) \right) = -\inf_{x \in F} \varphi^*(x), \end{aligned}$$

by Exercise 3.3.

In order to prove lower bound (13), we will use Proposition 3.1. We take $E_0 = \{x \in \mathbb{R} : \varphi^*(x) < +\infty\}$. By the continuity of φ^* , the set E_0 satisfies the properties of Proposition 3.1. Fix $x \in E_0$ such that $x > \mu$ and prove that

$$\lim_{r \rightarrow 0+} \overline{\lim}_{n \rightarrow \infty} \frac{1}{n} \ln \mathbb{P} \left\{ \frac{1}{n} S_n \in (x-r, x+r) \right\} \geq -\varphi^*(x). \quad (14)$$

We take $r > 0$ such that $x - \frac{r}{2} \geq \mu$ and note that $\varphi^*(x-r/2) < \varphi^*(x+r)$, according to Exercise 2.10 (v). Hence, by Exercise 4.1 below,

$$\begin{aligned} \overline{\lim}_{n \rightarrow \infty} \frac{1}{n} \ln \mathbb{P} \left\{ \frac{1}{n} S_n \in (x-r, x+r) \right\} &\geq \overline{\lim}_{n \rightarrow \infty} \frac{1}{n} \ln \mathbb{P} \left\{ \frac{1}{n} S_n \in \left[x - \frac{r}{2}, x+r \right) \right\} \\ &= \overline{\lim}_{n \rightarrow \infty} \frac{1}{n} \ln \left[\mathbb{P} \left\{ \frac{1}{n} S_n \geq x_0 - \frac{r}{2} \right\} - \mathbb{P} \left\{ \frac{1}{n} S_n \geq x_0 + r \right\} \right] = -\varphi^*(x_0 - r/2). \end{aligned} \quad (15)$$

Passing to the limit as $r \rightarrow 0+$, we obtain inequality (14). This finishes the proof of the proposition. If $x < \mu$, then similarly one can obtain (14) similarly. The case $x = \mu$ trivially follows from the law of large numbers and the fact that $\varphi^*(\mu) = 0$ (see Exercise 2.8 (ii)). \square

Exercise 4.1. Let $a_n > b_n$, $n \geq 1$, be positive real numbers such that there exist limits (probably infinite)

$$a := \lim_{n \rightarrow \infty} \frac{1}{n} \ln a_n \quad \text{and} \quad b := \lim_{n \rightarrow \infty} \frac{1}{n} \ln b_n$$

and $a > b$. Show that

$$\lim_{n \rightarrow \infty} \frac{1}{n} \ln(a_n - b_n) = a.$$

(Hint: Show that $\frac{b_n}{a_n} \rightarrow 0$, $n \rightarrow \infty$)

4.2 Multidimensional Cramer's theorem

In this section, we will state the LDP for empirical mean of random vectors. This result generalises Proposition 4.1.

Similarly to the one-dimensional case we introduce the **comulant generating function** associated with a random vector ξ in \mathbb{R}^d as follows

$$\varphi_\xi(\lambda) = \ln \mathbb{E} e^{\lambda \cdot \xi}, \quad \lambda \in \mathbb{R}^d,$$

where $a \cdot b = a_1 b_1 + \dots + a_d b_d$ for $a = (a_1, \dots, a_d)$ and $b = (b_1, \dots, b_d)$ from \mathbb{R}^d . As in one-dimensional case¹⁴, one can show that the function φ is convex. So, we can introduce the Fenchel-Legendre transform

$$\varphi_\xi^*(x) = \sup_{\lambda \in \mathbb{R}^d} \{\lambda \cdot x - \varphi(\lambda)\}, \quad x \in \mathbb{R}^d,$$

of a function φ .

Exercise 4.2. For any random vector $\xi \in \mathbb{R}^d$ and non-singular $d \times d$ matrix A , show that $\varphi_{A\xi}(\lambda) = \varphi_\xi(\lambda A)$ and $\varphi_{A\xi}^*(x) = \varphi_\xi^*(A^{-1}x)$.

Exercise 4.3. For any pair of independent random vectors ξ and η show that $\varphi_{\xi, \eta}(\lambda, \mu) = \varphi_\xi(\lambda) + \varphi_\eta(\mu)$ and $\varphi_{\xi, \eta}^*(x, y) = \varphi_\xi^*(x) + \varphi_\eta^*(y)$.

(Hint: To prove the second equality, use the equality $\sup_{\lambda, \mu} f(\lambda, \mu) = \sup_{\lambda} \sup_{\mu} f(\lambda, \mu)$)

The following theorem is multidimensional Cramer's theorem.

Theorem 4.1 (Cramer). *Let ξ_1, ξ_2, \dots be a sequence of independent identically distributed random vectors in \mathbb{R}^d with comulant generating function φ and let $S_n = \xi_1 + \dots + \xi_n$. If φ is finite in a neighborhood of 0 then the family $(\frac{1}{n}S_n)_{n \geq 1}$ satisfies the large deviation principle with good rate function φ^* , that is, for every Borel set $A \subset \mathbb{R}^d$*

$$-\inf_{x \in A^\circ} \varphi^*(x) \leq \liminf_{n \rightarrow \infty} \frac{1}{n} \ln \mathbb{P} \left\{ \frac{1}{n} S_n \in A \right\} \leq \overline{\lim}_{n \rightarrow \infty} \frac{1}{n} \ln \mathbb{P} \left\{ \frac{1}{n} S_n \in A \right\} \leq -\inf_{x \in \bar{A}} \varphi^*(x).$$

For proof of Theorem 4.1 see e.g. [RAS15, P.61] (for simpler proof in the case $\varphi(\lambda) < \infty$, $\lambda \in \mathbb{R}^d$, see e.g. [Var84, Theorem 3.1], [Kal02, Theorem 27.5] or [DZ98, Theorem 2.2.30].

Exercise 4.4. Let ξ_1, ξ_2, \dots be independent random vectors in \mathbb{R}^d whose coordinates are independent exponentially distributed random variables with rate γ .¹⁵ Show that the empirical means $(\frac{1}{n}S_n)_{n \geq 1}$ satisfies the LDP in \mathbb{R}^d and find the corresponding rate function I .

(Hint: Use Proposition 4.1. For computation of the rate function use exercises 2.7 and 4.3)

5 Lower semi-continuity and goodness of rate functions

5.1 Lower semi-continuity of rate functions

Let $(\xi_\varepsilon)_{\varepsilon > 0}$ satisfy the LDP in a metric space with rate function $I : E \rightarrow [0, \infty]$. In this section, we are going to answer the question when the rate function I is unique.

¹⁴see Exercise 2.1

¹⁵see also Example 2.2

Example 5.1. Let $\xi \sim N(0, 1)$. We know that the family $(\xi_\varepsilon := \sqrt{\varepsilon}\xi)_{\varepsilon>0}$ satisfies the LDP in \mathbb{R} with the good rate function $I(x) = \frac{x^2}{2}$.¹⁶ We take another function

$$\tilde{I}(x) = \begin{cases} \frac{x^2}{2} & \text{if } x \neq 0, \\ +\infty & \text{if } x = 0, \end{cases}$$

and show that the family $(\xi_\varepsilon)_{\varepsilon>0}$ also satisfies the LDP with rate function \tilde{I} .

Indeed, if G is an open set in \mathbb{R} , then trivially

$$\inf_{x \in G} I(x) = \inf_{x \in G} \tilde{I}(x).$$

For a closed F such that $F \neq \{0\}$ we also have

$$\inf_{x \in F} I(x) = \inf_{x \in F} \tilde{I}(x).$$

We have only to check the upper bound for the case $F = \{0\}$. We compute

$$\overline{\lim}_{\varepsilon \rightarrow 0} \varepsilon \ln \mathbb{P} \{ \xi_\varepsilon \in F \} = \overline{\lim}_{\varepsilon \rightarrow 0} \varepsilon \ln (\mathbb{P} \{ \sqrt{\varepsilon}\xi = 0 \}) = -\infty = - \inf_{x \in F} \tilde{I}(x).$$

This example shows that the same family of random variables can satisfy the LDP with different rate functions. In the rest of the section, we will impose some additional conditions on rate functions to provide the uniqueness.

Definition 5.1. A function $f : E \rightarrow [-\infty, +\infty]$ is called lower semi-continuous if

$$\underline{\lim}_{n \rightarrow \infty} f(x_n) \geq f(x) \quad \text{whenever } x_n \rightarrow x.$$

Remark 5.1. Note that the function \tilde{I} from Example 5.1 is not lower semi-continuous. Indeed, the inequality from Definition 5.1 does not hold e.g. for $x_n = \frac{1}{n}$.

Lemma 5.1. A function $f : E \rightarrow [-\infty, +\infty]$ is lower semi-continuous iff for each $\alpha \in [-\infty, +\infty]$ the level set $\{x \in E : f(x) \leq \alpha\}$ is a closed subset of E .

Proof. We assume that f is lower semi-continuous and show that for every $\alpha \in [-\infty, +\infty]$ the level set $L_\alpha = \{x \in E : f(x) \leq \alpha\}$ is closed. If $\alpha = +\infty$, then the closedness L_α is trivial due to $L_\alpha = E$. Let $\alpha < +\infty$. We assume that L_α is not close. Then there exists a convergent sequence $x_n \in L_\alpha$, $n \geq 1$, such that $x_n \rightarrow x$ and $x \notin L_\alpha$. Using the lower semi-continuity of f and the fact that $x_n \in L_\alpha$, $n \geq 1$, we get

$$\alpha < f(x) \leq \underline{\lim}_{n \rightarrow \infty} f(x_n) \leq \alpha.$$

This gives the contradiction.

Now, let L_α be closed for every $\alpha \in [-\infty, +\infty]$. Assume that f is not lower semi-continuous. Then there exists a sequence $x_n \rightarrow x$ such that

$$\underline{\lim}_{n \rightarrow \infty} f(x_n) < f(x).$$

¹⁶see Remark 3.2

We take α such that $\underline{\lim}_{n \rightarrow \infty} f(x_n) < \alpha < f(x)$. Then there exists a subsequence $\{x_{n_k}\}_{k \geq 1}$ of $\{x_n\}_{n \geq 1}$ such that

$$\lim_{k \rightarrow \infty} f(x_{n_k}) = \underline{\lim}_{n \rightarrow \infty} f(x_n) < \alpha$$

and $f(x_{n_k}) < \alpha$. Hence, $x_{n_k} \in L_\alpha$, $k \geq 1$. Since $x_{n_k} \rightarrow x$ as $k \rightarrow \infty$, the closability of L_α implies that $x \in L_\alpha$. Therefore, $f(x) \leq \alpha$. We have obtained the contradiction. \square

Let $C_0[0, T]$ denote the Banach space of continuous functions from $[0, T]$ satisfying $f(0) = 0$ endowed with the uniform norm.¹⁷ Let $H_0^2[0, T]$ be the set of all absolutely continuous¹⁸ functions $f \in C_0[0, T]$ with $\dot{f} \in L_2[0, T]$.

Exercise 5.1. Let $f \in C_0^1[0, T]$.¹⁹ Show that f is absolutely continuous and \dot{f} coincides with the classical derivative f' of f . Conclude that $f \in H_0^2[0, T]$.

Exercise 5.2. Show that the function $f(x) = 1 - |x - 1|$, $x \in [0, 2]$, belongs to $H_0^2[0, 2]$ but is not continuously differentiable.

(Hint: Show that $\dot{f}(x) = \begin{cases} 1 & \text{if } x \in [0, 1], \\ -1 & \text{if } x \in (1, 2]. \end{cases}$)

We consider a function from $C_0[0, T]$ to $[0, +\infty]$ defined as follows

$$I(f) = \begin{cases} \frac{1}{2} \int_0^T \dot{f}^2(x) dx & \text{if } f \in H_0^2[0, T], \\ +\infty & \text{otherwise.} \end{cases} \quad (17)$$

Exercise 5.3. Let $I : C_0[0, T] \rightarrow [0, +\infty]$ be defined by (17). Show that I is lower semi-continuous. (Hint: Use Lemma 5.1 and the Banach-Alaoglu theorem)

Next we are going to show that one can always replace a rate function by a lower semi-continuous rate function. Moreover, it turns out that a lower semi-continuous rate function is unique. For this, we introduce a transformation produces a lower semi-continuous function f_{lsc} from an arbitrary function $f : E \rightarrow [-\infty, +\infty]$ (for more details see [RAS15, Section 2.2]).

The **lower semi-continuous regularization** of f is defined by

$$f_{\text{lsc}}(x) = \sup \left\{ \inf_{y \in G} f(y) : G \ni x \text{ and } G \text{ is open} \right\}. \quad (18)$$

Exercise 5.4. Show that the function \tilde{I}_{lsc} coincides with $I(x) = \frac{x^2}{2}$, $x \in \mathbb{R}$, where \tilde{I} was defined in Example 5.1.

Exercise 5.5. Let $f(x) = \mathbb{I}_{\mathbb{Q}}(x)$, $x \in \mathbb{R}$, where \mathbb{Q} denotes the set of rational numbers. Find the function f_{lsc} .

¹⁷The uniform norm on $C_0[0, T]$ is defined as $\|f\|_C = \max_{x \in [0, T]} |f(x)|$. The space $C_0[0, T]$ endowed with this norm is a separable Banach space

¹⁸A continuous function $f \in C[0, T]$ (not necessarily $f(0) = 0$) is said to be **absolutely continuous** if there exists a function $h \in L_1[0, T]$ such that

$$f(t) = f(0) + \int_0^t h(s) ds, \quad t \in [0, T]. \quad (16)$$

Such a function h is denoted by \dot{f} and is called the derivative of f .

¹⁹ $C_0^m[0, T]$ consists of all functions from $C_0[0, T]$ which are m times continuously differentiable on $(0, T)$

Lemma 5.2. *The function f_{lsc} is lower semi-continuous and $f_{\text{lsc}}(x) \leq f(x)$ for all $x \in E$. If g is lower semi-continuous and satisfies $g(x) \leq f(x)$ for all x , then $g(x) \leq f_{\text{lsc}}(x)$ for all x . In particular, if f is lower semi-continuous, then $f = f_{\text{lsc}}$.*

The Lemma 5.2 says that the lower semi-continuous regularization f_{lsc} of f is the maximal lower semi-continuous function less or equal that f .

Proof of Lemma 5.2. The inequality $f_{\text{lsc}} \leq f$ is clear. To show that f_{lsc} is lower semi-continuous, we use Lemma 5.1. Let $x \in \{f_{\text{lsc}} > \alpha\}$. Then there is an open set G containing x such that $\inf_G f > \alpha$. Hence by the supremum in the definition of f_{lsc} , $f_{\text{lsc}}(y) \geq \inf_G f > \alpha$ for all $y \in G$. Thus G is an open neighborhood of x contained in $\{f_{\text{lsc}} > \alpha\}$. So $\{f_{\text{lsc}} > \alpha\}$ is open.

To show that $g \leq f_{\text{lsc}}$ one just needs to show that $g_{\text{lsc}} = g$. Indeed,

$$\begin{aligned} g(x) &= g_{\text{lsc}}(x) = \sup \left\{ \inf_G g : x \in G \text{ and } G \text{ is open} \right\} \\ &\leq \sup \left\{ \inf_G f : x \in G \text{ and } G \text{ is open} \right\} = f_{\text{lsc}}(x). \end{aligned}$$

We already know that $g_{\text{lsc}} \leq g$. To show the other direction let α be such that $g(x) > \alpha$. Then, $G = \{g > \alpha\}$ is an open set containing x and $\inf_G g \geq \alpha$. Thus, $g_{\text{lsc}}(x) \geq \alpha$. Now increasing α to $g(x)$, we obtain the needed inequality $g_{\text{lsc}}(x) \geq g(x)$. \square

Exercise 5.6. 1) Show that if $x_n \rightarrow x$, then $f_{\text{lsc}}(x) \leq \liminf_{n \rightarrow \infty} f(x_n)$.

(Hint: Use Lemma 5.2, namely that the function f_{lsc} is lower semi-continuous and $f_{\text{lsc}} \leq f$)

2) Show that for each the supremum in (18) can only be taken over all ball with center x , namely

$$f_{\text{lsc}}(x) = \sup_{r>0} \inf_{y \in B_r(x)} f(y) \tag{19}$$

(Hint: Use the fact that any open set G containing x also contains a ball $B_r(x)$ for some $r > 0$. It will allow to prove the inequality $f_{\text{lsc}}(x) \leq \sup_{r>0} \inf_{y \in B_r(x)} f(y)$. The inverse inequality just follows from the observation that supremum in the right hand side of (19) is taken over smaller family of open sets)

3) Prove that for each $x \in E$ there is a sequence $x_n \rightarrow x$ such that $f(x_n) \rightarrow f_{\text{lsc}}(x)$ (the constant sequence $x_n = x$ is allowed here). This gives the alternate definition

$$f_{\text{lsc}}(x) = \min \left\{ f(x), \liminf_{y \rightarrow x} f(y) \right\}.$$

(Hint: Use part 2) of the exercise to construct the corresponding sequence x_n , $n \geq 1$)

Proposition 5.1. *Let $(\xi_\varepsilon)_{\varepsilon>0}$ satisfy the LDP in a metric space E with rate function I . Then it satisfies the LDP in E with the rate function I_{lsc} . Moreover, there exists only unique lower semi-continuous associated rate function.*

Proof. We first show that $(\xi_\varepsilon)_{\varepsilon>0}$ satisfies the LDP in E with the lower semi-continuous function I_{lsc} . For this we check the inequalities of Lemma 3.1. We note that the upper bound immediately follows from the inequality $I_{\text{lsc}} \leq I$ (see Lemma 5.2). For the lower bound we observe that $\inf_G I_{\text{lsc}} = \inf_G I$ when G is open. Indeed, the inequality $\inf_G I_{\text{lsc}} \leq \inf_G I$ follows from $I_{\text{lsc}} \leq I$. In order to prove inverse inequality, we will use the definition of lower semi-continuous regularization. Remark that for every $x \in G$ one has $f_{\text{lsc}}(x) \geq \inf_G I$. Hence $\inf_G I_{\text{lsc}} \geq \inf_G I$.

To prove the uniqueness, assume that (11) holds for two lower semi-continuous functions I and J , and let $I(x) < J(x)$ for some $x \in E$. By the lower semi-continuity of J , we may choose a neighborhood G of x such that $\inf_G J > I(x)$, taking e.g. G as an open neighborhood of x such that $\bar{G} \subset \left\{ y : J(y) > I(x) + \frac{J(x)-I(x)}{2} \right\}$. Then applying (11) to both I and J yields the contradiction

$$-I(x) \leq -\inf_G I \leq \liminf_{\varepsilon \rightarrow 0} \varepsilon \ln \mathbb{P} \{ \xi_\varepsilon \in G \} \leq -\inf_G J < -I(x).$$

We obtained the contradiction with the assumption $I(x) < J(x)$. □

Exercise 5.7. Assume φ^* that is the Fenchel-Legendre transform of the cumulant generating function. Show that φ^* is lower semi-continuous.

(Hint: Show that supremum of a family of continuous functions is lower semi-continuous)

5.2 Goodness of rate functions

We remark that in many cases the rate function satisfies better properties than lower semi-continuity.

Definition 5.2. We say that a rate function $I : E \rightarrow [0, +\infty]$ is **good** if the level sets $\{x \in E : I(x) \leq \alpha\}$ are compact (rather than just closed) for all $\alpha \geq 0$.

Example 5.2. Show that the rate function $I(x) = \frac{\|x\|^2}{2}$, $x \in \mathbb{R}^d$, from Exercise 1.3 is good.

Remark 5.2. The rate functions from all previous examples are also good.

Now we consider another example of a good rate function which is the rate function for LDP for Brownian motion. We obtain the LDP for Brownian motion later and here we just show that the associated rate function is good.

Exercise 5.8. Let $I : C_0[0, T] \rightarrow [0, +\infty]$ be defined by (17). Show that the set $\{f \in C_0[0, T] : I(f) \leq \alpha\}$ is equicontinuous²⁰ and bounded in $C_0[0, T]$ for all $\alpha \geq 0$. Conclude that I is good.

(Hint: Using Hölder's inequality, show that $|f(t) - f(s)|^2 \leq |t - s| \int_0^T \dot{f}^2(x) dx$ for all $t, s \in [0, T]$ and each $f \in H_0^2[0, T]$)

6 Weak large deviation principle and exponential tightness

6.1 Weak large deviation principle

Proposition 3.1 shows that lower bound inequality (13) is enough to show only for open balls. Unfortunately, it is not enough for upper bound (12). Later, in Proposition 6.1, we will show that upper bound (12) for closed (or open) balls will only imply the upper bound for compact sets F . To have the

²⁰see Definition VI.3.7 [Con90]

upper bound for any closed set we need one extra condition, called exponential tightness, which we will discuss in the next section. Let us consider the following example taken from [DZ98, P. 7] which demonstrates that upper bound for all compact sets does not imply inequality (12) for any closed set.

Example 6.1. We consider random variables $\xi_\varepsilon := \frac{1}{\varepsilon}$, $\varepsilon > 0$, in \mathbb{R} and set $I(x) := +\infty$, $x \in \mathbb{R}$. Then for any compact set F in \mathbb{R} (which is also bounded) we have

$$\overline{\lim}_{\varepsilon \rightarrow 0} \varepsilon \ln \mathbb{P} \{ \xi_\varepsilon \in F \} = -\infty = - \inf_{x \in F} I(x)$$

because there exists $\varepsilon_0 > 0$ such that $\mathbb{P} \{ \xi_\varepsilon \in F \} = 0$ for all $\varepsilon \in (0, \varepsilon_0)$. But it is easily seen that this inequality is not preserved for the closed set $F = \mathbb{R}$. Indeed,

$$\overline{\lim}_{\varepsilon \rightarrow 0} \varepsilon \ln \mathbb{P} \{ \xi_\varepsilon \in \mathbb{R} \} = \overline{\lim}_{\varepsilon \rightarrow 0} \varepsilon \ln 1 = 0 \not\leq -\infty = \inf_{x \in \mathbb{R}} I(x).$$

We also remark here that the family $(\xi_\varepsilon)_{\varepsilon > 0}$ and the function I satisfy lower bound (13).

Consequently, it makes sense to introduce a relaxation of the *full* LDP, where we will require the upper bound only for compact sets.

Definition 6.1. We say that the family $(\xi_\varepsilon)_{\varepsilon > 0}$ satisfies the **weak large deviation principle (weak LDP) in E with rate function I** if

$$\overline{\lim}_{\varepsilon \rightarrow 0} \varepsilon \ln \mathbb{P} \{ \xi_\varepsilon \in F \} \leq - \inf_{x \in F} I(x) \tag{20}$$

for every *compact* set $F \subset E$, and


$$\underline{\lim}_{\varepsilon \rightarrow 0} \varepsilon \ln \mathbb{P} \{ \xi_\varepsilon \in G \} \geq - \inf_{x \in G} I(x) \tag{21}$$

for every open set $G \subset E$.

We remark that Example 6.1 shows that the family $(\xi_\varepsilon = \frac{1}{\varepsilon})_{\varepsilon > 0}$ satisfies the weak LDP in \mathbb{R} with good rate function $I(x) = +\infty$, $x \in \mathbb{R}$, but it does not satisfy the full LDP.

Let us consider another interesting example of a family of random elements in $C[0, T]$ which satisfies the weak LDP but it does not satisfies the full LDP for any rate function. This is a recent result obtained by V. Kuznetsov in [Kuz15].

Example 6.2 (Winding angle of Brownian trajectory around the origin). Let $w(t) = (w_1(t), w_2(t))$, $t \in [0, T]$, be a two dimensional Brownian motion started from the point $(1, 0)$. We denote for every $t \in [0, T]$ the angle between the vector $w(t)$ and the x -axis (the vector $(1, 0)$) by $\Phi(t)$ and set $\Phi_\varepsilon(t) = \Phi(\varepsilon t)$, $t \in [0, T]$. It turns out that the family $(\Phi_\varepsilon)_{\varepsilon > 0}$ satisfies only the weak LDP in the space of continuous functions $C[0, T]$.



Winding angle of Brownian motion

In the next section, we will consider conditions on a family $(\xi_\varepsilon)_{\varepsilon>0}$ which will guarantee the implication. Now we will give a useful statement which allows to check the upper bound for the weak LDP.

Proposition 6.1. *Let $(\xi_\varepsilon)_{\varepsilon>0}$ be a family of random variables in a metric space E and let I be a function from E to $[0, +\infty]$. Then upper bound (20) follows from*

$$\lim_{r \rightarrow 0} \overline{\lim}_{\varepsilon \rightarrow 0} \varepsilon \ln \mathbb{P} \{ \xi_\varepsilon \in B_r(x) \} \leq -I(x)$$

for all $x \in E$.

Proof. Let F be a compact set. We set $\alpha := \inf_{x \in F} I(x)$ and assume that $\alpha < \infty$. Remark that for every $x \in F$ $I(x) \geq \inf_{x \in F} I(x) = \alpha$. Hence, for any fixed $\delta > 0$

$$\lim_{r \rightarrow 0} \overline{\lim}_{\varepsilon \rightarrow 0} \varepsilon \ln \mathbb{P} \{ \xi_\varepsilon \in B_r(x) \} \leq -I(x) \leq -\alpha < -\alpha + \delta$$

for all $x \in F$. Consequently, by the definition of limit, for every $x \in F$ there exists $r_x > 0$ such that

$$\overline{\lim}_{\varepsilon \rightarrow 0} \varepsilon \ln \mathbb{P} \{ \xi_\varepsilon \in B_{r_x}(x) \} < -\alpha + \delta.$$

Since the family of balls $B_{r_x}(x)$, $x \in F$, is an open cover²¹ of F . By the compactness of F , there exists a finite subcover of F , i.e. there exist $x_1, \dots, x_m \in F$ such that $F \subset \bigcup_{k=1}^m B_{r_{x_k}}(x_k)$. Now we can estimate

$$\begin{aligned} \overline{\lim}_{\varepsilon \rightarrow 0} \varepsilon \ln \mathbb{P} \{ \xi_\varepsilon \in F \} &\leq \overline{\lim}_{\varepsilon \rightarrow 0} \varepsilon \ln \mathbb{P} \left\{ \xi_\varepsilon \in \bigcup_{k=1}^m B_{r_{x_k}}(x_k) \right\} \leq \overline{\lim}_{\varepsilon \rightarrow 0} \varepsilon \ln \left(\sum_{k=1}^m \mathbb{P} \{ \xi_\varepsilon \in B_{r_{x_k}}(x_k) \} \right) \\ &\stackrel{\text{Exercise 3.3}}{=} \max_{k=1, \dots, m} \overline{\lim}_{\varepsilon \rightarrow 0} \varepsilon \ln \mathbb{P} \{ \xi_\varepsilon \in B_{r_{x_k}}(x_k) \} < -\alpha + \delta = -\inf_{x \in F} I(x) + \delta. \end{aligned}$$

Making $\delta \rightarrow 0$, we obtain

$$\overline{\lim}_{\varepsilon \rightarrow 0} \varepsilon \ln \mathbb{P} \{ \xi_\varepsilon \in F \} \leq -\inf_{x \in F} I(x).$$

Similarly, one can show inequality (20) in the case $\alpha = +\infty$ replacing $-\alpha + \delta$ by $-\frac{1}{\delta}$. □

Exercise 6.1. Finish the proof of Proposition 6.1 in the case $\inf_{x \in F} I(x) = +\infty$.

²¹Each set $B_{r_x}(x)$ is open and $F \subset \bigcup_{x \in F} B_{r_x}(x)$

6.2 Exponential tightness

We start from the definition of exponential tightness.

Definition 6.2. A family of random elements $(\xi_\varepsilon)_{\varepsilon>0}$ is said to be **exponentially tight in E** if for any number $\beta > 0$ there exists a compact set $K \subset E$ such that

$$\overline{\lim}_{\varepsilon \rightarrow 0} \varepsilon \ln \mathbb{P} \{ \xi_\varepsilon \notin K \} \leq -\beta. \quad (22)$$

We remark that in the case of a countable family of random elements $(\xi_n)_{n \geq 1}$, the exponential tightness corresponds to the statement

$$\overline{\lim}_{n \rightarrow \infty} a_n \ln \mathbb{P} \{ \xi_n \notin K \} \leq -\beta \quad (23)$$

for some $a_n \rightarrow 0$.

Exercise 6.2. Prove that a family $(\xi_\varepsilon)_{\varepsilon>0}$ is exponentially tight in E if and only if for any $b > 0$ there exists a compact $K \subset E$ and $\varepsilon_0 > 0$ such that

$$\mathbb{P} \{ \xi_\varepsilon \notin K \} \leq e^{-\frac{1}{\varepsilon}b}, \quad \varepsilon \in (0, \varepsilon_0).$$

Exercise 6.2 shows that the exponential tightness is much more stronger than the tightness²².

Exercise 6.3. Let E be a complete and separable metric space.

a) Show that exponential tightness implies tightness for a countable family of random variables.

(Hint: Prove a similar inequality to one in Exercise 6.2 and then use the fact that any random element on a complete and separable metric space is tight (see Lemma 3.2.1 [EK86])

b) Show that tightness does not imply exponential tightness.

Example 6.3. Let ξ be a standard Gaussian vector in \mathbb{R}^d . We consider as before $\xi_\varepsilon = \sqrt{\varepsilon}\xi$ and check the family $(\xi_\varepsilon)_{\varepsilon>0}$ is exponentially tight in \mathbb{R}^d . For this we will use the fact that this family satisfies the LDP.

So, we fix $\beta > 0$ and take a compact set $K_a := [-a, a]^d$ such that $\inf_{\mathbb{R}^d \setminus (-a, a)^d} I \geq \beta$, where $I(x) = \frac{\|x\|^2}{2}$, $x \in \mathbb{R}^d$, is the rate function for the family $(\xi_\varepsilon)_{\varepsilon>0}$ ²³. Since the family $(\xi_\varepsilon)_{\varepsilon>0}$ satisfies the LDP with rate function I , we have

$$\overline{\lim}_{\varepsilon \rightarrow 0} \varepsilon \ln \mathbb{P} \{ \xi_\varepsilon \notin K_a \} \leq \overline{\lim}_{\varepsilon \rightarrow 0} \varepsilon \ln \mathbb{P} \left\{ \xi_\varepsilon \in \mathbb{R}^d \setminus (-a, a)^d \right\} \leq - \inf_{\mathbb{R}^d \setminus (-a, a)^d} I \leq -\beta.$$

Exercise 6.4. Let $(\xi_\varepsilon)_{\varepsilon>0}$ be a family of random variables in \mathbb{R} such that there exist $\lambda > 0$ and $\kappa > 0$ such that $\mathbb{E} e^{\frac{\lambda}{\varepsilon}|\xi_\varepsilon|} \leq \kappa \frac{1}{\varepsilon}$ for all $\varepsilon > 0$. Show that this family is exponentially tight.

(Hint: Use Chebyshev's inequality)

Proposition 6.2. *If a family $(\xi_\varepsilon)_{\varepsilon>0}$ is exponentially tight and satisfies a weak LDP in E with rate function I , then it satisfies a full LDP. Moreover, if I is lower semi-continuous, then I is good.*

²²A family of random variables (ξ_ε) is **tight** if for any $\delta > 0$ there exists a compact set $K \subset E$ such that $\mathbb{P} \{ \xi_\varepsilon \notin K \} \leq \delta$ for all ε

²³see Exercise 3.2

Proof. In order to prove a full LDP, we need to stay only upper bound (12) for each closed set $F \subset E$. So, let F be a given closed set and K and $\beta > 0$ be the corresponding set and constant from the definition of exponential tightness (see (22)). Then, using properties of probability, we have

$$\mathbb{P} \{ \xi_\varepsilon \in F \} \leq \mathbb{P} \{ \xi_\varepsilon \in F \cap K \} + \mathbb{P} \{ \xi_\varepsilon \in K^c \},$$

where $K = E \setminus K$ is the complement of K . Consequently, using Exercise 3.3 and the fact that the set $K \cap F$ is compact²⁴, one can estimate

$$\begin{aligned} \overline{\lim}_{\varepsilon \rightarrow 0} \varepsilon \ln \mathbb{P} \{ \xi_\varepsilon \in F \} &\leq \overline{\lim}_{\varepsilon \rightarrow 0} \varepsilon \ln (\mathbb{P} \{ \xi_\varepsilon \in F \cap K \} + \mathbb{P} \{ \xi_\varepsilon \in K^c \}) \\ &= \overline{\lim}_{\varepsilon \rightarrow 0} \varepsilon \ln \mathbb{P} \{ \xi_\varepsilon \in F \cap K \} \vee \overline{\lim}_{\varepsilon \rightarrow 0} \varepsilon \ln \mathbb{P} \{ \xi \notin K \} \\ &\leq \left(- \inf_{x \in F \cap K} I(x) \right) \vee (-\beta) \leq \left(- \inf_{x \in F} I(x) \right) \vee (-\beta). \end{aligned}$$

Letting $\beta \rightarrow +\infty$, we get upper bound (12).

We assume that I is lower semi-continuous. We fix $\alpha \geq 0$ and show that the level set $\{x \in E : I(x) \leq \alpha\}$ is compact. Let $K \subset E$ be the compact set from (22) in Definition 6.2 with $\beta = \alpha + 1$. Applying the lower bound of the definition of LDP to the open set K^c , we obtain

$$- \inf_{x \in K^c} I(x) \leq \underline{\lim}_{\varepsilon \rightarrow 0} \varepsilon \ln \mathbb{P} \{ \xi_\varepsilon \in K^c \} \leq -\beta < -\alpha.$$

Therefore, the closed set $\{x \in E : I(x) \leq \alpha\}$ is a subset of the compact set K , that implies that it is compact itself. \square

It turns out, that the full LDP implies exponential tightness, but only for a countable family $(\xi_\varepsilon)_{n \geq 1}$.

Proposition 6.3. *Let E be a complete and separable metric space. If $(\xi_n)_{n \geq 1}$ satisfies the full LDP with a good rate function I . Then it is exponentially tight.*

Proof. Since E is separable, for any $k \in \mathbb{N}$ we may cover E by some open balls B_{k1}, B_{k2}, \dots of radius $1/k$. We put $U_{km} = \bigcup_{j=1}^m B_{kj}$. We fix an arbitrary $\beta > 0$ and $k \in \mathbb{N}$ and show that there exists $m_k \geq 1$ such that for every $n \geq 1$

$$\mathbb{P} \{ \xi_n \notin U_{km_k} \} \leq \varepsilon^{-\frac{\beta k}{a_n}}. \quad (24)$$

We first remark that the level set $L_{\beta k} := \{x \in E : I(x) \leq \beta k + 1\}$ is compact due to the goodness of I . Since B_{k1}, B_{k2}, \dots is an open cover of $L_{\beta k}$, there exists a finite subcover $B_{kj_1}, \dots, B_{kj_l}$ of $L_{\beta k}$, where $j_1 < \dots < j_l$. Then trivially $L_{\beta k} \subseteq U_{k,j_l}$. By upper bound for the LDP,

$$\overline{\lim}_{n \rightarrow \infty} a_n \ln \mathbb{P} \{ \xi_n \in U_{k,j_l}^c \} \leq - \inf_{U_{k,j_l}^c} I \leq - \inf_{L_{\beta k}^c} I \leq -\beta k - 1.$$

According to the definition of the upper limit, we can choose $N \geq 1$ such that for every $n \geq N$

$$a_n \ln \mathbb{P} \{ \xi_n \in U_{k,j_l}^c \} \leq -\beta k$$

or equivalently

$$\mathbb{P} \{ \xi_n \notin U_{k,j_l} \} \leq e^{-\frac{\beta k}{a_n}}.$$

²⁴since $K \cap F$ is a closed subset of the compact set K

Next, using the fact that $\mathbb{P} \{ \xi_n \notin U_{k,m} \} \rightarrow 0$ as $m \rightarrow \infty$, we can find $m_k \geq j_l$ such that inequality (24) for every $1 \leq n < N$. Moreover, for every $n \geq N$

$$\mathbb{P} \{ \xi_n \notin U_{k,m_k} \} \leq \mathbb{P} \{ \xi_n \notin U_{k,j_l} \} \leq e^{-\frac{\beta k}{a_n}}.$$

This proves (24) for every $n \geq 1$.

Summing (24) over k , we obtain

$$\mathbb{P} \left\{ \xi_n \notin \bigcap_{k=1}^{\infty} U_{k,m_k} \right\} \leq \sum_{k=1}^{\infty} \mathbb{P} \{ \xi_n \notin U_{k,m_k} \} \leq \sum_{k=1}^{\infty} e^{-\frac{\beta k}{a_n}} = \frac{e^{-\beta/a_n}}{1 - e^{-\beta/a_n}}.$$

Let K be the closure of the set $\bigcap_{k=1}^{\infty} U_{k,m_k}$, which is compact, since $\bigcap_{k=1}^{\infty} U_{k,m_k}$ is totally bounded.²⁵ Therefore,

$$\overline{\lim}_{n \rightarrow \infty} a_n \ln \mathbb{P} \{ \xi_n \notin K \} \leq \overline{\lim}_{n \rightarrow \infty} a_n \ln \mathbb{P} \left\{ \xi_n \notin \bigcap_{k=1}^{\infty} U_{k,m_k} \right\} \leq -\beta.$$

This finishes the proof of the proposition. □

Exercise 6.5. Find a simpler proof of Proposition 6.3 in the case $E = \mathbb{R}^d$.

(Hint: Cover a level set $\{x \in \mathbb{R}^d : I(x) \leq \beta\}$ by an open ball and use the upper bound)

7 Large deviation principle for Brownian motion

7.1 Schilder's theorem

We start this section with computation of the rate function for finite dimensional distributions of a Brownian motion. So, let $w(t)$, $t \in [0, T]$, denote a standard Brownian motion on \mathbb{R} .²⁶ We take a partition $0 = t_0 < t_1 < \dots < t_d = T$ and consider the random vector $\xi = (w(t_1), \dots, w(t_d))$ in \mathbb{R}^d . Let ξ_1, ξ_2, \dots be independent copies of ξ . Then the distribution of

$$\frac{1}{n} S_n = \frac{1}{n} \sum_{k=1}^n \xi_k$$

coincides with the distribution of $\frac{1}{\sqrt{n}}(w(t_1), \dots, w(t_d))$. Consequently, one can use Theorem 4.1 to conclude that the family $\left(\frac{1}{\sqrt{n}}(w(t_1), \dots, w(t_d)) \right)_{n \geq 1}$ satisfies the LDP with good rate function φ_{ξ}^* . Next we compute φ_{ξ}^* to see the precise form of the rate function. We remark that the random vector

$$\eta = \left(\frac{w(t_1)}{\sqrt{t_1}}, \frac{w(t_2) - w(t_1)}{\sqrt{t_2 - t_1}}, \dots, \frac{w(t_d) - w(t_{d-1})}{\sqrt{t_d - t_{d-1}}} \right)$$

is a standard Gaussian vector in \mathbb{R}^d . According to exercises 4.3 and 2.7,

$$\varphi_{\eta}^*(x) = \frac{\|x\|^2}{2}, \quad x \in \mathbb{R}^d.$$

²⁵A set A is totally bounded in a metric space E if for every $r > 0$ it can be covered by a finite number of balls of radius r (see e.g. Definition I.14 [DS88]). By Theorem I.15 [DS88], the closure of a totally bounded set is compact in E , if E is complete.

²⁶ $w(t)$, $t \in [0, T]$ is a Brownian motion with $w(0) = 0$ and $\text{Var } w(t) = t$

We observe that

$$\eta = A \begin{pmatrix} w(t_1) \\ \dots \\ w(t_d) \end{pmatrix},$$

where A is some non-singular $d \times d$ -matrix. Thus, by Exercise 4.2,

$$\varphi_\xi^*(x) = \varphi_{A^{-1}\eta}^*(x) = \varphi_\eta^*(Ax) = \frac{\|Ax\|^2}{2} = \frac{1}{2} \sum_{k=1}^n \frac{(x_k - x_{k-1})^2}{t_k - t_{k-1}},$$

where $x_0 = 0$.

Let us denote

$$w_\varepsilon(t) = \sqrt{\varepsilon}w(t), \quad t \in [0, T].$$

Then taking a function $f \in C[0, T]$, we should expect that the family $(w_\varepsilon)_{\varepsilon>0}$ will satisfy the LDP in $C[0, T]$ with rate function

$$I(f) = \frac{1}{2} \int_0^T f'^2(t) dt.$$

Now we give a rigorous statement about the LDP for a Brownian motion. So, let $H_0^2[0, T]$ be a set of all absolutely continuous functions $h \in C_0[0, T]$ with $\dot{h} \in L_2[0, T]$ (see also Section 5 for more details).

Theorem 7.1 (Schilder's theorem). *The family $(w_\varepsilon)_{\varepsilon>0}$ satisfies the large deviation principle in $C_0[0, T]$ with good rate function*

$$I(f) = \begin{cases} \frac{1}{2} \int_0^T \dot{f}^2(t) dt & \text{if } f \in H_0^2[0, T], \\ +\infty & \text{otherwise.} \end{cases}$$

In order to prove Schilder's theorem, we are going estimate probabilities $\mathbb{P} \{w_\varepsilon \in B_r(f)\}$, where $B_r(f) = \{g \in C[0, T] : \|g - f\|_C < r\}$ is the ball in $C[0, T]$ with center f and radius r . This will be enough to prove the weak LDP according to Proposition 6.1 and Proposition 3.1. Then we will prove the exponential tightness of $(w_\varepsilon)_{\varepsilon>0}$ that will guarantee the full LDP by Proposition 6.2. The fact that the rate function I is good was considered as an exercise (see Exercise 5.8 above).

Exercise 7.1. Let $N(t)$, $t \geq 0$, be a Poisson process. Define $N_n(t) = \frac{1}{n}N(nt)$, $t \geq 0$, for all $n \geq 1$.

1. Show that for every $t > 0$ the family $(N_n(t))_{n \geq 1}$ satisfies the LDP in \mathbb{R} (with $a_n = \frac{1}{n}$) and find the corresponding rate function.
2. Show that for every $t_1 < t_2 < \dots < t_d$ the family $((N_n(t_1), \dots, N_n(t_d)))_{n \geq 1}$ satisfies the LDP in \mathbb{R}^d (with $a_n = \frac{1}{n}$) and find the corresponding rate function.
3. Which form should have the rate function in the LDP for the family of processes $\{N_n(t), t \in [0, T]\}_{n \geq 1}$ in the space $C_0[0, T]$?²⁷

(Hint: Express $N_n(t)$, $t \geq 0$, as the empirical mean of independent copies of the Poisson process $N(t)$, $t \geq 0$. Then use Cramer's theorem (see Proposition 4.1 and Theorem 4.1) for 1. and 2. For the computation of the Fenchel-Legendre transform in 2., use the same approach as in this section)

²⁷The proof of LDP for processes with independent increments (in particular, for a Poisson process) can be found e.g. in [LS87]

7.2 Cameron-Martin formula

In order to estimate the probability $\mathbb{P}\{w_\varepsilon \in B_r(f)\} = \mathbb{P}\{\|w_\varepsilon - f\|_C < r\}$, we need to work with distribution of the process $w_\varepsilon(t) - f(t)$, $t \in [0, T]$. We start the section with a simple observation. Let a random variable $\eta \sim N(0, 1)$ be given on a probability space $(\Omega, \mathcal{F}, \mathbb{P})$ and $a \in \mathbb{R}$. It turns out that one can change the probability measure \mathbb{P} in such a way that the random variable $\eta - a$ has a standard normal distributed. We note that

$$e^{-\frac{(x-a)^2}{2}} = e^{ax - \frac{a^2}{2}} e^{-\frac{x^2}{2}}.$$

Considering the new probability measure on Ω defined as

$$\mathbb{P}^a \{A\} = \mathbb{E} \mathbb{I}_A e^{a\eta - \frac{a^2}{2}}, \quad A \in \mathcal{F},$$

we claim that the random variable $\eta - a$ has a standard normal distribution on $(\Omega, \mathcal{F}, \mathbb{P}^a)$.

Exercise 7.2. Show that \mathbb{P}^a is a probability measure on Ω , i.e. $\mathbb{P}^a \{\Omega\} = 1$.

Indeed, for any $z \in \mathbb{R}$ we have

$$\begin{aligned} \mathbb{P}^a \{\eta - a \leq z\} &= \mathbb{E} \mathbb{I}_{\{\eta - a \leq z\}} e^{a\eta - \frac{a^2}{2}} = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{z+a} e^{ax - \frac{a^2}{2}} e^{-\frac{x^2}{2}} dx \\ &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{z+a} e^{-\frac{(x-a)^2}{2}} dx = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^z e^{-\frac{x^2}{2}} dx. \end{aligned}$$

It turns out that for a Brownian motion we can do the same. So, let $w_{\sigma^2}(t)$, $t \in [0, T]$, be a Brownian motion with diffusion rate²⁸ σ^2 defined on a probability space $(\Omega, \mathcal{F}, \mathbb{P})$. We introduce a new probability measure on Ω defined as follows

$$\mathbb{P}^h \{A\} = \mathbb{E} \mathbb{I}_A e^{\int_0^T h(t) dw_{\sigma^2}(t) - \frac{\sigma^2}{2} \int_0^T h^2 dt}, \quad A \in \mathcal{F},$$

where h is a fixed function from $L_2[0, T]$.

Proposition 7.1. *The process*

$$w_{\sigma^2}(t) - \sigma^2 \int_0^t h(s) ds, \quad t \in [0, T],$$

is a Brownian motion with diffusion rate σ^2 on the probability space $(\Omega, \mathcal{F}, \mathbb{P}^h)$.

We remark that the statement of Proposition 7.1 is a consequence of more general Cameron-Martin theorem about admissible shifts of Brownian motion (see [Kal02, Theorems 18.22]).

Exercise 7.3. Let $w(t)$, $t \in [0, T]$, be a Brownian motion with diffusion rate σ^2 defined on $(\Omega, \mathcal{F}, \mathbb{P})$ and $f \in H_0^2[0, T]$. Find a probability measure $\tilde{\mathbb{P}}$ such that $w(t) - f(t)$, $t \in [0, T]$, is a Brownian motion on $(\Omega, \mathcal{F}, \tilde{\mathbb{P}})$.

(Hint: Use Proposition 7.1 and definition of absolutely continuous functions)

Exercise 7.4. Show that for every $a \in \mathbb{R}$ and $\delta > 0$

$$\mathbb{P}\{w_{\sigma^2}(t) + at < \delta, \quad t \in [0, T]\} > 0.$$

(Hint: Use Proposition 7.1 and the fact that $\sup_{t \in [0, T]} w_{\sigma^2}(t)$ and $|w_{\sigma^2}(T)|$ have the same distribution²⁹)

²⁸ $w(t)$, $t \in [0, T]$, is a Brownian motion with $\text{Var } w(t) = \sigma^2 t$

²⁹see Proposition 13.13 [Kal02]

7.3 Proof of Schilder's theorem

7.3.1 Proof of weak LDP for Brownian motion

The goal of this lecture is to prove the LDP for a family $(w_\varepsilon)_{\varepsilon>0}$ of Brownian motions, where $w_\varepsilon(t) = \sqrt{\varepsilon}w(t)$, $t \in [0, T]$. The rigorous statement was formulated in Section 7.1 (see Theorem 7.1). In this section, we will prove the weak LDP.

For the proof of **the lower bound** we use Proposition 3.1. So, we need to show that

1) for every $f \in C_0^2[0, T]$

$$\lim_{r \rightarrow 0} \overline{\lim}_{\varepsilon \rightarrow 0} \varepsilon \ln \mathbb{P} \{w_\varepsilon \in B_r(f)\} \geq -I(f);$$

where

$$I(f) = \begin{cases} \frac{1}{2} \int_0^T \dot{f}^2(t) dt & \text{if } f \in H_0^2[0, T], \\ +\infty & \text{otherwise.} \end{cases}$$

2) For every $f \in H_0^2[0, T]$ there exists a sequence f_n , $n \geq 1$, from $C_0^2[0, T]$ such that $f_n \rightarrow f$ in $C_0[0, T]$ and $I(f_n) \rightarrow I(f)$, $n \rightarrow \infty$.

We start from checking 1). Take a function $f \in C_0^2[0, T]$ and estimate $\mathbb{P} \{\|w_\varepsilon - f\|_C < r\}$ from below, using Proposition 7.1. We set $h := f'$ and consider the following transformation of the probability measure \mathbb{P}

$$\mathbb{P}^{h, \varepsilon} \{A\} = \mathbb{E} \mathbb{I}_A e^{\int_0^T \frac{h(t)}{\varepsilon} dw_\varepsilon(t) - \frac{\varepsilon}{2} \int_0^T \frac{h^2(t)}{\varepsilon^2} dt} = \mathbb{E} \mathbb{I}_A e^{\frac{1}{\varepsilon} [\int_0^T h(t) dw_\varepsilon(t) - \frac{1}{2} \int_0^T h^2(t) dt]}.$$

Then the process

$$w_\varepsilon(t) - \varepsilon \int_0^t \frac{h(s)}{\varepsilon} ds = w_\varepsilon(t) - \int_0^t f'(s) ds = w_\varepsilon(t) - f(t), \quad t \in [0, T],$$

is a Brownian motion on the probability space $(\Omega, \mathcal{F}, \mathbb{P}^{h, \varepsilon})$ with diffusion rate ε , according to Proposition 7.1. Integrating by parts in the first integral,³⁰ we have

$$\int_0^T h(t) dw_\varepsilon(t) - \frac{1}{2} \int_0^T h^2(t) dt = h(T)w_\varepsilon(T) - \int_0^T h'(t)w_\varepsilon(t) dt - \frac{1}{2} \int_0^T h^2(t) dt =: \Phi(h, w_\varepsilon).$$

Now, we can estimate

$$\begin{aligned} \mathbb{P} \{w_\varepsilon \in B_r(f)\} &= \mathbb{E} \left[\mathbb{I}_{\{w_\varepsilon \in B_r(f)\}} e^{\frac{1}{\varepsilon} \Phi(h, w_\varepsilon)} e^{-\frac{1}{\varepsilon} \Phi(h, w_\varepsilon)} \right] \\ &\geq \mathbb{E} \left[\mathbb{I}_{\{w_\varepsilon \in B_r(f)\}} e^{\frac{1}{\varepsilon} \Phi(h, w_\varepsilon)} e^{-\frac{1}{\varepsilon} \sup_{g \in B_r(f)} \Phi(h, g)} \right] \\ &= e^{-\frac{1}{\varepsilon} \sup_{g \in B_r(f)} \Phi(h, g)} \mathbb{P}^{\varepsilon, h} \{\|w_\varepsilon - f\|_C < r\} = e^{-\frac{1}{\varepsilon} \sup_{g \in B_r(f)} \Phi(h, g)} \mathbb{P} \{\|w_\varepsilon\|_C < r\}, \end{aligned} \tag{25}$$

where the latter equality follows from Proposition 7.1. Hence,

$$\begin{aligned} \lim_{r \rightarrow 0} \overline{\lim}_{\varepsilon \rightarrow 0} \varepsilon \ln \mathbb{P} \{w_\varepsilon \in B_r(f)\} &\geq - \lim_{r \rightarrow 0} \sup_{g \in B_r(f)} \Phi(h, g) \\ &= \Phi(h, f) = h(T)f(T) - \int_0^T h'(t)f(t) dt - \frac{1}{2} \int_0^T h^2(t) dt \\ &= \int_0^T h^2(t) dt - \frac{1}{2} \int_0^T h^2(t) dt = I(f) \end{aligned}$$

³⁰see Exercise 7.7 below

because $\mathbb{P} \{ \|w_\varepsilon\|_C < r \} \rightarrow 1$ as $\varepsilon \rightarrow 0$,³¹ and the function Φ is continuous on $C_0[0, T]$ in the second argument. This finishes the proof of 1). The proof of 2) is proposed as an exercise (see Exercise 7.6).

Exercise 7.5. Let $w_\varepsilon(t)$, $t \in [0, T]$, denote a Brownian motion with diffusion rate ε for every $\varepsilon > 0$. Show that $\mathbb{P} \{ \|w_\varepsilon\|_C < r \} \rightarrow 1$ as $\varepsilon \rightarrow 0$, for all $r > 0$.

Exercise 7.6. Show that for any $f \in H_0^2[0, T]$ there exists a sequence $(f_n)_{n \geq 1}$ from E_0 such that $f_n \rightarrow f$ in $C_0[0, T]$ and $I(f_n) \rightarrow I(f)$ as $n \rightarrow \infty$.

(Hint: Use first the fact that $C^1[0, T]$ is dense in $L_2[0, T]$. Then show that if $h_n \rightarrow h$ in $L_2[0, T]$, then $\int_0^T h_n(s) ds$ tends to $\int_0^T h(s) ds$ in $C_0[0, T]$, using Hölder's inequality)

Exercise 7.7. Let $h \in C^1[0, T]$ and $w(t)$, $t \in [0, T]$, be a Brownian motion. Show that

$$\int_0^T h(t) dw(t) = h(T)w(T) - h(0)w(0) - \int_0^T h'(t)w(t) dt.$$

(Hint: Take a partition $0 = t_0 < t_1 < \dots < t_n = T$ and check first that functions $h_n = \sum_{k=1}^n h(t_k) \mathbb{I}_{[t_{k-1}, t_k]}$ converge to h in $L_2[0, T]$ as the mesh of partition goes to 0, using e.g. the uniform continuity of h on $[0, T]$. Next show that

$$\sum_{k=1}^n h(t_{k-1})(w(t_k) - w(t_{k-1})) = h(t_{n-1})w(T) - h(0)w(0) - \sum_{k=1}^{n-1} w(t_k)(h(t_k) - h(t_{k-1}))$$

Then prove that the first partial sum converges to the integral $\int_0^T h(t) dw(t)$ in L_2 and the second partial sum converges to $\int_0^T w(t) dh(t)$ a.s. as the mesh of partition goes to 0)

To prove **the upper bound**³² (20) for any compact set $F \subset C_0[0, T]$, we will use Proposition 6.1. We are going to show that for any $f \in C_0[0, T]$

$$\lim_{r \rightarrow 0} \overline{\lim}_{\varepsilon \rightarrow 0} \varepsilon \ln \mathbb{P} \{ w_\varepsilon \in B_r(f) \} \leq -I(f).$$

So we fix any $f \in C_0[0, T]$ and $h \in C^1[0, T]$, and estimate

$$\begin{aligned} \mathbb{P} \{ w_\varepsilon \in B_r(f) \} &= \mathbb{E} \left[\mathbb{I}_{\{w_\varepsilon \in B_r(f)\}} e^{\frac{1}{\varepsilon} \Phi(h, w_\varepsilon)} e^{-\frac{1}{\varepsilon} \Phi(h, w_\varepsilon)} \right] \\ &\leq \mathbb{E} \left[\mathbb{I}_{\{w_\varepsilon \in B_r(f)\}} e^{\frac{1}{\varepsilon} \Phi(h, w_\varepsilon)} e^{-\frac{1}{\varepsilon} \inf_{g \in B_r(f)} \Phi(h, g)} \right] \\ &\leq e^{-\frac{1}{\varepsilon} \inf_{g \in B_r(f)} \Phi(h, g)} \mathbb{E} e^{\frac{1}{\varepsilon} \Phi(h, w_\varepsilon)} = e^{-\frac{1}{\varepsilon} \inf_{g \in B_r(f)} \Phi(h, g)}, \end{aligned}$$

because $\mathbb{E} e^{\frac{1}{\varepsilon} \Phi(h, w_\varepsilon)} = 1$. The last equality follows from the fact that $\mathbb{P}^{h, \varepsilon}$ is a probability measure and $\mathbb{E} e^{\frac{1}{\varepsilon} \Phi(h, w_\varepsilon)}$ is the expectation of 1 with respect to $\mathbb{P}^{h, \varepsilon}$. Consequently,

$$\lim_{r \rightarrow 0} \overline{\lim}_{\varepsilon \rightarrow 0} \varepsilon \ln \mathbb{P} \{ w_\varepsilon \in B_r(f) \} \leq -\lim_{r \rightarrow 0} \inf_{g \in B_r(f)} \Phi(h, g) = -\Phi(h, f).$$

Now, taking infimum over all $h \in C^1[0, T]$, we obtain

$$\lim_{r \rightarrow 0} \overline{\lim}_{\varepsilon \rightarrow 0} \varepsilon \ln \mathbb{P} \{ w_\varepsilon \in B_r(f) \} \leq \inf_{h \in C^1[0, T]} (-\Phi(h, f)) = -\sup_{h \in C^1[0, T]} \Phi(h, f).$$

It remains only to show that

$$\sup_{h \in C^1[0, T]} \Phi(h, f) = I(f). \tag{26}$$

³¹see Exercise 7.5

³²The method applied here was taken from [DMRYZ04], see also [KvR19] for an infinite dimensional state space

7.3.2 A variational problem

We will first check equality (26) for the case $f \in C_0^2[0, T]$ because it is much more simple. The general case is based on the Riesz representation theorem and will be state in Proposition 7.2 below (see also [KvR19, Proposition 4.6]). We observe that for $f \in C_0^2[0, T]$ and any $h \in C^1[0, T]$

$$\begin{aligned}\Phi(h, f) &= h(T)f(T) - \int_0^T h'(t)f(t)dt - \frac{1}{2} \int_0^T h^2(t)dt \\ &= \int_0^T h(t)f'(t)dt - \frac{1}{2} \int_0^T h^2(t)dt \leq \frac{1}{2} \int_0^T f'^2(t)dt = I(f),\end{aligned}$$

where we first used the integration by parts and then the trivial inequality $ab \leq \frac{a^2}{2} + \frac{b^2}{2}$, $a, b \in \mathbb{R}$. Moreover, we see that the last inequality becomes an equality if $h = f'$. So, the supremum is attained at the point $h = f'$ and $\Phi(f', f) = I(f)$. This proves (26).

Proposition 7.2. *For each $f \in C_0[0, T]$*

$$\sup_{h \in C^1[0, T]} \Phi(h, f) = I(f),$$

where $\Phi(h, f) = h(T)f(T) - \int_0^T h'(t)f(t)dt - \frac{1}{2} \int_0^T h^2(t)dt$.

Proof. We first prove the assertion of the proposition for f satisfying

$$J(f) := \sup_{h \in C^1[0, T]} \Phi(h, f) < \infty.$$

Replacing h by θh , $\theta \in \mathbb{R}$, and using the linearity of $C^1[0, T]$, we get

$$J(f) = \sup_{h \in C^1[0, T]} \Phi(\theta h, f)$$

for all $\theta \in \mathbb{R}$. Next, we note that for each $h \in C^1[0, T]$ the function

$$\theta \mapsto \Phi(\theta h, f) = \theta G(h, f) - \frac{\theta^2}{2} \int_0^T h^2(t)dt,$$

where

$$G(h, f) = h(T)f(T) - \int_0^T h'(t)f(t)dt,$$

reaches its maximum at the point

$$\theta_h^{\max} = \frac{G(h, f)}{\int_0^T h^2(t)dt}$$

Consequently,

$$J(f) = \sup_{h \in C^1[0, T]} \Phi(\theta h, f) = \sup_{h \in C^1[0, T]} \Phi(\theta_h^{\max} h, f),$$

which implies

$$J(f) = \frac{1}{2} \sup_{h \in C^1[0, T]} \frac{G^2(h, f)}{\int_0^T h^2(t)dt} < \infty. \quad (27)$$

We can consider $C^1[0, T]$ as a linear subspace of $L_2[0, T]$, which is dense in $L_2[0, T]$. Therefore, the linear form

$$G_f : h \rightarrow G(h, f),$$

which is continuous on $C^1[0, T]$, by (27), can be extended to the space $L_2[0, T]$. Using the Riesz theorem, there exists a function $g_f \in L_2[0, T]$ such that

$$G_f(h) = G(h, f) = \int_0^T g_f(t)h(t)dt. \quad (28)$$

Exercise 7.8 (ii) and equality (28) imply that f is absolutely continuous and $\dot{f} = g_f$. Applying the Cauchy-Schwarz inequality to (28), we get

$$G(h, f)^2 \leq \int_0^T g_f^2(t)dt \cdot \int_0^T h^2(t)dt = 2I(f) \int_0^T h^2(t)dt,$$

with equality for h proportional to g_f . The latter inequality yields $J(f) \leq I(f)$ and since $C^1[0, T]$ is dense in $L_2[0, T]$, we get the equality $J(f) = I(f)$.

If $I(f) < \infty$, then f is absolutely continuous and $g_f = \dot{f}$ in (28), by Exercise 7.8 (i). So, $J(f) \leq I(f) < \infty$ and, consequently, we have $J(f) = I(f)$. This completes the proof of the proposition. \square

Exercise 7.8. Let $f \in C_0[0, T]$.

(i) Let f be absolutely continuous. Show that for every $h \in C^1[0, T]$

$$h(T)f(T) - \int_0^T h'(t)f(t)dt = \int_0^T h(t)\dot{f}(t)dt. \quad (29)$$

(Hint: Check first the equality if $f \in C[0, T]$. Then, in the general case, approximate f in $L_1[0, T]$ by continuous functions)

(ii) Let $g \in L_2[0, T]$ and for every $h \in C^1[0, T]$

$$h(T)f(T) - \int_0^T h'(t)f(t)dt = \int_0^T h(t)g(t)dt.$$

Show that f is absolutely continuous with $\dot{f} = g$.

(Hint: Consider the function $\tilde{f}(t) = \int_0^t g(s)ds$ and apply to $\int_0^T h(t)g(t)dt$ the integration by parts formula)

7.3.3 Exponential tightness of I

To finish the proof of Schilder's theorem, it remains to prove that $(w_\varepsilon)_{\varepsilon>0}$ is exponentially tight. Exercise 6.4 shows that the estimate

$$\mathbb{E} e^{\lambda|\xi_\varepsilon|} \leq \kappa \frac{1}{\varepsilon}$$

for some $\kappa > 0$, $\lambda > 0$ and all $\varepsilon > 0$, is enough to conclude the exponential tightness of $(\xi_\varepsilon)_{\varepsilon>0}$ in \mathbb{R} . It turns out, that a similar estimate allows to get exponential tightness in the space of continuous functions. However, one has to control the Hölder norm.

Proposition 7.3. Let $(\xi_\varepsilon)_{\varepsilon>0}$ be a family of random elements in $C_0[0, T]$. If there exist positive constants γ , λ and κ such that for all $s, t \in [0, T]$, $s < t$, and $\varepsilon > 0$

$$\mathbb{E} e^{\frac{\lambda}{\varepsilon} \frac{|\xi_\varepsilon(t) - \xi_\varepsilon(s)|}{(t-s)^\gamma}} \leq \kappa \frac{1}{\varepsilon},$$

then the family $(\xi_\varepsilon)_{\varepsilon>0}$ is exponentially tight in $C_0[0, T]$.

The proof of Proposition 7.3 follows from Corollary 7.1 [Sch97].

Lemma 7.1. Let $w(t)$, $t \in [0, T]$, be a standard Brownian motion. Then the family $(\sqrt{\varepsilon}w)_{\varepsilon>0}$ is exponentially tight in $C_0[0, T]$.

Proof. To check the exponential tightness of $(\sqrt{\varepsilon}w)_{\varepsilon>0}$, we will use Proposition 7.3. We first remark that $\mathbb{E} e^{\alpha(w(t)-w(s)) - \frac{\alpha^2}{2}(t-s)} = 1$ for all $\alpha \in \mathbb{R}$ and $s, t \in [0, T]$, $s < t$ (see Exercise 7.9 below). So, we can estimate for $\varepsilon > 0$, $s < t$ and $\alpha > 0$

$$\begin{aligned} \mathbb{E} e^{\alpha|\sqrt{\varepsilon}w(t) - \sqrt{\varepsilon}w(s)|} &\leq \mathbb{E} e^{\alpha(\sqrt{\varepsilon}w(t) - \sqrt{\varepsilon}w(s))} + \mathbb{E} e^{-\alpha(\sqrt{\varepsilon}w(t) - \sqrt{\varepsilon}w(s))} \\ &= 2\mathbb{E} e^{\alpha(\sqrt{\varepsilon}w(t) - \sqrt{\varepsilon}w(s))} = 2\mathbb{E} e^{\alpha\sqrt{\varepsilon}(w(t)-w(s)) - \frac{\alpha^2\varepsilon}{2}(t-s) + \frac{\alpha^2\varepsilon}{2}(t-s)} = 2e^{\frac{\alpha^2\varepsilon}{2}(t-s)}. \end{aligned}$$

Taking $\alpha := \frac{\sqrt{2}}{\varepsilon\sqrt{t-s}}$, we obtain

$$\mathbb{E} e^{\frac{\sqrt{2}|\sqrt{\varepsilon}w(t) - \sqrt{\varepsilon}w(s)|}{\varepsilon\sqrt{t-s}}} \leq 2\varepsilon^{\frac{1}{2}} \leq (2\varepsilon)^{\frac{1}{\varepsilon}}.$$

This implies the exponential tightness. □

Exercise 7.9. Let $w(t)$, $t \in [0, T]$, be a standard Brownian motion. Show directly that for each $\alpha \in \mathbb{R}$, $s, t \in [0, T]$, $s < t$

$$\mathbb{E} e^{\alpha(w(t)-w(s)) - \frac{\alpha^2}{2}(t-s)} = 1.$$

(Hint: Use Exercise 7.2)

Remark 7.1. Let $w(t)$, $t \in [0, T]$, be a standard Brownian motion on \mathbb{R}^d . Then using the same argument, one can prove that the family $(w_\varepsilon)_{\varepsilon>0}$ satisfies the LDP in $C_0([0, T], \mathbb{R}^d)$ with rate function

$$I(f) = \begin{cases} \frac{1}{2} \int_0^T \|\dot{f}(t)\|^2 dt & \text{if } f \in H_0^2([0, T], \mathbb{R}^d), \\ +\infty & \text{otherwise,} \end{cases}$$

where $C_0([0, T], \mathbb{R}^d) = \{f = (f_1, \dots, f_d) : f_i \in C_0[0, T], i = 1, \dots, d\}$, $H_0^2([0, T], \mathbb{R}^d) = \{f = (f_1, \dots, f_d) : f_i \in H_0^2[0, T], i = 1, \dots, d\}$ and $\dot{f} = (\dot{f}_1, \dots, \dot{f}_d)$.

8 Contraction principle and Freidlin-Wentzell theory

8.1 Contraction principle

The goal of this section is the transformation of LDP under a continuous map.

Theorem 8.1 (Contraction principle). *Consider a continuous function f between two metric spaces E and S , and let ξ_ε be random elements in E . If $(\xi_\varepsilon)_{\varepsilon>0}$ satisfies the LDP in E with rate function I , then the images $f(\xi_\varepsilon)$ satisfy the LDP in S with rate function*

$$J(y) = \inf \{I(x) : f(x) = y\} = \inf_{f^{-1}(\{y\})} I, \quad y \in S. \quad (30)$$

Moreover, J is a good rate function on S whenever the function I is good on E .

Proof. We take a closed set $F \subset S$ and denote $f^{-1}(F) = \{x : f(x) \in F\} \subset E$. Then

$$\begin{aligned} \overline{\lim}_{\varepsilon \rightarrow 0} \varepsilon \ln \mathbb{P} \{f(\xi_\varepsilon) \in F\} &= \overline{\lim}_{\varepsilon \rightarrow 0} \varepsilon \ln \mathbb{P} \{\xi_\varepsilon \in f^{-1}(F)\} \\ &\leq - \inf_{x \in f^{-1}(F)} I(x) = - \inf_{y \in F} \inf_{f(x)=y} I(x) = - \inf_{y \in F} J(y). \end{aligned}$$

The lower bound can be proved similarly.

When I is good, we claim that

$$\{J \leq \alpha\} = f(\{I \leq \alpha\}) = \{f(x) : I(x) \leq \alpha\}, \quad \alpha \geq 0. \quad (31)$$

To see this, fix any $\alpha \geq 0$, and let $x \in \{I \leq \alpha\}$, i.e. $I(x) \leq \alpha$. Then

$$J(f(x)) = \inf \{I(u) : f(u) = f(x)\} \leq I(x) \leq \alpha,$$

which means that $f(x) \in \{J \leq \alpha\}$. Since I is good and f is continuous, the infimum in (30) is attained at some $x \in E$, and we get $y = f(x)$ with $I(x) \leq \alpha$. Thus, $y \in f(\{I \leq \alpha\})$, which completes the proof of (31). Since continuous maps preserve compactness, $\{J \leq \alpha\}$ is compact, by (31). \square

Exercise 8.1. Let I be a good rate function on E and f be a continuous function from E to S . Show that the infimum in (30) is attained, that is, there exists $x \in E$ such that $f(x) = y$ and $J(y) = I(x)$.

Exercise 8.2. Let $w(t)$, $t \in [0, T]$, be a Brownian motion on \mathbb{R} with diffusion rate σ^2 and $w(0) = x_0$. Show that $(w(\varepsilon \cdot))_{\varepsilon>0}$ satisfies the LDP in $C[0, T]$ and find the associated rate function.

(Hint: Take the continuous map $\Phi(f)(t) = \sigma f(t) + x_0$, and use the contraction principle and Schilder's Theorem 7.1)

Remark 8.1. Let us explain the form of the rate function for Brownian motion using a concept of white noise and contraction principle. We recall that the white noise $\dot{w}(t)$, $t \in [0, T]$, formally can be defined as a Gaussian process with covariance $\mathbb{E} \dot{w}(t)\dot{w}(s) = \delta_0(t-s)$, where δ_0 denotes the Dirac delta function. One should interpret the white noise as a family of uncountable numbers of independent identically distributed Gaussian random variables. Similarly, as for Gaussian vectors, where the rate function is given by the formula $\frac{\|x\|^2}{2} = \sum_{k=1}^d \frac{x_k^2}{2}$, the rate function for the family $(\sqrt{\varepsilon}\dot{w})_{\varepsilon>0}$ should be

$$I_{\dot{w}}(x) = \frac{1}{2} \int_0^T x^2(t) dt.$$

We remark that a Brownian motion formally appears as a continuous function of white noise, namely the process $w(t) := \int_0^t \dot{w}(r) dr$, $t \in [0, T]$, defines a standard Brownian motion. Indeed, it is a Gaussian process with covariance

$$\mathbb{E} \left(\int_0^s \dot{w}(r_1) dr_1 \int_0^t \dot{w}(r_2) dr_2 \right) = \int_0^s \int_0^t \mathbb{E} \dot{w}(r_1)\dot{w}(r_2) dr_1 dr_2 = \int_0^s \int_0^t \delta_0(r_1-r_2) dr_1 dr_2 = \int_0^s 1 dr_1 = s$$

if $s \leq t$. So, $w = \Phi(\dot{w})$, where Φ denotes the integration procedure. By the contraction principle, the rate function of the family $(w_\varepsilon = \Phi(\dot{w}_\varepsilon))_{\varepsilon>0}$ has to be

$$I_w(x) = I_{\dot{w}}(\Phi^{-1}(x)) = \frac{1}{2} \int_0^T x'^2(t) dt.$$

Exercise 8.3. Let $B(t) = w(t) - tw(1)$, $t \in [0, 1]$, be a Brownian bridge on \mathbb{R} , where w is a standard Brownian motion on \mathbb{R} . Show that the family $(\sqrt{\varepsilon}B)_{\varepsilon>0}$ satisfy the LDP in $C_0[0, 1]$ with good rate function

$$I(f) = \begin{cases} \frac{1}{2} \int_0^1 f'^2(t) dt & \text{if } f \in H_0^2[0, 1] \text{ and } f(1) = 0, \\ +\infty & \text{otherwise} \end{cases}$$

(Hint: Use the contraction principle)

Exercise 8.4. Let $\xi = (\xi_n)_{n \geq 1}$ be a sequence of i.i.d. $N(0, 1)$ random variables. Use Schilder's theorem to show that the family $(\sqrt{\varepsilon}\xi)_{\varepsilon>0}$ satisfies the LDP in \mathbb{R}^∞ with the good rate function

$$I(x) = \begin{cases} \frac{1}{2} \sum_{n=1}^{\infty} x_n^2 & \text{if } x = (x_n)_{n \geq 1} \in l^2, \\ +\infty & \text{otherwise.} \end{cases}$$

(Hint: Consider the map $\Phi(f) = \left(\frac{f(t_n) - f(t_{n-1})}{\sqrt{t_n - t_{n-1}}} \right)_{n \geq 1}$ for an infinite partition $0 = t_0 < t_1 < \dots < t_n < \dots \rightarrow 1$ of the interval $[0, 1]$ and use the contraction principle)

8.2 Freidlin-Wentzell theory

In this section, we prove the LDP for solutions of stochastic differential equations (shortly SDE). Let consider a family $(z_\varepsilon)_{\varepsilon>0}$ of solutions to the following SDEs

$$dz_\varepsilon(t) = a(z_\varepsilon(t))dt + \sqrt{\varepsilon}dw(t), \quad z_\varepsilon(0) = 0, \quad (32)$$

where $a : \mathbb{R} \rightarrow \mathbb{R}$ is a bounded Lipschitz continuous³³ function and $w(t)$, $t \in [0, T]$, is a standard Brownian motion. We recall that a continuous process $z_\varepsilon(t)$, $t \in [0, T]$, is a solution to (32) if

$$z_\varepsilon(t) = \int_0^t a(z_\varepsilon(s))ds + \sqrt{\varepsilon}w(t), \quad t \in [0, T].$$

By Theorem 21.3 [Kal02], equation (32) has a unique solution.

Theorem 8.2 (Freidlin-Wentzell theorem). *For any bounded Lipschitz continuous function $a : \mathbb{R} \rightarrow \mathbb{R}$ the solutions the family $(z_\varepsilon)_{\varepsilon>0}$ satisfies the large deviation principle in $C_0[0, T]$ with good rate function*

$$I(f) = \begin{cases} \frac{1}{2} \int_0^T [\dot{f}(t) - a(f(t))]^2 dt & \text{if } f \in H_0^2[0, T], \\ +\infty & \text{otherwise.} \end{cases} \quad (33)$$

³³ $a : \mathbb{R} \rightarrow \mathbb{R}$ is Lipschitz continuous function if there exists a constant L such that $|a(x_1) - a(x_2)| \leq L|x_1 - x_2|$ for all $x_1, x_2 \in \mathbb{R}$

Proof. To prove the theorem, we will use the contraction principle. We first remark that the equation

$$z(t) = \int_0^t a(z(s))ds + g(t), \quad t \in [0, T], \quad (34)$$

has a unique solution for any $g \in C_0[0, T]$, since the function a is bounded and Lipschitz continuous. So, there exists a function $\Phi : C_0[0, T] \rightarrow C_0[0, T]$ such that $z = \Phi(g)$. Let us show that Φ is continuous. Take $g_1, g_2 \in C_0[0, T]$ and set $z_1 := \Phi(g_1)$, $z_2 := \Phi(g_2)$. Then one can estimate

$$\begin{aligned} |z_1(t) - z_2(t)| &= |\Phi(g_2) - \Phi(g_1)| = \left| \int_0^t a(z_1(s)) - a(z_2(s))ds + g_1(t) - g_2(t) \right| \\ &\leq \int_0^t |a(z_1(s)) - a(z_2(s))|ds + |g_1(t) - g_2(t)| \\ &\leq L \int_0^t |z_1(s) - z_2(s)|ds + \|g_1 - g_2\|_C. \end{aligned}$$

Gronwall's Lemma 21.4 [Kal02] yields $|z_1(t) - z_2(t)| \leq \|g_1 - g_2\|_C e^{Lt}$ for all $t \in [0, T]$. Hence,

$$\|\Phi(g_1) - \Phi(g_2)\|_C = \|z_1 - z_2\|_C \leq e^{LT} \|g_1 - g_2\|_C,$$

which shows that Φ is continuous. Using Schilder's theorem 7.1 along with the contraction principle (see Theorem 8.1), we conclude that the family $(z_\varepsilon)_{\varepsilon>0}$ satisfies the LDP in $C_0[0, T]$ with the good rate function

$$I(f) = \inf \{I_w(g) : \Phi(g) = f\} = \inf \left\{ I_w(g) : f(t) = \int_0^t a(f(s))ds + g(t) \right\},$$

where I_w is defined in Theorem 7.1. Due to the uniqueness of solutions to differential equation (34), the function Φ is bijective. Moreover, g and $f = \Phi(g)$ lie simultaneously in $H_0^2[0, T]$, in which case $\dot{g} = \dot{f} - a(f)$ almost everywhere.³⁴ Thus

$$I(f) = \begin{cases} \int_0^T (\dot{f}(t) - a(f(t)))^2 dt & \text{if } f \in H_0^2[0, T], \\ +\infty & \text{otherwise.} \end{cases}$$

□

Exercise 8.5. Let $\Phi : C_0[0, T] \rightarrow C_0[0, T]$ be defined in the proof of Theorem 8.2.

- 1) Show that the function Φ is bijective.
- 2) Prove that $g \in H_0^2[0, T]$ if and only if $f = \Phi(g) \in H_0^2[0, T]$.
 (Hint: Use equation (34) and the definition of $H_0^2[0, T]$)
- 3) Show that $\dot{g} = \dot{f} - a(f)$ almost everywhere for every $g \in H_0^2[0, T]$ and $f = \Phi(g)$.

³⁴see Exercise 8.5

8.3 Contraction principle for some discontinuous functions

In the first section, we showed that LDP is preserved under a continuous transformation. But very often one must work with discontinuous transformations. It turns out that LDP can also be preserved in some cases. Let us consider the following example which was taken from [DO10].

Given a standard Brownian motion $w(t)$, $t \in [0, T]$, in \mathbb{R}^d and a closed set $B \subset \mathbb{R}^d$, we consider the stopping time

$$\tau := \inf \{t : w(t) \in B\} \wedge T,$$

where $a \wedge b = \min \{a, b\}$. Let $y(t) := w(t \wedge \tau)$, $t \in [0, T]$, denote the stopped Brownian motion and $y_\varepsilon(t) := y(\varepsilon t)$, $t \in [0, T]$. We are interested in the LDP for the family $(y_\varepsilon)_{\varepsilon > 0}$. We remark that the process y_ε is obtained as an image of $w_\varepsilon(t) = w(\varepsilon t)$, $t \in [0, T]$. Indeed, let us define for a function $f \in C_0([0, T], \mathbb{R}^d)$

$$\tau(f) := \inf \{t : f(t) \in B\} \wedge T,$$

and

$$\Phi(f)(t) := f(t \wedge \tau(f)), \quad t \in [0, T]. \quad (35)$$

Then, by Exercise 8.6, Φ is a map from $C_0([0, T], \mathbb{R}^d)$ to $C_0([0, T], \mathbb{R}^d)$ and $y_\varepsilon = \Phi(w_\varepsilon)$. Unfortunately, we cannot apply the contraction principle here since Φ is discontinuous. But still, one can use the idea of contraction principle to obtain the LDP for $(y_\varepsilon)_{\varepsilon > 0}$. We remark also that the set B could be chosen by such a way that the set of discontinuous points of the map Φ has a positive Wiener measure³⁵ (for more details see Example 4.1 [DO10]).

Proposition 8.1. *The family $(y_\varepsilon)_{\varepsilon > 0}$ satisfies the LDP in $C_0([0, T], \mathbb{R}^d)$ with rate function*

$$I(f) = \begin{cases} \frac{1}{2} \int_0^T \dot{f}^2(t) dt & \text{if } f \in H_0^2([0, T], \mathbb{R}^d) \cap \text{Im } \Phi, \\ +\infty & \text{otherwise,} \end{cases} \quad (36)$$

where $\text{Im } \Phi = \{\Phi(f) : f \in C_0([0, T], \mathbb{R}^d)\}$.

Proof. A detailed proof of the proposition can be found in [DO10, Section 4]. We present here only the main idea. For the proof of the lower bound we take a closed set $F \subset C_0([0, T], \mathbb{R}^d)$ and estimate from above the upper limit

$$\begin{aligned} \overline{\lim}_{\varepsilon \rightarrow 0} \varepsilon \ln \mathbb{P} \{y_\varepsilon \in F\} &= \overline{\lim}_{\varepsilon \rightarrow 0} \varepsilon \ln \mathbb{P} \{\Phi(w_\varepsilon) \in F\} = \overline{\lim}_{\varepsilon \rightarrow 0} \varepsilon \ln \mathbb{P} \{w_\varepsilon \in \Phi^{-1}(F)\} \\ &\leq \overline{\lim}_{\varepsilon \rightarrow 0} \varepsilon \ln \mathbb{P} \left\{ w_\varepsilon \in \overline{\Phi^{-1}(F)} \right\} \leq - \frac{\inf_{\Phi^{-1}(F)} I_w}{\Phi^{-1}(F)}, \end{aligned}$$

where I_w is the rate function defined in Theorem 7.1 and $\Phi^{-1}(F) = \{f \in C_0([0, T], \mathbb{R}^d) : \Phi(f) \in F\}$. So, in order to obtain the upper bound (12), one needs to prove

$$\inf_{\Phi^{-1}(F)} I_w = \inf_{\Phi^{-1}(F)} I_w \left(= \inf_F \tilde{I} \right). \quad ^{36}$$

³⁵ $\mathbb{P} \{w \text{ is a point of discontinuity of } \Phi\} > 0$

³⁶We remark that this equality and the equality for open G trivially holds if Φ is continuous, since $\Phi^{-1}(F)$ is closed and $\Phi^{-1}(G)$ is open

Similarly, for the proof of the lower bound (13), one needs to show that

$$\inf_{\Phi^{-1}(G)^\circ} I_w = \inf_{\Phi^{-1}(G)} I_w \quad \left(= \inf_G \tilde{I} \right)$$

for any open set $G \subset C_0([0, T], \mathbb{R}^d)$. The prove of those equalities can be found in [DO10, Section 4] \square

Exercise 8.6. Let Φ be defined by (35) for $d = 1$ and $B = \{0\}$. Show that Φ maps $C[0, T]$ to $C[0, T]$. Prove that it is discontinuous.

9 Sanov's Theorem

Let $U = \{u_1, \dots, u_d\}$ be a finite set. We consider a family of i.i.d. random variables X_1, X_2, \dots taking values in U and for every $n \geq 1$ define the random probability measure on U as follows

$$\mu_n := \frac{1}{n} \sum_{k=1}^n \delta_{X_k}, \quad (37)$$

that is, for every $\varphi : U \rightarrow \mathbb{R}$

$$\int_U \varphi(u) \mu_n(du) = \frac{1}{n} \sum_{k=1}^n \varphi(X_k).$$

The random measures μ_n take a values in a metric space $\mathcal{P}(U)$ of all probability measures endowed with the distance of total variation.

Exercise 9.1. (i) Let $|\nu|_{TV}$ denote the total variation of a signed measure³⁷ on U . Show that

$$|\mu - \nu|_{TV} = \sum_{i=1}^d |\mu(\{u_i\}) - \nu(\{u_i\})|.$$

Therefore, the convergence of a sequence $(\nu_n)_{n \geq 1}$ to ν in $\mathcal{P}(U)$ is equivalent to the convergence of $\nu_n(\{u_i\}) \rightarrow \nu(\{u_i\})$, $n \rightarrow \infty$, for each $i \in [d]$.

(ii) Show that a sequence $(\nu_n)_{n \geq 1}$ converges in ν in $\mathcal{P}(U)$ if and only it $\nu_n \rightarrow \nu$ weakly.

(iii) Prove that the space $\mathcal{P}(U)$ is complete and separable.

(Hint: Use the isometry between $\mathcal{P}(U)$ and the simplex $\Delta = \{(x_1, \dots, x_d) \in \mathbb{R}^d : x_1 + \dots + x_d = 1\}$)

We remark that according to the strong law of large numbers the sequence of probability measures $(\mu_n)_{n \geq 1}$ converges almost surely to the distribution of X_1 denoted by μ (see Exercise 9.2). In this section, we are interested in the large deviation principle for the family $(\mu_n)_{n \geq 1}$ in the space $\mathcal{P}(U)$.

Exercise 9.2. Let μ_n be defined by (37) and μ be the distribution of X_1 , i.e. $\mu(\{u_i\}) = \mathbb{P}\{X_1 = u_i\}$, $i \in [d]$. Show that $\mu_n \rightarrow \mu$ in $\mathcal{P}(U)$ a.s.

(Hint: Use Exercise 9.1 and the strong law of large numbers)

³⁷The total variation $|\nu|_{TV}$ of a signed measure $\nu \in \mathcal{P}(U)$ is defined as $|\nu|_{TV} = \sup_{\pi} \sum_{A \in \pi} |\nu(A)|$, where is taken over all partitions π of the set U

9.1 Relative entropy

For simplicity of notation, we will further assume that the distribution μ of X_n , $n \geq 1$, from the previous section satisfies the property $\mu(\{u_i\}) > 0$ for all $i \in [d]$. For every $\nu \in \mathcal{P}(U)$ we define the **relative entropy** of ν given μ by

$$H(\nu|\mu) := \int_U \ln \left(\frac{\nu(\{u\})}{\mu(\{u\})} \right) \nu(du) = \sum_{i=1}^d \ln \left(\frac{\nu(\{u_i\})}{\mu(\{u_i\})} \right) \nu(\{u_i\}).$$

Exercise 9.3. (i) Show that the function $H(\cdot|\mu) : \mathcal{P}(U) \rightarrow \mathbb{R}$ is continuous.

(ii) Prove that $H(\nu|\mu) > 0$ for every $\nu \neq \mu$ and $H(\mu|\mu) = 0$.

(iii) Show that the function $H(\cdot|\mu)$ is good, that is, the level sets $\{\nu \in \mathcal{P}(U) : H(\nu|\mu) \leq \alpha\}$, $\alpha \geq 0$, are compact in $\mathcal{P}(U)$.

We are going to show that $(\mu_n)_{n \geq 1}$ satisfies the LDP in $\mathcal{P}(U)$ with the good rate function $I = H(\cdot|\mu)$. In order to prove this, we will need the following estimate.

Lemma 9.1. For every $f : U \rightarrow \mathbb{R}$ and $\nu \in \mathcal{P}(U)$ one has

$$\int_U f(u)\nu(du) - \ln \int_U e^{f(u)}\mu(du) \leq H(\nu|\mu).$$

Moreover, the equality is reached if and only if the function $\frac{e^{f(u)}\mu(\{u\})}{\nu(\{u\})}$ is constant on $\{u \in U : \nu(\{u\}) > 0\}$.

Proof. Using the Jensen inequality, we can estimate

$$\begin{aligned} \int_U f(u)\nu(du) - H(\nu|\mu) &= \int_U f(u)\nu(du) - \int_U \ln \left(\frac{\nu(\{u\})}{\mu(\{u\})} \right) \nu(du) \\ &= \int_U \ln e^{f(u)}\nu(du) - \int_U \ln \left(\frac{\nu(\{u\})}{\mu(\{u\})} \right) \nu(du) \\ &= \int_U \ln \left(\frac{e^{f(u)}\mu(\{u\})}{\nu(\{u\})} \right) \nu(du) \leq \ln \left(\int_U \frac{e^{f(u)}\mu(\{u\})}{\nu(\{u\})} \nu(du) \right) \\ &= \ln \left(\int_U e^{f(u)}\mu(du) \right). \end{aligned}$$

Since the function \ln is strictly concave, the equality is reached if and only if the function $\frac{e^{f(u)}\mu(\{u\})}{\nu(\{u\})}$, $u \in U$, is constant on $\{u : \nu(u) > 0\}$. The lemma is proved. \square

9.2 Sanov's theorem (particular case)

Theorem 9.1 (Sanov). Let X_1, X_2, \dots be i.i.d. U -valued random variables with distribution μ , where U is a finite set and $\mu(\{u\}) > 0$ for every $u \in U$. The family

$$\mu_n = \frac{1}{n} \sum_{k=1}^n \delta_{X_k}, \quad n \geq 1,$$

satisfies the LDP in $\mathcal{P}(U)$ with rate function $H(\cdot|\mu)$.

Proof. We will prove the theorem combining multidimensional Cramer Theorem 4.1 and the contraction principle. We consider the simplex $\Delta = \{(x_1, \dots, x_d) \in \mathbb{R}^d : x_1 + \dots + x_d = 1, x_i \geq 0, i \in [d]\}$ as a metric subspace of \mathbb{R}^d . Since the set Δ is closed in \mathbb{R}^d , the space Δ is a complete separable metric space. We next define the following map

$$\Phi(x) = \sum_{i=1}^d x_i \delta_{u_i}, \quad x = (x_i)_{i \in [d]},$$

from Δ to $\mathcal{P}(U)$. By Exercise 9.1, the map Φ is continuous. We consider the following random vectors $\xi_n = (\mathbb{I}_{\{X_n = u_i\}})_{i \in [d]}$, $n \geq 1$. It is trivial that $\xi_n \in \Delta$ for all $n \geq 1$. Hence, the random vectors $\frac{S_n}{n} = \frac{\xi_1 + \dots + \xi_n}{n}$, $n \geq 1$, take values from Δ . We remark that

$$\Phi\left(\frac{1}{n}S_n\right) = \mu_n, \quad n \geq 1.$$

Indeed,

$$\Phi\left(\frac{1}{n}S_n\right) = \sum_{i=1}^d \left(\frac{1}{n} \sum_{k=1}^n \mathbb{I}_{\{X_k = u_i\}}\right) \delta_{u_i} = \frac{1}{n} \sum_{k=1}^n \left(\sum_{i=1}^d \mathbb{I}_{\{X_k = u_i\}} \delta_{u_i}\right) = \frac{1}{n} \sum_{k=1}^n \delta_{X_k} = \mu_n.$$

Therefore, the LDP for $(\mu_n)_{n \geq 1}$ will directly follow from the contraction principle and the LDP for $(\frac{1}{n}S_n)_{n \geq 1}$ in Δ .

By multidimensional Cramer Theorem 4.1, the empirical means $(\frac{1}{n}S_n)_{n \geq 1}$ satisfy the LDP in \mathbb{R}^d with rate function φ^* that is the Fenchel-Legendre transform of the cumulant generating function

$$\varphi(\lambda) = \ln \mathbb{E} e^{\lambda \cdot \xi_1}, \quad \lambda \in \mathbb{R}^d,$$

where trivially the function $\varphi(\lambda) < \infty$ for every $\lambda \in \mathbb{R}^d$. Let us show that this family satisfies the LDP in the metric space Δ with rate function $\varphi^*(x)$, $x \in \Delta$. The upper bound immediately follows from the fact that every closed set F in Δ is also closed in \mathbb{R}^d . We will only prove the lower bound. Let G be an open set in Δ . Then there exists an open set \tilde{G} in \mathbb{R}^d such that $G = \tilde{G} \cap \Delta$. Using the LDP for $(\frac{1}{n}S_n)_{n \geq 1}$ in \mathbb{R}^d , we get

$$\liminf_{n \rightarrow \infty} \frac{1}{n} \ln \mathbb{P} \left\{ \frac{1}{n}S_n \in G \right\} = \liminf_{n \rightarrow \infty} \frac{1}{n} \ln \mathbb{P} \left\{ \frac{1}{n}S_n \in \tilde{G} \right\} \geq - \inf_{x \in \tilde{G}} \varphi^*(x) \geq - \inf_{x \in G} \varphi^*(x).$$

By Theorem 8.1, the family $(\mu_n)_{n \geq 1}$ satisfies the LDP in $\mathcal{P}(U)$ with rate function

$$I(\nu) = \inf \{ \varphi^*(x) : \Phi(x) = \nu, x \in \Delta \}, \quad \nu \in \mathcal{P}(U).$$

Since the function Φ is bijective with

$$\Phi^{-1}(\nu) = (\nu(\{u_i\}))_{i \in [d]}, \quad \nu \in \mathcal{P}(U),$$

we can conclude that $I(\nu) = \varphi^*((\nu(\{u_i\}))_{i \in [d]})$ for all $\nu \in \mathcal{P}(U)$.

In order to show that $I(\nu) = H(\nu|\mu)$, we first compute the cumulant generating function φ of ξ_1 . For $\lambda = (\lambda_i)_{i \in [d]} \in \mathbb{R}^d$ we have

$$\begin{aligned} \varphi(\lambda) &= \ln \left(\mathbb{E} e^{\sum_{i=1}^d \lambda_i \mathbb{I}_{\{X_1 = u_i\}}} \right) = \ln \mathbb{E} \left(\sum_{i=1}^d e^{\lambda_i} \mathbb{I}_{\{X_1 = u_i\}} \right) \\ &= \ln \left(\sum_{i=1}^d e^{\lambda_i} \mu(\{u_i\}) \right). \end{aligned}$$

Then for $\nu \in \mathcal{P}(U)$

$$\begin{aligned} I(\nu) &= \sup_{\lambda \in \mathbb{R}^d} \left\{ \sum_{i=1}^d \lambda_i \nu(\{u_i\}) - \varphi(\lambda) \right\} = \sup_{\lambda \in \mathbb{R}^d} \left\{ \sum_{i=1}^d \lambda_i \nu(\{u_i\}) - \ln \left(\sum_{i=1}^d e^{\lambda_i} \mu(\{u_i\}) \right) \right\} \\ &= \sup_{\lambda \in \mathbb{R}^d} \left\{ \int_U f_\lambda(u) \nu(du) - \ln \left(\int_U e^{f_\lambda(u)} \mu(du) \right) \right\} \leq H(\nu|\mu), \end{aligned}$$

by Lemma 9.1, where the function $f : U \rightarrow \mathbb{R}$ is defined as $f_\lambda(u_i) = \lambda_i$, $i \in [d]$. Taking $\lambda \in \mathbb{R}^d$ such that $\frac{e^{f_\lambda(u_i)} \mu(\{u_i\})}{\nu(\{u_i\})} = \frac{e^{\lambda_i} \mu(\{u_i\})}{\nu(\{u_i\})}$ is constant on $\{i : \nu(\{u_i\}) > 0\}$, we get that $I(\nu) = H(\nu|\mu)$, according to the same lemma.

The fact that $H(\cdot|\mu)$ is a good rate function follows from Exercise 9.3 (iii). □

Exercise 9.4. Let ξ_1, ξ_2, \dots be independent Bernoulli distributed random variables with parameter $p \in (0, 1)$. Using Sanov Theorem 9.1 and the contraction principle show that the family $(\frac{1}{n} S_n)_{n \geq 1}$ satisfies the large deviation principle with good rate function

$$I(x) = \begin{cases} x \ln \frac{x}{p} + (1-x) \ln \frac{1-x}{1-p} & \text{if } x \in [0, 1], \\ +\infty & \text{otherwise,} \end{cases}$$

where $S_n = \xi_1 + \dots + \xi_n$.

9.3 Sanov's theorem (general case)

In this section, we will extend Sanov's theorem to the case of a complete separable metric space U . Let U be a complete separable metric space and $\mathcal{P}(U)$ is the space of probability measures on $\mathcal{P}(U)$, equipped with the topology of weak convergence, under which $\mathcal{P}(U)$ is a complete separable metric space. Let also μ be a fixed probability measure from $\mathcal{P}(U)$. We first define on the space $\mathcal{P}(U)$ the relative entropy $H(\cdot|\mu)$.

We recall that by the Radon-Nikodym theorem, ν has a density³⁸ $\frac{d\nu}{d\mu}$ with respect to μ if and only if ν is **absolutely continuous** with respect to μ , i.e. $\nu(A) = 0$ for all $A \in \mathcal{B}(U)$ such that $\mu(A) = 0$. We write $\nu \ll \mu$ if ν is absolutely continuous with respect to μ . For every $\nu \in \mathcal{P}(U)$ we define the **relative entropy** $H(\nu|\mu)$ of ν given μ as

$$H(\nu|\mu) := \begin{cases} \int_U \ln \left(\frac{d\nu}{d\mu}(u) \right) \nu(du) & \text{if } \nu \ll \mu, \\ +\infty & \text{otherwise.} \end{cases}$$

Exercise 9.5. Prove that $H(\nu|\mu) \geq 0$ for every $\nu \in \mathcal{P}(U)$.

Theorem 9.2 (Sanov). Let X_1, X_2, \dots be i.i.d. random variables taking values in U with the distribution function μ , where U is a complete separable metric space and $\mu \in \mathcal{P}(U)$. Let

$$\mu_n := \frac{1}{n} \sum_{k=1}^n \delta_{X_k}, \quad n \geq 1,$$

be the empirical laws of $(X_n)_{n \geq 1}$. Then the family $(\mu_n)_{n \geq 1}$ satisfies the LDP in $\mathcal{P}(U)$ with good rate function $H(\cdot|\mu)$.

³⁸A function $f : U \rightarrow [0, \infty)$ is a density of ν with respect to μ if $\int_U f(u) \mu(du) < \infty$ and $\nu(A) = \int_A f(u) \mu(du)$ for all $A \in \mathcal{B}(U)$. Such a function is uniquely defined up to a.s. equality with respect to μ

Proof. The proof of the theorem can be found e.g. in [Swa12, Theorem 2.14] or [DZ98, Teorem 6.2.10]. \square

10 Varadhan's lemma

In this section, we discuss an equivalent version of the upper and lower bounds in the large deviation principle (see (12) and (13), respectively) via continuous bounded functions. We start from the following simple observation. Let $\xi \sim N(0, 1)$. We consider a bounded function f and formally compute the following limit

$$\lim_{\varepsilon \rightarrow 0} \varepsilon \ln \mathbb{E} e^{f(\sqrt{\varepsilon}\xi)/\varepsilon} = \lim_{\varepsilon \rightarrow 0} \varepsilon \ln \int_{-\infty}^{+\infty} e^{\frac{f(x)}{\varepsilon}} \frac{1}{\sqrt{2\pi\varepsilon}} e^{-\frac{x^2}{2\varepsilon}} dx.$$

In spirit of Exercise 3.3, we should expect that

$$\begin{aligned} \lim_{\varepsilon \rightarrow 0} \varepsilon \ln \mathbb{E} e^{f(\sqrt{\varepsilon}\xi)/\varepsilon} &= \lim_{\varepsilon \rightarrow 0} \varepsilon \ln \int_{-\infty}^{+\infty} e^{\frac{f(x)}{\varepsilon}} \frac{1}{\sqrt{2\pi\varepsilon}} e^{-\frac{x^2}{2\varepsilon}} dx \quad \text{"="} \quad \sup_{x \in \mathbb{R}} \left(\lim_{\varepsilon \rightarrow 0} \varepsilon \ln \frac{1}{\sqrt{2\pi\varepsilon}} e^{\frac{f(x)}{\varepsilon} - \frac{x^2}{2\varepsilon}} \right) \\ &= \sup_{x \in \mathbb{R}} \left(f(x) - \frac{x^2}{2} \right) = \sup_{x \in \mathbb{R}} (f(x) - I(x)), \end{aligned}$$

where I is the rate function associated with the family $(\sqrt{\varepsilon}\xi)_{\varepsilon > 0}$. It turns out, that the same equality takes place for any family $(\xi_\varepsilon)_{\varepsilon > 0}$ satisfying the LDP with a rate function I .

Lemma 10.1 (Varadhan). *Let the family $(\xi_\varepsilon)_{\varepsilon > 0}$ satisfy the LDP in a metric space E with a rate function I , and a function $f : E \rightarrow \mathbb{R}$ be continuous and bounded above, then*

$$\Lambda_f := \lim_{\varepsilon \rightarrow 0} \varepsilon \ln \mathbb{E} e^{f(\xi_\varepsilon)/\varepsilon} = \sup_{x \in E} (f(x) - I(x)). \quad (38)$$

Proof. We first show the lower bound for Λ_f . Let $x \in E$ and $G := B_r(x) \subset E$ be the open ball with the center x and a radius $r > 0$. We estimate

$$\begin{aligned} \underline{\lim}_{\varepsilon \rightarrow 0} \varepsilon \ln \mathbb{E} e^{f(\xi_\varepsilon)/\varepsilon} &\geq \underline{\lim}_{\varepsilon \rightarrow 0} \varepsilon \ln \mathbb{E} e^{f(\xi_\varepsilon)/\varepsilon} \mathbb{I}_{\{\xi_\varepsilon \in G\}} \geq \underline{\lim}_{\varepsilon \rightarrow 0} \varepsilon \ln \left(\inf_{y \in G} e^{f(y)/\varepsilon} \mathbb{P} \{ \xi_\varepsilon \in G \} \right) \\ &= \underline{\lim}_{\varepsilon \rightarrow 0} \left(\varepsilon \ln e^{\inf_{y \in G} f(y)/\varepsilon} + \varepsilon \ln \mathbb{P} \{ \xi_\varepsilon \in G \} \right) \\ &= \inf_{y \in G} f(y) - \inf_{y \in G} I(y) \geq \inf_{y \in G} f(y) - I(x). \end{aligned}$$

Making $r \rightarrow 0$ and using the continuity of f , we get

$$\underline{\lim}_{\varepsilon \rightarrow 0} \varepsilon \ln \mathbb{E} e^{f(\xi_\varepsilon)/\varepsilon} \geq f(x) - I(x)$$

for every $x \in E$. Therefore,

$$\underline{\lim}_{\varepsilon \rightarrow 0} \varepsilon \ln \mathbb{E} e^{f(\xi_\varepsilon)/\varepsilon} \geq \sup_{x \in E} (f(x) - I(x)).$$

In order to prove the upper bound, we fix $n \geq 1$ and choose finitely many closed sets $B_1, \dots, B_m \subset E$ (non necessarily disjoint) such that $f(x) \leq -n$ for all $x \in B_0 := (\bigcup_{k=1}^m B_k)^c$ and the oscillation of f on each B_k is at most $\frac{1}{n}$, i.e.

$$\sup_{x \in B_k} f(x) - \inf_{x \in B_k} f(x) \leq \frac{1}{n}, \quad k \in [m].$$

The existence of such a collection of closed sets follows from Exercise 10.1. Then, using Exercise 10.2, we estimate

$$\begin{aligned} \overline{\lim}_{\varepsilon \rightarrow 0} \varepsilon \ln \mathbb{E} e^{f(\xi_\varepsilon)/\varepsilon} &\leq \overline{\lim}_{\varepsilon \rightarrow 0} \varepsilon \ln \mathbb{E} \left(\sum_{k=0}^m e^{f(\xi_\varepsilon)/\varepsilon} \mathbb{1}_{\{\xi_\varepsilon \in B_k\}} \right) \\ &= \max_{k \in [m] \cup \{0\}} \left\{ \overline{\lim}_{\varepsilon \rightarrow 0} \varepsilon \ln \mathbb{E} \left(e^{f(\xi_\varepsilon)/\varepsilon} \mathbb{1}_{\{\xi_\varepsilon \in B_k\}} \right) \right\} \\ &\leq \max_{k \in [m] \cup \{0\}} \left\{ \overline{\lim}_{\varepsilon \rightarrow 0} \varepsilon \ln \left(\sup_{x \in B_k} e^{f(x)/\varepsilon} \mathbb{P} \{ \xi_\varepsilon \in B_k \} \right) \right\} \\ &\leq \max_{k \in [m]} \left\{ \sup_{x \in B_k} f(x) - \inf_{x \in B_k} I(x) \right\} \vee (-n) \\ &\leq \max_{k \in [m]} \left\{ \sup_{x \in B_k} \left(f(x) - I(x) + \frac{1}{n} \right) \right\} \vee (-n) \\ &\leq \sup_{x \in E} \left(f(x) - I(x) + \frac{1}{n} \right) \vee (-n). \end{aligned}$$

Making $n \rightarrow +\infty$, we obtain the upper bound. This implies the statement of the lemma. \square

Exercise 10.1. Let $f : E \rightarrow \mathbb{R}$ be a continuous and bounded above function. Show that for every $n \geq 1$ there exists a family of closed subsets B_k , $k \in [m]$, of E such that $f \leq -n$ on $B_0 := (\bigcup_{k=1}^m B_k)^c$ and the oscillation of f on each B_k is at most $\frac{1}{n}$.

Hint: Consider the sets $f^{-1} \left(\left[\frac{k-1}{n}, \frac{k}{n} \right] \right)$, $k \in \mathbb{Z}$.

Exercise 10.2. Let B be a subset of E and $f, g : A \rightarrow \mathbb{R}$ and $\inf_{x \in A} g(x) > -\infty$. Prove that

$$\inf_{x \in A} f(x) - \inf_{x \in A} g(x) \leq \sup_{x \in A} (f(x) - g(x)).$$

We remark that the inverse statement to Varadan's lemma also holds. Denote the family of bounded continuous function $f : E \rightarrow \mathbb{R}$ by $C_b(E)$.

Theorem 10.1 (Bruc). *Let the family $(\xi_\varepsilon)_{\varepsilon>0}$ be exponentially tight and the limit Λ_f in (38) exist for every $f \in C_b(E)$. Then $(\xi_\varepsilon)_{\varepsilon>0}$ satisfies the LDP with good rate function*

$$I(x) = \sup_{f \in C_b(E)} (f(x) - \Lambda_f(x)), \quad x \in E.$$

Proof. The proof of the theorem can be found in [Kal02, Theorem 27.10]. \square

11 Exponential equivalence

In order to prove that a family $(\xi_\varepsilon)_{\varepsilon>0}$ satisfies a large deviation principle with a given rate function, it is often convenient to replace the random elements ξ_ε by some other random elements η_ε that are “sufficiently close”, so that the LDP for $(\eta_\varepsilon)_{\varepsilon>0}$ implies the LDP for $(\xi_\varepsilon)_{\varepsilon>0}$. We consider the following example.

Example 11.1. Let $a, b : \mathbb{R} \rightarrow \mathbb{R}$ be bounded Lipschitz continuous functions and $w(t)$, $t \in [0, T]$ be a standard Brownian motion. We are interested in the LDP in $C_0[0, T]$ for solutions to the following perturbed SDEs

$$dx_\varepsilon(t) = a(x_\varepsilon(t))dt + \varepsilon b(x_\varepsilon(t))dt + \sqrt{\varepsilon}dw(t), \quad x_\varepsilon(0) = 0.$$

Let us compare the solutions to those equations with the solutions to the non-perturbed SDEs

$$dz_\varepsilon(t) = a(z_\varepsilon(t))dt + \sqrt{\varepsilon}dw(t), \quad z_\varepsilon(0) = 0.$$

for which we have already proved the LDP (see Section 8.2). We estimate

$$\begin{aligned} |x_\varepsilon(t) - z_\varepsilon(t)| &= \left| \int_0^t (a(x_\varepsilon(s)) - a(z_\varepsilon(s)))ds + \varepsilon \int_0^t b(x_\varepsilon(s))ds \right| \\ &\leq \int_0^t |a(x_\varepsilon(s)) - a(z_\varepsilon(s))|ds + \varepsilon \|b\|_C T \\ &\leq L \int_0^t |x_\varepsilon(s) - z_\varepsilon(s)|ds + \varepsilon \|b\|_C T, \quad t \in [0, T], \end{aligned}$$

where $\|\cdot\|_C$ denotes the supremum norm in $C_0[0, T]$ and L is the Lipschitz constant for a . Using Gronwall’s Lemma 21.4 [Kal02], we get $|x_\varepsilon(t) - z_\varepsilon(t)| \leq \varepsilon \|b\|_C T e^{LT}$, $t \in [0, T]$. Hence

$$\|x_\varepsilon - z_\varepsilon\|_C \leq \varepsilon \|b\|_C T.$$

In particular, this implies that for every $\delta > 0$

$$\lim_{\varepsilon \rightarrow 0} \varepsilon \ln \mathbb{P} \{ \|x_\varepsilon - z_\varepsilon\| > \delta \} = -\infty.$$

As we will see later this is enough to conclude that the family $(x_\varepsilon)_{\varepsilon>0}$ satisfies the LDP in $C_0[0, T]$ with the same good rate function as $(z_\varepsilon)_{\varepsilon>0}$, which is defined by (33).

Let (E, d) be a complete separable metric space.

Definition 11.1. We will say that families $(\xi_\varepsilon)_{\varepsilon>0}$ and $(\eta_\varepsilon)_{\varepsilon>0}$ of random elements in E are **exponentially equivalent** if for every $\delta > 0$

$$\lim_{\varepsilon \rightarrow 0} \varepsilon \ln \mathbb{P} \{ d(\xi_\varepsilon, \eta_\varepsilon) > \delta \} = -\infty.$$

Proposition 11.1. Let families $(\xi_\varepsilon)_{\varepsilon>0}$ and $(\eta_\varepsilon)_{\varepsilon>0}$ of random elements in a separable metric space E be exponentially equivalent. Then $(\xi_\varepsilon)_{\varepsilon>0}$ satisfies the LDP with a good rate function I iff the same LDP holds for $(\eta_\varepsilon)_{\varepsilon>0}$.

Proof. Suppose that the LDP holds for $(\xi_\varepsilon)_{\varepsilon>0}$ with rate function I . We fix any closed set $F \subseteq E$, and denote the closed δ -neighborhood of F by F^δ , i.e.

$$F^\delta = \{x \in E : d(x, F) \leq \delta\},$$

where $d(x, F) = \inf_{y \in F} d(x, y)$. Then one has

$$\begin{aligned} \mathbb{P} \{\eta_\varepsilon \in F\} &\leq \mathbb{P} \{\eta_\varepsilon \in F, d(\xi_\varepsilon, \eta_\varepsilon) \leq \delta\} + \mathbb{P} \{d(\xi_\varepsilon, \eta_\varepsilon) > \delta\} \\ &\leq \mathbb{P} \{\xi_\varepsilon \in F^\delta\} + \mathbb{P} \{d(\xi_\varepsilon, \eta_\varepsilon) > \delta\} \end{aligned}$$

for all $\varepsilon > 0$. Using the LDP for $(\xi_\varepsilon)_{\varepsilon>0}$, the exponential equivalence and Exercise 3.3, we can estimate

$$\begin{aligned} \overline{\lim}_{\varepsilon \rightarrow 0} \varepsilon \ln \mathbb{P} \{\eta_\varepsilon \in F\} &\leq \overline{\lim}_{\varepsilon \rightarrow 0} \varepsilon \ln \left(\mathbb{P} \{\xi_\varepsilon \in F^\delta\} + \mathbb{P} \{d(\xi_\varepsilon, \eta_\varepsilon) > \delta\} \right) \\ &\leq \max \left\{ \overline{\lim}_{\varepsilon \rightarrow 0} \varepsilon \ln \mathbb{P} \{\xi_\varepsilon \in F^\delta\}, \overline{\lim}_{\varepsilon \rightarrow 0} \varepsilon \ln \mathbb{P} \{d(\xi_\varepsilon, \eta_\varepsilon) > \delta\} \right\} \\ &\leq \max \left\{ - \inf_{x \in F^\delta} I(x), -\infty \right\} = - \inf_{x \in F^\delta} I(x). \end{aligned}$$

Since I is a good rate function, one can show that $\inf_{x \in F^\delta} I(x) \rightarrow \inf_{x \in F} I(x)$ as $\delta \rightarrow 0$ (see Exercise 11.2).

This implies the upper bound (12) for the family $(\eta_\varepsilon)_{\varepsilon>0}$.

We next prove the lower bound (13) for $(\eta_\varepsilon)_{\varepsilon>0}$. Let G be a fixed open subset of E and $x \in G$. If $d(x, G^c) > 3\delta > 0$, then the ball $B_\delta(x)$ is contained in G . We estimate as before

$$\begin{aligned} \mathbb{P} \{\xi_\varepsilon \in B_\delta(x)\} &\leq \mathbb{P} \{\xi_\varepsilon \in B_\delta(x), d(\xi_\varepsilon, \eta_\varepsilon) \leq \delta\} + \mathbb{P} \{d(\xi_\varepsilon, \eta_\varepsilon) > \delta\} \\ &\leq \mathbb{P} \{\eta_\varepsilon \in G\} + \mathbb{P} \{d(\xi_\varepsilon, \eta_\varepsilon) > \delta\}. \end{aligned}$$

Therefore, using the LDP for $(\xi_\varepsilon)_{\varepsilon>0}$ and the exponential equivalence, we get

$$\begin{aligned} -I(x) &\leq - \inf_{y \in B_\delta(x)} I(y) \leq \underline{\lim}_{\varepsilon \rightarrow 0} \varepsilon \ln \mathbb{P} \{\xi_\varepsilon \in B_\delta(x)\} \leq \underline{\lim}_{\varepsilon \rightarrow 0} \varepsilon \ln (\mathbb{P} \{\eta_\varepsilon \in G\} + \mathbb{P} \{d(\xi_\varepsilon, \eta_\varepsilon) > \delta\}) \\ &\leq \max \left\{ \underline{\lim}_{\varepsilon \rightarrow 0} \varepsilon \ln \mathbb{P} \{\eta_\varepsilon \in G\}, \overline{\lim}_{\varepsilon \rightarrow 0} \varepsilon \ln \mathbb{P} \{d(\xi_\varepsilon, \eta_\varepsilon) > \delta\} \right\} = \underline{\lim}_{\varepsilon \rightarrow 0} \varepsilon \ln \mathbb{P} \{\eta_\varepsilon \in G\}. \end{aligned}$$

Taking the supremum over all $x \in G$, we obtain the lower bound. The proposition is proved. \square

Exercise 11.1. Let F be a closed subset of E . Show that the closed δ -neighborhood $F^\delta = \{x \in E : d(x, F) \leq \delta\}$ of F is a closed set and $\bigcap_{\delta>0} F^\delta = F$.

Exercise 11.2. Let $I : E \rightarrow [0, \infty]$ be good, F be a closed set and F^δ be the closed δ -neighborhood of F . Show that $\inf_{x \in F^\delta} I(x) \rightarrow \inf_{x \in F} I(x)$ as $\delta \rightarrow 0$.

12 Some applications of large deviations

12.1 Curie-Weiss model of ferromagnetism

This section is taken from [RAS15, Section 3.4].

In this section, we consider an application of LDP in statistical mechanics, using a toy model of ferromagnetism. Let us imagine that a piece of material is magnetized by subjecting it to a magnetic field. Then assume that the field is turned off. We are interesting if the magnetization persist. To answer this question, we introduce a model called the Curie-Weiss ferromagnet and will try to understand this using large deviation theory.

Let us start from the description of the model. Consider n atoms each of them have a ± 1 valued spin ω_i , $i = 1, \dots, n$. The space of n -spin configurations is $\Omega_n = \{-1, 1\}^n$. The energy of the system is given by the **Hamiltonian**

$$\mathcal{H}_n(\omega) = -\frac{J}{2n} \sum_{i,j=1}^n \omega_i \omega_j - h \sum_{j=1}^n \omega_j = -\frac{J}{2} \sum_{i=1}^n \omega_i \left(\frac{1}{n} \sum_{j=1}^n \omega_j \right) - h \sum_{j=1}^n \omega_j. \quad (39)$$

A ferromagnet has a positive coupling constant $J > 0$ and $h \in \mathbb{R}$ is the external magnetic field. Since nature prefers low energy, ferromagnet spins tend to align with each other and with the magnetic field h , if $h \neq 0$. The **Gibbs measure** for n spins is

$$\gamma_n(\omega) = \frac{1}{Z_n} e^{-\beta \mathcal{H}_n(\omega)} P_n(\omega), \quad \omega \in \Omega_n.$$

Here $P_n(\omega) = \frac{1}{2^n}$, $\beta > 0$ is the **inverse temperature** and Z_n is the normalization constant.

The Gibbs measure captures the competition between the ordering tendency of the energy term $\mathcal{H}(\omega)$ and the randomness represented by P_n . Indeed, let $h = 0$. If the temperature is high (β close to 0), then noise dominates and complete disorder reigns at the limit, $\lim_{\beta \rightarrow 0} \gamma_n(\omega) = P_n$. But if temperature goes to zero, then the limit $\lim_{\beta \rightarrow \infty} \gamma_n(\omega) = \frac{1}{2}(\delta_{\omega=1} + \delta_{\omega=-1})$ is concentrated on the two **ground states**. The key question is the existence of phase transition: namely, if there is a **critical temperature** β_c^{-1} (**Curie point**) at which the infinite model undergoes a transition that reflects the order/disorder dichotomy of the finite model.

Let a random vector (η_1, \dots, η_n) have distribution γ_n . We define magnetization as the expectation $M_n(\beta, h) = \mathbb{E} S_n$ of the total spin $S_n = \sum_{i=1}^n \eta_i$. We show that $\frac{1}{n} S_n$ converges and there exists a limit

$$m(\beta, h) = \lim_{n \rightarrow \infty} \frac{1}{n} M_n(\beta, h).$$

Then we will see something interesting as $h \rightarrow 0$. We remark, that $m(\beta, 0) = 0$, since $\gamma_n(\omega) = \gamma_n(-\omega)$.

Proposition 12.1. *The family $(\frac{1}{n} S_n)_{n \geq 1}$ satisfies the LDP in \mathbb{R} with rate function*

$$I(x) = \begin{cases} \frac{1}{2}(1-x) \ln(1-x) + \frac{1}{2}(1+x) \ln(1+x) - \frac{1}{2} J \beta x^2 - \beta h x - c & \text{if } x \in [-1, 1], \\ +\infty & \text{otherwise,} \end{cases}$$

where $c = \inf_{x \in [-1, 1]} \left\{ \frac{1}{2}(1-x) \ln(1-x) + \frac{1}{2}(1+x) \ln(1+x) - \frac{1}{2} J \beta x^2 - \beta h x \right\}$.

For the proof of the proposition see [RAS15, Section 3.4].

In order to understand limit of $\frac{1}{n} S_n$, we find the minimizers of the rate function I . Critical points satisfy $I'(x) = 0$ that is equivalent to the equation


$$\frac{1}{2} \ln \frac{1+x}{1-x} = J \beta x + \beta h, \quad x \in [-1, 1]. \quad (40)$$

Theorem 12.1. Let $0 < \beta, J < \infty$ and $h \in \mathbb{R}$.

- (i) For $h \neq 0$, $m(\beta, h)$ is the unique solution of (40) that has the same sign as h .
- (ii) Let $h = 0$ and $\beta < J^{-1}$. Then $m(\beta, 0) = 0$ is the unique solution of (40) and $m(\beta, \tilde{h}) \rightarrow 0$ as $\tilde{h} \rightarrow 0$.
- (iii) Let $h = 0$ and $\beta > J^{-1}$. Then (40) has two nonzero solutions $m(\beta, +) > 0$ and $m(\beta, -) = -m(\beta, +)$. Spontaneous magnetization happens: for $\beta > J^{-1} =: \beta_c$,

$$\lim_{\tilde{h} \rightarrow 0+} m(\beta, \tilde{h}) = m(\beta, +) \quad \text{and} \quad \lim_{\tilde{h} \rightarrow 0-} m(\beta, \tilde{h}) = m(\beta, -).$$

We note that statements (i) and (ii) follows directly from the form of equation (40). Statement (iii) is the direct consequence of the further proposition and the dominated convergence theorem.



The graphs of the rate function I . Top plots have $\beta > J^{-1}$ while bottom plots have $\beta \leq J^{-1}$. Top left to right: $h = 0$, $0 < h < h_0(J, \beta)$ and $h > h_0(J, \beta)$. Bottom left to right, $h = 0$ and $h > 0$. The case $h < 0$ is symmetric to that of $h > 0$.

Proposition 12.2. (i) Suppose that either $h \neq 0$, or $h = 0$ and $\beta \leq J^{-1}$. Then $\frac{1}{n}S_n \rightarrow m(\beta, h)$.

(ii) If $h = 0$ and $\beta > J^{-1}$, then $\frac{1}{n}S_n \rightarrow \zeta$ weakly, where $\mathbb{P} \{ \zeta = m(\beta, +) \} = \mathbb{P} \{ \zeta = m(\beta, -) \} = \frac{1}{2}$.

Proof. We note that the first part of the proposition follows from the fact that the rate function I has a unique minimizer. Indeed,

$$\overline{\lim}_{n \rightarrow \infty} \frac{1}{n} \ln \mathbb{P} \left\{ \left| \frac{1}{n}S_n - m(\beta, h) \right| \geq \varepsilon \right\} \leq - \inf_{|x - m(\beta, h)| \geq \varepsilon} I(x) < 0.$$

For part (ii) the large deviation upper bound can be obtained similarly

$$\lim_{n \rightarrow \infty} \mathbb{P} \left\{ \left| \frac{1}{n}S_n - m(\beta, -) \right| < \varepsilon \quad \text{or} \quad \left| \frac{1}{n}S_n - m(\beta, +) \right| < \varepsilon \right\} = 1.$$

Form $\gamma_n(\omega) = \gamma_n(-\omega)$ it follows that S_n is symmetric and so

$$\lim_{n \rightarrow \infty} \mathbb{P} \left\{ \left| \frac{1}{n} S_n - m(\beta, -) \right| < \varepsilon \right\} = \lim_{n \rightarrow \infty} \mathbb{P} \left\{ \left| \frac{1}{n} S_n - m(\beta, +) \right| < \varepsilon \right\} = \frac{1}{2}.$$

This shows the weak convergence of $\frac{1}{n} S_n$ to ζ . □

12.2 Varadhan formula

The goal of the present section is to show a connection of diffusion processes with the underlying geometry of the state space. This result was obtained by Varadhan in [Var67]. So, we are interesting in deviations of solution $x(t)$ of the SDE in \mathbb{R}^d

$$dx(t) = \sigma(x(t))dw(t), \quad x(0) = x_0, \quad (41)$$

from the initial value of x_0 as $t \rightarrow 0$, where $w(t)$, $t \in [0, 1]$, denotes a standard Brownian motion in \mathbb{R}^d and the $d \times d$ -matrix σ is Lipschitz continuous.

We first consider the following family of SDEs

$$dx_\varepsilon(t) = \sigma(x_\varepsilon(t))dw_\varepsilon(t), \quad x(0) = x_0, \quad (42)$$

where $w_\varepsilon(t) = \sqrt{\varepsilon}w(t)$, $t \in [0, 1]$. For every $\varepsilon > 0$ the solution is the diffusion process corresponding to the operator

$$L_\varepsilon(f) = \frac{\varepsilon}{2} \sum_{i,j=1}^d a_{ij} \frac{\partial^2 f}{\partial x_i \partial x_j},$$

with $a = \sigma\sigma^*$. We also assume that the matrix a is bounded and uniformly elliptic, that is, there exists $c > 0$ and $C > 0$ such that for all $\lambda = (\lambda_1, \dots, \lambda_d) \in \mathbb{R}^d$

$$c\|\lambda\|^2 \leq \lambda a \lambda \leq C\|\lambda\|^2,$$

where $\lambda a \lambda = \sum_{i,j=1}^d a_{ij} \lambda_i \lambda_j$. We remark that for every $\varepsilon > 0$ SDE (42) has a unique solution x_ε on the space $C([0, 1], \mathbb{R}^d)$.³⁹ The proof of the following theorem can be found in [Var84, Section 6].

Theorem 12.2. *The family $(x_\varepsilon)_{\varepsilon>0}$ satisfies the LDP in $C([0, 1], \mathbb{R}^d)$ with rate function*

$$I(f) = \begin{cases} \frac{1}{2} \int_0^1 \dot{f}(t) a^{-1}(f(t)) \dot{f}(t) dt & \text{if } f \in H_{x_0}^2([0, 1], \mathbb{R}^d), \\ +\infty & \text{otherwise,} \end{cases}$$

where $H_{x_0}^2([0, T]; \mathbb{R}^d)$ is defined similarly as $H_0^2([0, T]; \mathbb{R}^d)$, the only difference is $f(0) = x_0$.

Now we are going to obtain the LDP for the family $(x(\varepsilon))_{\varepsilon>0}$ where $x(t)$, $t \in [0, 1]$, is a solution to equation (41). It is easily to see that $x(\varepsilon) = x_\varepsilon(1)$. So, we can apply the contraction principle to the LDP for $(x_\varepsilon)_{\varepsilon>0}$. We take the following continuous map on $C([0, T], \mathbb{R}^d)$

$$\Phi(f) = f(1), \quad f \in C([0, T], \mathbb{R}^d).$$

³⁹see e.g. Theorem 21.3 [Kal02]

Then the family $(x(\varepsilon) = \Phi(x_\varepsilon))_{\varepsilon>0}$ satisfies the LDP with rate function

$$\begin{aligned} I_{x_0}(x_1) &= \inf \left\{ I(f) : f \in H_0^2([0, T], \mathbb{R}^d), f(1) = x_1 \right\} \\ &= \frac{1}{2} \inf_{f(0)=x_0, f(1)=x_1} \int_0^1 \dot{f}(t) a^{-1}(f(t)) \dot{f}(t) dt =: \frac{d^2(x_0, x_1)}{2}, \end{aligned}$$

where the later infimum is taken over all functions $f \in H_{x_0}^2([0, 1], \mathbb{R}^d)$ which end at x_1 (and begin at x_0).

Let us define locally the metric on \mathbb{R}^d as

$$ds^2 = \sum_{i,j=1}^d a_{ij} dx_i dx_j.$$

Then the distance

$$d(x_0, x_1) = \left(\inf \left\{ \int_0^1 \dot{f}(t) a^{-1}(f(t)) \dot{f}(t) dt : f \in H_{x_0}^2([0, 1], \mathbb{R}^d), f(1) = 1 \right\} \right)^{\frac{1}{2}}, \quad x_0, x_1 \in \mathbb{R}^d,$$

coincides with the global geodesic distance

$$d_{geod}(x_0, x_1) = \inf \left\{ \int_0^1 \sqrt{\dot{f}(t) a^{-1}(f(t)) \dot{f}(t)} dt : f \in H_{x_0}^2([0, 1], \mathbb{R}^d), f(1) = 1 \right\}, \quad x_0, x_1 \in \mathbb{R}^d,$$

induced by this metric.

Exercise 12.1. Show that d_{geod} is a distance of \mathbb{R}^d .

We remark that the operator L is the Laplace-Beltrami operator on the Riemannian manifold \mathbb{R}^d (with metric ds^2) and the associated process $x(t)$, $t \in [0, 1]$, plays a role of Brownian motion on this space.

For further applications of large deviation principle see also [Var08].

References

- [Con90] John B. Conway, *A course in functional analysis*, second ed., Graduate Texts in Mathematics, vol. 96, Springer-Verlag, New York, 1990. MR 1070713
- [DE97] Paul Dupuis and Richard S. Ellis, *A weak convergence approach to the theory of large deviations*, Wiley Series in Probability and Statistics: Probability and Statistics, John Wiley & Sons, Inc., New York, 1997, A Wiley-Interscience Publication. MR 1431744
- [dH00] Frank den Hollander, *Large deviations*, Fields Institute Monographs, vol. 14, American Mathematical Society, Providence, RI, 2000. MR 1739680
- [DMRYZ04] C. Donati-Martin, A. Rouault, M. Yor, and M. Zani, *Large deviations for squares of Bessel and Ornstein-Uhlenbeck processes*, Probab. Theory Related Fields **129** (2004), no. 2, 261–289. MR 2063378
- [DO10] A. A. Dorogovtsev and O. V. Ostapenko, *Large deviations for flows of interacting Brownian motions*, Stoch. Dyn. **10** (2010), no. 3, 315–339. MR 2671379
- [DS88] Nelson Dunford and Jacob T. Schwartz, *Linear operators. Part I*, Wiley Classics Library, John Wiley & Sons, Inc., New York, 1988, General theory, With the assistance of William G. Bade and Robert G. Bartle, Reprint of the 1958 original, A Wiley-Interscience Publication. MR 1009162
- [DS89] Jean-Dominique Deuschel and Daniel W. Stroock, *Large deviations*, Pure and Applied Mathematics, vol. 137, Academic Press, Inc., Boston, MA, 1989. MR 997938
- [DZ98] Amir Dembo and Ofer Zeitouni, *Large deviations techniques and applications*, second ed., Applications of Mathematics (New York), vol. 38, Springer-Verlag, New York, 1998. MR 1619036
- [EK86] Stewart N. Ethier and Thomas G. Kurtz, *Markov processes*, Wiley Series in Probability and Mathematical Statistics: Probability and Mathematical Statistics, John Wiley & Sons, Inc., New York, 1986, Characterization and convergence. MR 838085
- [Ell06] Richard S. Ellis, *Entropy, large deviations, and statistical mechanics*, Classics in Mathematics, Springer-Verlag, Berlin, 2006, Reprint of the 1985 original. MR 2189669
- [FW84] M. I. Freidlin and A. D. Wentzell, *Random perturbations of dynamical systems*, Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], vol. 260, Springer-Verlag, New York, 1984, Translated from the Russian by Joseph Szücs. MR 722136
- [Kal02] Olav Kallenberg, *Foundations of modern probability*, second ed., Probability and its Applications (New York), Springer-Verlag, New York, 2002. MR 1876169
- [Kuz15] V. A. Kuznetsov, *On the large-deviation principle for the winding angle of a Brownian trajectory around the origin*, Theory Stoch. Process. **20** (2015), no. 2, 63–84. MR 3510229
- [KvR19] Vitalii Konarovskiy and Max-K. von Renesse, *Modified massive Arratia flow and Wasserstein diffusion*, Comm. Pure Appl. Math. **72** (2019), no. 4, 764–800. MR 3914882

- [LS87] James Lynch and Jayaram Sethuraman, *Large deviations for processes with independent increments*, Ann. Probab. **15** (1987), no. 2, 610–627. MR 885133
- [Mö8] Peter Mörters, *Large deviation theory and applications*, Lecture notes, 2008.
- [RAS15] Firas Rassoul-Agha and Timo Seppäläinen, *A course on large deviations with an introduction to Gibbs measures*, Graduate Studies in Mathematics, vol. 162, American Mathematical Society, Providence, RI, 2015. MR 3309619
- [Sch97] Alexander Schied, *Moderate deviations and functional LIL for super-Brownian motion*, Stochastic Process. Appl. **72** (1997), no. 1, 11–25. MR 1483609
- [Swa12] Jan Swart, *Large deviation theory*, Lecture notes, 2012.
- [Var67] S. R. S. Varadhan, *On the behavior of the fundamental solution of the heat equation with variable coefficients*, Comm. Pure Appl. Math. **20** (1967), 431–455. MR 0208191
- [Var84] ———, *Large deviations and applications*, CBMS-NSF Regional Conference Series in Applied Mathematics, vol. 46, Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, 1984. MR 758258
- [Var08] ———, *Large deviations*, Ann. Probab. **36** (2008), no. 2, 397–419. MR 2393987