

SYMPLECTIC GEOMETRY

Problem Set 6

1. For a function $a : \mathbb{R}^4 \rightarrow \mathbb{R}$, we consider the almost complex structure J_a on the manifold $M = \mathbb{R}^4$ which in the global coordinates (x_1, x_2, y_1, y_2) has the form

$$J_a(p) = \begin{pmatrix} 0 & 0 & -1 & 0 \\ a(p) & 0 & 0 & -1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & -a(p) & 0 \end{pmatrix}, \text{ i.e. } J_a \left(\frac{\partial}{\partial x_1} \right) = a(p) \frac{\partial}{\partial x_2} + \frac{\partial}{\partial y_1} \text{ etc.}$$

- a) Prove that if $|a(p)| \leq 1$ for all $p \in \mathbb{R}^4$, then J_a is tamed by the standard symplectic form $\omega_{\text{st}} = dx_1 \wedge dy_1 + dx_2 \wedge dy_2$!

Hint: Recall that the taming condition means that $\omega(v, Jv) > 0$ for all non-zero v , but ω need not be J -invariant, so that the bilinear form $\omega(\cdot, J\cdot)$ need not be symmetric.

- b) Under which conditions on the function a is the almost complex structure J_a on \mathbb{R}^4 integrable?

Hint: Argue that in order to determine N_{J_a} on any two vectors $v, w \in T_p\mathbb{R}^4$, it suffices to know $N_{J_a} \left(\frac{\partial}{\partial x_1}(p), \frac{\partial}{\partial x_2}(p) \right)$, and then compute this.

2. In the lecture, we introduced the Fubini-Study form ω_{FS} on $\mathbb{C}P^n$ as the unique 2-form satisfying

$$\pi^* \omega_{\text{FS}} = \iota^* \omega_{\text{st}},$$

where $\pi : S^{2n+1} \rightarrow \mathbb{C}P^n$ and $\iota : S^{2n+1} \rightarrow \mathbb{C}^{n+1}$ are the standard projection and the standard embedding.

- a) Prove that the form

$$\tilde{\omega} := \frac{1}{\|z\|^2} \cdot \omega_{\text{st}}$$

on $\mathbb{C}^{n+1} \setminus \{0\}$ is invariant under the \mathbb{C}^* -action by rescaling, and notice that $\iota^* \tilde{\omega} = \iota^* \omega_{\text{st}}$.

- b) Recall that we defined¹ $H_z := \text{span}_{\mathbb{C}}(z)^\perp \subset T_z\mathbb{C}^{n+1}$. Now use a coordinate expression for the orthogonal projection $T_z\mathbb{C}^{n+1} \rightarrow H_z$ and part **a**) to prove that in homogeneous coordinates $[z_0 : \dots : z_n]$ on $\mathbb{C}P^n$ the Fubini-Study form is given by

$$\omega_{\text{FS}} = \frac{i}{2} \left(\sum_j \frac{dz_j \wedge d\bar{z}_j}{\|z\|^2} - \sum_{j,k} \frac{\bar{z}_j z_k dz_j \wedge d\bar{z}_k}{\|z\|^4} \right).$$

- c) The open sets $U_i = \{[z_0 : \dots : z_n] \in \mathbb{C}P^n \mid z_i \neq 0\}$ cover $\mathbb{C}P^n$. Prove that the maps

$$\begin{aligned} \psi_i : U_i &\rightarrow \mathbb{C}^n \\ [z_0 : \dots : z_n] &\mapsto \left(\frac{z_0}{z_i}, \dots, \widehat{\frac{z_i}{z_i}}, \dots, \frac{z_n}{z_i} \right) \end{aligned}$$

give complex charts for $\mathbb{C}P^n$, i.e. the transition maps are holomorphic. What is $\mathbb{C}P^n \setminus U_i$ diffeomorphic to? What is $U_i \cap U_j$ diffeomorphic to?

- d) Derive an expression for the Fubini-Study form in inhomogeneous coordinates $\zeta_j := \frac{z_j}{z_0}$ on the open set $U_0 \subset \mathbb{C}P^n$.
- e) Compute an explicit expression for $\eta = \frac{i}{2} \partial \bar{\partial} (\log(1 + \|\zeta\|^2))$ on \mathbb{C}^n and compare with the result for $(\psi_0^{-1})^* \omega_{\text{FS}}$ you obtained in **d**).
- f) Compute the integral

$$\int_{\mathbb{C}P^1} \omega_{\text{FS}}.$$

¹In class we did this for $z \in S^{2n+1}$, but the same definition works in general.