

SQUARES, ASCENT PATHS, AND CHAIN CONDITIONS

CHRIS LAMBIE-HANSON AND PHILIPP LÜCKE

ABSTRACT. With the help of various square principles, we obtain results concerning the consistency strength of several statements about trees containing ascent paths, special trees, and strong chain conditions. Building on a result that shows that Todorčević's principle $\square(\kappa)$ implies an indexed version of $\square(\kappa, \lambda)$, we show that for all infinite, regular cardinals $\lambda < \kappa$, the principle $\square(\kappa)$ implies the existence of a κ -Aronszajn tree containing a λ -ascent path. We then provide a complete picture of the consistency strengths of statements relating the interactions of trees with ascent paths and special trees. As a part of this analysis, we construct a model of set theory in which \aleph_2 -Aronszajn trees exist and all such trees contain \aleph_0 -ascent paths. Finally, we use our techniques to show that the assumption that the κ -Knaster property is countably productive and the assumption that every κ -Knaster partial order is κ -stationarily layered both imply the failure of $\square(\kappa)$.

1. INTRODUCTION

The existence or non-existence of cofinal branches is one of the most fundamental properties of set-theoretic trees¹ of uncountable regular height. Important examples of trees without cofinal branches are given by *special trees*. Given an infinite cardinal μ , a tree of height μ^+ is special if it can be decomposed into μ -many antichains. This notion was generalized by Todorčević to the class of all trees of uncountable regular heights (see Definition 2.3). It is easy to see that a special tree does not contain a cofinal branch, not only in the ground model V , but also in all outer models of V in which its height remains a regular cardinal.

In contrast, it is possible to use the concept of *ascent paths*, introduced by Laver, to obtain interesting examples of branchless, non-special trees of uncountable regular height. Given infinite regular cardinals $\lambda < \kappa$, a λ -ascent path through a tree \mathbb{T} of height κ is a sequence $\langle b_\alpha : \lambda \longrightarrow \mathbb{T} \mid \alpha < \kappa \rangle$ of functions with the property

2010 *Mathematics Subject Classification*. Primary 03E05; Secondary 03E35, 03E55.

Key words and phrases. Square principles, ascent paths, special trees, productivity of chain conditions, Knaster property, layered posets, walks on ordinals.

During the preparation of this paper, the first author was partially supported by the Israel Science Foundation under the grant #1630/14, and the second author was partially supported by the Deutsche Forschungsgemeinschaft under the grant LU2020/1-1. The initial results of this paper were obtained while the authors were participating in the *Intensive Research Program on Large Cardinals and Strong Logics* at the Centre de Recerca Matemàtica in Barcelona during the fall of 2016. The authors would like to thank the organizers for the opportunity to participate in the program. Further results were obtained while the first author was visiting the second author in Bonn during the spring of 2017. The first author would like to thank the Deutsche Forschungsgemeinschaft for the financial support of this visit through the above grant. Finally, the authors would like to thank the anonymous referee for the careful reading of the manuscript and several helpful comments.

¹A summary of basic definitions concerning set-theoretic trees can be found in Section 2.

that $b_\alpha(i)$ is contained in the α -th level of \mathbb{T} for all $\alpha < \kappa$ and $i < \lambda$ and, for all $\alpha < \beta < \kappa$, there is an $i < \lambda$ with $b_\alpha(j) <_{\mathbb{T}} b_\beta(j)$ for all $i \leq j < \lambda$. A theorem of Shelah (see [22, Lemma 3]) then shows that, if μ is an uncountable cardinal and $\lambda < \mu$ is a regular cardinal with $\lambda \neq \text{cof}(\mu)$, then every tree of height μ^+ that contains a λ -ascent path is not special. Note that this shows that trees containing ascent paths are non-special in a very absolute way, because it implies that they remain non-special in every outer model of V in which μ and μ^+ remain cardinals and $\text{cof}(\lambda) \neq \text{cof}(\mu)$ holds. This result was later strengthened by Todorćević and Torres Pérez in [27, Proposition 2.3] and by the second author in [18, Lemma 1.6] (see Lemma 2.6 below).

Many authors have dealt with the construction of trees of various types containing ascending paths (see, for example, [2], [5], [9], [15], [18], [22] and [26]). In particular, the constructions of Shelah and Stanley in [22] and Todorćević in [26] show that, given infinite, regular cardinals $\lambda < \kappa$, the existence of a κ -Aronszajn tree containing a λ -ascent path follows from the existence of a $\square(\kappa)$ -sequence that avoids² a stationary subset S of κ consisting of limit ordinals of cofinality λ (see [18, Theorem 4.12] and [21, Section 3]). Our first main result shows that such a tree can be constructed from a $\square(\kappa)$ -sequence without additional properties. This answers [18, Questions 6.5 and 6.6].

Theorem 1.1. *Let $\lambda < \kappa$ be infinite, regular cardinals. If $\square(\kappa)$ holds, then there is a κ -Aronszajn tree with a λ -ascent path.*

It is easy to see that, if κ is a weakly compact cardinal and \mathbb{T} is tree of height κ containing a λ -ascent path with $\lambda < \kappa$, then \mathbb{T} contains a cofinal branch. Moreover, basic arguments, presented in [18, Section 3], show that, if κ is a weakly compact cardinal, $\mu < \kappa$ is a regular, uncountable cardinal, and G is $\text{Col}(\mu, < \kappa)$ -generic over V , then every tree of height κ in $V[G]$ that contains a λ -ascent path with $\lambda < \mu$ already has a cofinal branch. Since seminal results of Jensen and Todorćević show that, for uncountable regular cardinals κ , a failure of $\square(\kappa)$ implies that κ is weakly compact in Gödel's constructible universe L (see [11, Section 6] and [25, (1.10)]), the above theorem directly yields the following corollary showing that the existence of regular cardinals $\lambda < \mu$ such that there are no μ^+ -Aronszajn trees with λ -ascent paths is equiconsistent with the existence of a weakly compact cardinal.

Corollary 1.2. *Let κ be an uncountable regular cardinal. If there is an infinite regular cardinal $\lambda < \kappa$ with the property that there are no κ -Aronszajn trees with λ -ascent paths, then κ is a weakly compact cardinal in L . \square*

Starting with the above theorem, we provide a complete picture of the consistency strengths of statements relating the interactions of trees with ascent paths and special trees. For concreteness, we will speak here about \aleph_2 -Aronszajn trees and \aleph_0 -ascent paths, but the same results will hold for μ^+ -Aronszajn trees and λ -ascent paths, provided $\lambda < \mu$ are infinite, regular cardinals. In what follows, if \mathbb{T} is an \aleph_2 -Aronszajn tree, then $S(\mathbb{T})$ denotes the assertion that \mathbb{T} is special and $A(\mathbb{T})$ denotes the assertion that \mathbb{T} has an \aleph_0 -ascent path. Table 1 provides a complete picture of the precise consistency strengths of various assertions relating the existence of special trees and the existence of trees with ascent paths, where the

²For a set A of ordinals, we define $\text{acc}(A) = \{\alpha \in A \mid \sup(A \cap \alpha) = \alpha > 0\}$. Given an uncountable regular cardinal κ , a $\square(\kappa)$ -sequence $\langle C_\alpha \mid \alpha < \kappa \rangle$ avoids a stationary subset S of κ if $\text{acc}(C_\alpha) \cap S = \emptyset$ holds for all $\alpha \in \text{acc}(\kappa)$.

background assumption is that there are \aleph_2 -Aronszajn trees, and the quantification is over the set of \aleph_2 -Aronszajn trees.

	$\forall \mathbb{T}.A(\mathbb{T})$	$\exists \mathbb{T}.A(\mathbb{T})$	$\forall \mathbb{T}.\neg A(\mathbb{T})$
$\forall \mathbb{T}.S(\mathbb{T})$	$0 = 1$	$0 = 1$	Weakly compact
$\exists \mathbb{T}.S(\mathbb{T})$	$0 = 1$	ZFC	Weakly compact
$\forall \mathbb{T}.\neg S(\mathbb{T})$	Weakly compact	Mahlo	Weakly compact

TABLE 1. The consistency strengths of interactions between trees with ascent paths and special trees.

Besides Theorem 1.1, the following result is the other main new ingredient in the determination of the consistency strengths in Table 1. The results of Section 4 will show that, for successors of regular cardinals and inaccessible cardinals, the consistency of the hypotheses of this theorem can be established from a weakly compact cardinal. Given an infinite, regular cardinal κ , we let $\text{Add}(\kappa, 1)$ denote the partial order that adds a Cohen subset to κ . Moreover, given an uncountable, regular cardinal κ , we let $\text{TP}(\kappa)$ denote the statement that the *tree property* holds at κ , i.e. the statement that there are no κ -Aronszajn trees.

Theorem 1.3. *Let $\lambda < \kappa$ be infinite, regular cardinals such that $\kappa = \kappa^{<\kappa}$ and $\mathbb{1}_{\text{Add}(\kappa, 1)} \Vdash \text{TP}(\check{\kappa})$. Then the following statements hold in a cofinality-preserving forcing extension of the ground model:*

- (1) *There are κ -Aronszajn trees.*
- (2) *Every κ -Aronszajn tree contains a λ -ascent path.*

In the last part of this paper, we use the techniques developed in it to study chain conditions of partial orders. There is a close connection between ascent paths and the infinite productivity of chain conditions, given by the fact that a result of Baumgartner (see [1, Theorem 8.2]) shows that for every tree \mathbb{T} of uncountable regular height κ without cofinal branches, the canonical partial order $\mathbb{P}(\mathbb{T})$ that specializes \mathbb{T} using finite partial functions $f : \mathbb{T} \xrightarrow{\text{part}} \omega$ (see Definition 7.3) satisfies the κ -chain condition, and that every λ -ascent path $\langle b_\alpha : \lambda \rightarrow \mathbb{T} \mid \alpha < \kappa \rangle$ through \mathbb{T} induces an antichain $\{p_\alpha \mid \alpha < \kappa\}$ in the full support product $\prod_{i < \lambda} \mathbb{P}(\mathbb{T})$ with $\text{dom}(p_\alpha(i)) = \{b_\alpha(i)\}$ and $p_\alpha(i)(b_\alpha(i)) = 0$ for all $\alpha < \kappa$ and $i < \lambda$ (see [18, Section 2] for more details on this connection).

Our first application deals with failures of the infinite productivity of the κ -Knaster property. Remember that, given an uncountable regular cardinal κ , a partial order \mathbb{P} is κ -Knaster if every collection of κ -many conditions in \mathbb{P} contains a subcollection of cardinality κ that consists of pairwise compatible conditions. This strengthening of the κ -chain condition is of great interest because of its product behavior. In particular, the product of two κ -Knaster partial orders is κ -Knaster and the product of a κ -Knaster partial order with a partial order satisfying the κ -chain condition again satisfies the κ -chain condition. An easy argument (see [4, Proposition 1.1]) shows that, if κ is a weakly compact cardinal, then the class of κ -Knaster partial orders is closed under μ -support products for all $\mu < \kappa$. A combination of [4, Theorem 1.13] with [18, Theorem 1.12] shows that the question of whether, for uncountable regular cardinals κ , the countable productivity of the κ -Knaster is equivalent to the weak compactness of κ is independent of the axioms

of ZFC. The construction in [4], producing a model of set theory in which this characterization of weak compactness fails, starts from a model of ZFC containing a weakly compact cardinal. The following result and its corollary show that this assumption is necessary.

Theorem 1.4. *Let κ be an uncountable regular cardinal with the property that $\square(\kappa)$ holds. If $\lambda < \kappa$ is an infinite, regular cardinal, then there is a partial order \mathbb{P} with the following properties:*

- (1) *If $\mu < \lambda$ is a (possibly finite) cardinal with $\nu^\mu < \kappa$ for all $\nu < \kappa$, then \mathbb{P}^μ is κ -Knaster.*
- (2) *\mathbb{P}^λ does not satisfy the κ -chain condition.*

The results of [4] and [18] leave open the question whether it is consistent that the κ -Knaster property is countably productive for accessible uncountable regular cardinals, like \aleph_2 . It is easy to see that this productivity implies certain cardinal arithmetic statements. Namely, if $\lambda < \kappa$ are infinite, regular cardinals, $\nu < \kappa$ is a cardinal with $\nu^\lambda \geq \kappa$ and \mathbb{P} is a partial order of cardinality ν containing an antichain of size ν (e.g. the lottery sum of ν -many copies of *Cohen forcing* $\text{Add}(\omega, 1)$), then \mathbb{P} is κ -Knaster and the full support product \mathbb{P}^λ contains an antichain of size ν^λ .

Corollary 1.5. *Let $\lambda < \kappa$ be infinite, regular cardinals. If the class of κ -Knaster partial orders is closed under λ -support products, then κ is weakly compact in \mathbb{L} and $\nu^\lambda < \kappa$ for all $\nu < \kappa$. \square*

Our second application deals with a strengthening of the κ -Knaster property introduced by Cox in [3]. Given an uncountable regular cardinal κ , a partial order \mathbb{P} is κ -stationarily layered if the collection of all regular suborders³ of \mathbb{P} of cardinality less than κ is stationary⁴ in the collection $\mathcal{P}_\kappa(\mathbb{P})$ of all subsets of \mathbb{P} of cardinality less than κ . In [3], Cox shows that this property implies the κ -Knaster property. The main result of [4] shows that an uncountable regular cardinal is weakly compact if and only if every partial order satisfying the κ -chain condition is κ -stationarily layered. Moreover, it is shown that the assumption that every κ -Knaster partial order is κ -stationarily layered implies that κ is a Mahlo cardinal with the property that every stationary subset of κ reflects. In particular, it follows that this assumption characterizes weak compactness in certain models of set theory. In contrast, it is also shown in [4] that there is consistently a non-weakly compact cardinal κ such that every κ -Knaster partial order is κ -stationarily layered. The model of set theory witnessing this consistency is again constructed assuming the existence of a weakly compact cardinal. The following result shows that this assumption is necessary, answering [4, Questions 7.1 and 7.2].

Theorem 1.6. *Let κ be an uncountable, regular cardinal. If every κ -Knaster partial order is κ -stationarily layered, then $\square(\kappa)$ fails.*

2. TREES AND ASCENT PATHS

In this short section, we recall some fundamental definitions and results dealing with trees, special trees and ascent paths.

³Given a partial order \mathbb{P} , a suborder \mathbb{Q} of \mathbb{P} is a *regular suborder* if the inclusion map preserves incompatibility and maximal antichains in \mathbb{Q} are maximal in \mathbb{P} .

⁴This definition refers to Jech's notion of stationarity in $\mathcal{P}_\kappa(A)$: a subset of $\mathcal{P}_\kappa(A)$ is stationary in $\mathcal{P}_\kappa(A)$ if it meets every subset of $\mathcal{P}_\kappa(A)$ which is \subseteq -continuous and cofinal in $\mathcal{P}_\kappa(A)$.

Definition 2.1. A partial order \mathbb{T} is a *tree* if, for all $t \in \mathbb{T}$, the set

$$\text{pred}_{\mathbb{T}}(t) = \{s \in \mathbb{T} \mid s <_{\mathbb{T}} t\}$$

is well-ordered by the relation $<_{\mathbb{T}}$.

Definition 2.2. Let \mathbb{T} be a tree.

- (1) For all $t \in \mathbb{T}$, we let $\text{ht}_{\mathbb{T}}(t)$ denotes the order type of $\langle \text{pred}_{\mathbb{T}}(t), <_{\mathbb{T}} \rangle$.
- (2) For all $\alpha \in \text{On}$, we set $\mathbb{T}(\alpha) = \{t \in \mathbb{T} \mid \text{ht}_{\mathbb{T}}(t) = \alpha\}$.
- (3) We let $\text{ht}(\mathbb{T})$ denote the least ordinal α such that $\mathbb{T}(\alpha) = \emptyset$. This ordinal is referred to as the *height* of \mathbb{T} .
- (4) If $S \subseteq \text{ht}(\mathbb{T})$, then $\mathbb{T} \upharpoonright S$ is the suborder of \mathbb{T} whose underlying set is $\bigcup \{\mathbb{T}(\alpha) \mid \alpha \in S\}$.
- (5) A *branch through* \mathbb{T} is a subset B of \mathbb{T} that is linearly ordered by $\leq_{\mathbb{T}}$. A branch B is *cofinal* if the set $\{\text{ht}_{\mathbb{T}}(t) \mid t \in B\}$ is cofinal in $\text{ht}(\mathbb{T})$.
- (6) If κ is a regular cardinal, then \mathbb{T} is a κ -*tree* if $\text{ht}(\mathbb{T}) = \kappa$ and $\mathbb{T}(\alpha)$ has cardinality less than κ for all $\alpha < \kappa$. A κ -*Aronszajn tree* is a κ -tree without cofinal branches.

The concept defined next was introduced by Todorćević in [23] for the case $\kappa = \aleph_1$ and later generalized to arbitrary uncountable regular cardinals in [24].

Definition 2.3 (Todorćević). Let κ be an uncountable regular cardinal, let \mathbb{T} be a tree of height κ , and let S be a subset of κ .

- (1) A map $r : \mathbb{T} \upharpoonright S \rightarrow \mathbb{T}$ is *regressive* if $r(t) <_{\mathbb{T}} t$ holds for all $t \in \mathbb{T} \upharpoonright S$ with $\text{ht}_{\mathbb{T}}(t) > 0$.
- (2) The subset S is *non-stationary with respect to* \mathbb{T} if there is a regressive map $r : \mathbb{T} \upharpoonright S \rightarrow \mathbb{T}$ such that, for every $t \in \mathbb{T}$, there is a $\theta_t < \kappa$ and a function $c_t : r^{-1}(\theta_t) \rightarrow \theta_t$ that is injective on $<_{\mathbb{T}}$ -chains.
- (3) The tree \mathbb{T} is *special* if κ is non-stationary with respect to \mathbb{T} .

A result of Todorćević (see [24, Theorem 14]) shows that for successor cardinals, the above notion of special trees coincides with the notion mentioned in Section 1. One of the reasons for interest in special trees is that they are branchless in a very absolute way.

Fact 2.4. *Suppose that κ is an uncountable regular cardinal, \mathbb{T} is a tree of height κ , and there is a stationary $S \subseteq \kappa$ such that S is non-stationary with respect to \mathbb{T} . Then there are no cofinal branches through \mathbb{T} . \square*

Definition 2.5. Let $\lambda < \kappa$ be cardinals with κ uncountable and regular, let \mathbb{T} be a tree of height κ , and let $\vec{b} = \langle b_{\alpha} : \lambda \rightarrow \mathbb{T}(\alpha) \mid \alpha < \kappa \rangle$ be a sequence of functions.

- (1) The sequence \vec{b} is a λ -*ascending path through* \mathbb{T} if, for all $\alpha < \beta < \kappa$, there are $i, j < \lambda$ with $b_{\alpha}(i) <_{\mathbb{T}} b_{\beta}(j)$.
- (2) The sequence \vec{b} is a λ -*ascent path through* \mathbb{T} if, for all $\alpha < \beta < \kappa$, there is an $i < \lambda$ such that $b_{\alpha}(j) <_{\mathbb{T}} b_{\beta}(j)$ holds for all $i \leq j < \lambda$.
- (3) Suppose that \vec{b} is a λ -ascent path through \mathbb{T} . Given $I < \lambda$ and a cofinal subset B of κ , the pair $\langle I, B \rangle$ is a *true cofinal branch* through \vec{b} if the following statements hold:
 - (a) If $\alpha, \beta \in B$ with $\alpha < \beta$, then $b_{\alpha}(i) <_{\mathbb{T}} b_{\beta}(i)$ for all $I \leq i < \lambda$.
 - (b) If $\beta \in B$ and $\alpha < \beta$ with $b_{\alpha}(i) <_{\mathbb{T}} b_{\beta}(i)$ for all $I \leq i < \lambda$, then $\alpha \in B$.

A λ -ascent path through a tree \mathbb{T} is clearly a λ -ascending path through \mathbb{T} . The notion of a λ -ascent path is due to Laver and grew out of his work on higher Souslin Hypotheses in [17]. Ascent paths and ascending paths can be seen as generalized cofinal branches and, like cofinal branches, they provide concrete obstructions to a tree being special. The best current result in this direction is due to the second author, building upon work of Shelah in [22] and Todorćević and Torres Pérez in [27]. Given an uncountable regular cardinal κ and a cardinal $\lambda < \kappa$, we let $E_{>\lambda}^\kappa$ denote the set of all $\alpha \in \text{acc}(\kappa)$ with $\text{cof}(\alpha) > \lambda$. The sets $E_{\geq\lambda}^\kappa$, E_λ^κ , etc. are defined analogously.

Lemma 2.6 ([18, Lemma 1.6]). *Let $\lambda < \kappa$ be infinite cardinals with κ uncountable and regular, let \mathbb{T} be a tree of height κ and let $S \subseteq S_{>\lambda}^\kappa$ be stationary in κ . If κ is not the successor of a cardinal of cofinality at most λ and the set S is non-stationary with respect to \mathbb{T} , then there are no λ -ascending paths through \mathbb{T} .*

In particular, if κ is either weakly inaccessible or the successor of a regular cardinal and λ is a cardinal with $\lambda^+ < \kappa$, then special trees of height κ do not contain λ -ascending paths. In contrast, the first author showed in [15] that, if λ is a singular cardinal, then Jensen's principle \square_λ implies the existence of a special tree of height λ^+ containing a $\text{cof}(\lambda)$ -ascent path.

3. SQUARE PRINCIPLES

In the following, we recall the definitions of several square principles that will be used in this paper.

Definition 3.1. Given an uncountable regular cardinal κ and a cardinal $1 < \lambda \leq \kappa$, a sequence $\langle \mathcal{C}_\alpha \mid \alpha < \kappa \rangle$ is a $\square(\kappa, < \lambda)$ -sequence if the following statements hold:

- (1) For all $\alpha \in \text{acc}(\kappa)$, \mathcal{C}_α is a collection of club subsets of α with $0 < |\mathcal{C}_\alpha| < \lambda$.
- (2) If $\alpha, \beta \in \text{acc}(\kappa)$, $C \in \mathcal{C}_\beta$, and $\alpha \in \text{acc}(C)$, then $C \cap \alpha \in \mathcal{C}_\alpha$.
- (3) There is no club D in κ such that, for all $\alpha \in \text{acc}(D)$, we have $D \cap \alpha \in \mathcal{C}_\alpha$.

We let $\square(\kappa, < \lambda)$ denote the assertion that there is a $\square(\kappa, < \lambda)$ -sequence. The principle $\square(\kappa, < \lambda^+)$ is typically written as $\square(\kappa, \lambda)$, and $\square(\kappa, 1)$ is written as $\square(\kappa)$. Finally, a sequence $\langle C_\alpha \mid \alpha < \kappa \rangle$ is a $\square(\kappa)$ -sequence if the sequence $\{\langle C_\alpha \rangle \mid \alpha < \kappa\}$ witnesses that $\square(\kappa)$ holds.

We next introduce an indexed version of $\square(\kappa, \lambda)$. The definition is taken from [16] and is a modification of similar indexed square notions studied in [7] and [8].

Definition 3.2. Let $\lambda < \kappa$ be infinite regular cardinals. A $\square^{\text{ind}}(\kappa, \lambda)$ -sequence is a matrix $\vec{C} = \langle C_{\alpha,i} \mid \alpha < \kappa, i(\alpha) \leq i < \lambda \rangle$ satisfying the following statements:

- (1) If $\alpha \in \text{acc}(\kappa)$, then $i(\alpha) < \lambda$.
- (2) If $\alpha \in \text{acc}(\kappa)$ and $i(\alpha) \leq i < \lambda$, then $C_{\alpha,i}$ is a club subset of α .
- (3) If $\alpha \in \text{acc}(\kappa)$ and $i(\alpha) \leq i < j < \lambda$, then $C_{\alpha,i} \subseteq C_{\alpha,j}$.
- (4) If $\alpha, \beta \in \text{acc}(\kappa)$ and $i(\beta) \leq i < \lambda$, then $\alpha \in \text{acc}(C_{\beta,i})$ implies that $i \geq i(\alpha)$ and $C_{\alpha,i} = C_{\beta,i} \cap \alpha$.
- (5) If $\alpha, \beta \in \text{acc}(\kappa)$ with $\alpha < \beta$, then there is an $i(\beta) \leq i < \lambda$ such that $\alpha \in \text{acc}(C_{\beta,i})$.
- (6) There is no club subset D of κ such that, for all $\alpha \in \text{acc}(D)$, there is $i < \lambda$ such that $C_{\alpha,i} = D \cap \alpha$ holds.

We let $\square^{\text{ind}}(\kappa, \lambda)$ denote the assertion that there is a $\square^{\text{ind}}(\kappa, \lambda)$ -sequence.

A proof of the following statement can be found in [16, Section 6].

Proposition 3.3. *Definition 3.2 is unchanged if we replace condition (6) by the following seemingly weaker condition:*

- (6') *There is no club subset D of κ and $i < \lambda$ such that $i \geq i(\alpha)$ and $C_{\alpha,i} = D \cap \alpha$ for all $\alpha \in \text{acc}(D)$.*

It is immediate that the principle $\square^{\text{ind}}(\kappa, \lambda)$ implies $\square(\kappa, \lambda)$. We next show that $\square(\kappa)$ implies all relevant instances of $\square^{\text{ind}}(\kappa, \lambda)$.

Theorem 3.4. *Let $\lambda < \kappa$ be infinite regular cardinals and assume that $\square(\kappa)$ holds. Given a stationary subset S of κ , there is a $\square^{\text{ind}}(\kappa, \lambda)$ -sequence*

$$\vec{C} = \langle C_{\alpha,i} \mid \alpha \in \text{acc}(\kappa), i(\alpha) \leq i < \lambda \rangle$$

with the following properties:

- (1) *If $i < \lambda$, then the set $\{\alpha \in S \mid i(\alpha) = i\}$ is stationary in κ .*
- (2) *There is a $\square(\kappa)$ -sequence $\langle D_\alpha \mid \alpha < \kappa \rangle$ such that $\text{acc}(D_\alpha) \subseteq \text{acc}(C_{\alpha,i(\alpha)})$ holds for all $\alpha \in \text{acc}(\kappa)$.*

Proof. By a result of Rinot (see [20, Lemma 3.2]) and our assumptions, there is a $\square(\kappa)$ -sequence $\vec{D} = \langle D_\alpha \mid \alpha < \kappa \rangle$ with the property that, for all $\eta < \kappa$, the set $S^\eta = \{\alpha \in S \mid \min(D_\alpha) = \eta\}$ is stationary. Fix a partition $\langle A_i \mid i < \lambda \rangle$ of κ into disjoint, non-empty sets. Given $\alpha \in \text{acc}(\kappa)$, define $i(\alpha)$ to be the unique $i < \lambda$ with $\min(D_\alpha) \in A_i$. Then the set $\{\alpha \in S \mid i(\alpha) = i\}$ is stationary in κ for every $i < \lambda$ and, if $\alpha \in \text{acc}(\kappa)$ and $\beta \in \text{acc}(D_\alpha)$, then $i(\alpha) = i(\beta)$.

By induction on $\alpha \in \text{acc}(\kappa)$, we define a matrix $\langle C_{\alpha,i} \mid \alpha \in \text{acc}(\kappa), i(\alpha) < i < \lambda \rangle$ satisfying clauses (1)–(5) listed in Definition 3.2 together with the assumption that $\text{acc}(D_\alpha) \subseteq \text{acc}(C_{\alpha,i(\alpha)})$ holds for every $\alpha \in \text{acc}(\kappa)$. In the following, fix a limit ordinal $\alpha < \kappa$ and assume that we already have constructed a matrix with the above properties up to α . There are a number of cases to consider:

Case 1: $\alpha = \omega$. Define $C_{\omega,i} = \omega$ for all $i(\omega) \leq i < \lambda$. Then all of the desired requirements are trivially satisfied.

Case 2a: $\alpha = \beta + \omega$ for a limit ordinal β , and $\text{acc}(D_\alpha) = \emptyset$. Set $j = \max\{i(\alpha), i(\beta)\}$, $C_{\alpha,i} = \{\beta + n \mid n < \omega\}$ for all $i(\alpha) \leq i < j$ and $C_{\alpha,i} = C_{\beta,i} \cup \{\beta + n \mid n < \omega\}$ for all $j \leq i < \lambda$. Then it is easy to see that clauses (1)–(3) and (5) of Definition 3.2 hold. To verify clause (4), suppose $i(\alpha) \leq i < \lambda$ and $\gamma \in \text{acc}(C_{\alpha,i})$. By our construction, it follows that $i \geq j \geq i(\beta)$ and $\gamma \in \text{acc}(C_{\beta,i}) \cup \{\beta\}$. By the induction hypothesis applied to β , it follows that $i \geq i(\gamma)$ and $C_{\gamma,i} = C_{\beta,i} \cap \gamma = C_{\alpha,i} \cap \gamma$.

Case 2b: $\alpha = \beta + \omega$ for a limit ordinal β , and $\text{acc}(D_\alpha) \neq \emptyset$. In the following, we set $\alpha_0 = \max(\text{acc}(D_\alpha)) \leq \beta$. Then the above remarks show that $i(\alpha_0) = i(\alpha)$. If $\alpha_0 < \beta$, then we let j be minimal such that $i(\alpha) \leq j < \lambda$ and $\alpha_0 \in \text{acc}(C_{\beta,j})$. Otherwise, we set $j = i(\alpha)$. Then $j \geq i(\beta)$, because either $\alpha_0 = \beta$, $\beta \in \text{acc}(D_\alpha)$ and $i(\beta) = i(\alpha_0) = j$ or $\alpha_0 < \beta$, $\alpha_0 \in \text{acc}(C_{\beta,j})$ and $j \geq i(\beta)$. Define

$$C_{\alpha,i} = C_{\alpha_0,i} \cup \{\alpha_0\} \cup \{\beta + n \mid n < \omega\}$$

for all $i(\alpha) \leq i < j$ and $C_{\alpha,i} = C_{\beta,i} \cup \{\beta + n \mid n < \omega\}$ for all $j \leq i < \lambda$.

Since our induction hypothesis ensures that $C_{\alpha_0,i} \cup \{\alpha_0\} \subseteq C_{\alpha_0,j} \cup \{\alpha_0\} \subseteq C_{\beta,j}$ holds for all $i(\alpha) \leq i < j$, it is easy to see that Clauses (1)–(3) and (5) from Definition 3.2 hold in this case. We thus verify clause (4). Suppose $i(\alpha) \leq i < \lambda$ and $\gamma \in \text{acc}(C_{\alpha,i})$. If $i(\alpha) \leq i < j$, then either $\gamma = \alpha_0$ or $\gamma \in \text{acc}(C_{\alpha_0,i})$. In both

instances, we have $i(\gamma) \leq i$ and $C_{\gamma,i} = C_{\alpha_0,i} \cap \gamma = C_{\alpha,i} \cap \gamma$. On the other hand, if $j \leq i < \lambda$, then $\gamma \in \text{acc}(C_{\beta,i}) \cup \{\beta\}$, so $i \geq i(\gamma)$ and $C_{\gamma,i} = C_{\beta,i} \cap \gamma = C_{\alpha,i} \cap \gamma$. Finally, the above definitions ensure that

$$\text{acc}(D_\alpha) = \text{acc}(D_{\alpha_0}) \cup \{\alpha_0\} \subseteq C_{\alpha_0,i(\alpha_0)} \cup \{\alpha_0\} \subseteq C_{\alpha,i(\alpha)}.$$

Case 3a: α is a limit of limit ordinals and $\text{acc}(D_\alpha) = \emptyset$. Pick a strictly increasing sequence $\langle \alpha_n \mid n < \omega \rangle$ of limit ordinals cofinal in α . Given $n < \omega$, let $i(\alpha) \leq j_n < \lambda$ be minimal with $i(\alpha_m) \leq j_n$ and $\alpha_m \in \text{acc}(C_{\alpha_{n+1},j_n})$ for all $m \leq n$. Then our induction hypothesis implies that $j_n \leq j_{n+1}$ for all $n < \omega$. Set $j_\omega = \sup_{n < \omega} j_n \leq \lambda$. If $i(\alpha) \leq i < j_0$, then we define $C_{\alpha,i} = \{\alpha_n \mid n < \omega\}$. Next, if $j_n \leq i < j_{n+1}$ for some $n < \omega$, then $i \geq i(\alpha_n)$, and we define $C_{\alpha,i} = C_{\alpha_n,i} \cup \{\alpha_m \mid n \leq m < \omega\}$. Finally, if $j_\omega \leq i < \lambda$, then $i \geq i(\alpha_n)$ for all $n < \omega$, and we define $C_{\alpha,i} = \bigcup \{C_{\alpha_n,i} \mid n < \omega\}$.

The above definitions and our induction hypothesis directly ensure that clauses (1) and (5) from Definition 3.2 hold. If $m < n < \omega$ and $i \geq j_{n-1}$, then the above definitions ensure that $\alpha_m \in \text{acc}(C_{\alpha_n,i}) \subseteq C_{\alpha_n,i}$ holds. In particular, we have $\{\alpha_n \mid n < \omega\} \subseteq C_{\alpha,i}$ for all $i(\alpha) \leq i < \lambda$. Moreover, we have $C_{\alpha_m,i} = C_{\alpha_n,i} \cap \alpha_m$ for all $j_\omega \leq i < \lambda$ and $m < n < \omega$. This shows that clause (2) from Definition 3.2 holds. Now, fix $i(\alpha) \leq i < j < \lambda$. If either i or j is contained in $[i(\alpha), j_0) \cup [j_\omega, \lambda)$, then the above remark and our induction hypothesis imply that $C_{\alpha,i} \subseteq C_{\alpha,j}$ holds. Next, if there is an $n < \omega$ with $j_n \leq i < j < j_{n+1}$, then our induction hypothesis implies that $C_{\alpha_n,i} \subseteq C_{\alpha_n,j}$ and therefore $C_{\alpha,i} \subseteq C_{\alpha,j}$ also holds in this case. Finally, if there are $m < n < \omega$ with $j_m \leq i < j_{m+1} \leq j_n \leq j < j_{n+1}$, then $\alpha_m \in \text{acc}(C_{\alpha_n,j})$, $C_{\alpha_m,i} \subseteq C_{\alpha_m,j} = C_{\alpha_n,j} \cap \alpha_m$ and, in combination with the above remarks, this implies that $C_{\alpha,i} \subseteq C_{\alpha,j}$ holds. These computations show that clause (3) of Definition 3.2 holds in this case. We finally verify clause (4) of Definition 3.2. To this end, fix $i(\alpha) \leq i < \lambda$ and $\gamma \in \text{acc}(C_{\alpha,i})$. Then $i \geq j_0$. If there is an $n < \omega$ with $j_n \leq i < j_{n+1}$, then it follows that $\gamma \in \text{acc}(C_{\alpha_n,i}) \cup \{\alpha_n\}$, in which case the induction hypothesis implies that $i \geq i(\gamma)$ and $C_{\gamma,i} = C_{\alpha_n,i} \cap \gamma = C_{\alpha,i} \cap \gamma$. In the other case, assume that $j_\omega \leq i < \lambda$ and let $n < \omega$ be least such that $\gamma < \alpha_n$. By the above remarks and our induction hypothesis, we then have $C_{\alpha,i} \cap \gamma = C_{\alpha_n,i} \cap \gamma$, $\gamma \in \text{acc}(C_{\alpha_n,i})$, $i \geq i(\gamma)$ and $C_{\gamma,i} = C_{\alpha_n,i} \cap \gamma = C_{\alpha,i} \cap \gamma$.

Case 3b: α is a limit of limit ordinals and $\text{acc}(D_\alpha) \neq \emptyset$ is bounded below α . Pick a strictly increasing sequence $\langle \alpha_n \mid n < \omega \rangle$ of limit ordinals cofinal in α with $\alpha_0 = \max(\text{acc}(D_\alpha))$. Then $\alpha_0 \in \text{acc}(D_\alpha)$ implies that $i(\alpha_0) = i(\alpha)$. Define a sequence $\langle j_n \mid n \leq \omega \rangle$ as in Case 3a. If $i(\alpha) \leq i < j_0$, then $i \geq i(\alpha_0)$, and we define $C_{\alpha,i} = C_{\alpha_0,i} \cup \{\alpha_n \mid n < \omega\}$. Next, if $j_n \leq i < j_{n+1}$ for some $n < \omega$, then $i \geq i(\alpha_n)$, and we define $C_{\alpha,i} = C_{\alpha_n,i} \cup \{\alpha_m \mid n \leq m < \omega\}$. Finally, if $j_\omega \leq i < \lambda$, then $i \geq i(\alpha_n)$ for all $n < \omega$, and we define $C_{\alpha,i} = \bigcup \{C_{\alpha_n,i} \mid n < \omega\}$.

As above, it is easy to see that clauses (1) and (5) from Definition 3.2 hold. Moreover, we again have $\alpha_m \in \text{acc}(C_{\alpha_n,i}) \subseteq C_{\alpha_n,i}$ for all $m < n < \omega$ and $i \geq j_{n-1}$ and, together with our induction hypothesis, this implies that

$$C_{\alpha_0,i} \cup \{\alpha_n \mid n < \omega\} \subseteq C_{\alpha,i}$$

for all $i(\alpha) \leq i < \lambda$ and $C_{\alpha_m,i} = C_{\alpha_n,i} \cap \alpha_m$ for all $j_\omega \leq i < \lambda$ and $m < n < \omega$. Hence clause (2) from Definition 3.2 holds. As in Case 3a, clauses (3) and (4) from Definition 3.2 are a direct consequence of our induction hypothesis and the above observations. Finally, our construction and the induction hypothesis ensure that

$$\text{acc}(D_\alpha) = \text{acc}(D_{\alpha_0}) \cup \{\alpha_0\} \subseteq \text{acc}(C_{\alpha_0,i(\alpha_0)}) \cup \{\alpha_0\} \subseteq \text{acc}(C_{\alpha,i(\alpha)}).$$

Case 4: $\text{acc}(D_\alpha)$ is unbounded in α . Note that, for all $\beta, \gamma \in \text{acc}(D_\alpha)$ with $\beta > \gamma$, our construction and the induction hypothesis imply that $i(\alpha) = i(\beta) = i(\gamma)$, $\gamma \in \text{acc}(C_{\beta, i(\alpha)})$, and therefore $C_{\gamma, i} = C_{\beta, i} \cap \gamma$ for all $i(\alpha) \leq i < \lambda$. If we now define $C_{\alpha, i} = \bigcup \{C_{\beta, i} \mid \beta \in \text{acc}(D_\alpha)\}$ for all $i(\alpha) \leq i < \lambda$, then it is easy to see that clauses (1)–(3) and (5) of Definition 3.2 hold. To verify clause (4), fix $i(\alpha) \leq i < \lambda$ and $\gamma \in \text{acc}(C_{\alpha, i})$. Let $\beta = \min(\text{acc}(D_\alpha) \setminus (\gamma + 1))$. It follows that $\gamma \in \text{acc}(C_{\beta, i})$, so, by the induction hypothesis, we have $i \geq i(\gamma)$ and $C_{\gamma, i} = C_{\beta, i} \cap \gamma = C_{\alpha, i} \cap \gamma$. Finally, our induction hypothesis implies that

$$\begin{aligned} \text{acc}(D_\alpha) &= \bigcup \{\text{acc}(D_\beta) \mid \beta \in \text{acc}(D_\alpha)\} \\ &\subseteq \bigcup \{\text{acc}(C_{\beta, i(\beta)}) \mid \beta \in \text{acc}(D_\alpha)\} \subseteq \text{acc}(C_{\alpha, i(\alpha)}). \end{aligned}$$

We have thus constructed a matrix \vec{C} satisfying clauses (1)–(5) of Definition 3.2. We finish the proof by verifying condition (6') from Proposition 3.3. To this end, suppose for the sake of a contradiction that there is $i < \lambda$ and a club E in κ such that $i \geq i(\alpha)$ holds for all $\alpha \in \text{acc}(E)$. Let $T = \{\alpha \in S \mid i(\alpha) = i + 1\}$. Since T is stationary in κ , there is $\alpha \in \text{acc}(E) \cap T$. But then $i \geq i(\alpha) = i + 1$, a contradiction. \square

In the proof of the following result, we use Todorćević's method of *walks on ordinals* to construct trees with ascent paths from suitable square principles.

Theorem 3.5. *If $\lambda < \kappa$ are infinite regular cardinals and $\square^{\text{ind}}(\kappa, \lambda)$ holds, then there is a κ -Aronszajn tree with a λ -ascent path.*

Proof. Fix a $\square^{\text{ind}}(\kappa, \lambda)$ -sequence $\vec{C} = \langle C_{\alpha, i} \mid \alpha < \kappa, i(\alpha) \leq i < \lambda \rangle$. Given $i < \lambda$, let $\vec{C}_i = \langle C_\alpha^i \mid \alpha < \kappa \rangle$ denote the unique C -sequence⁵ with $C_\alpha^i = C_{\alpha, i(\alpha)}$ for all $\alpha \in \text{acc}(\kappa)$ with $i < i(\alpha)$ and $C_\alpha^i = C_{\alpha, i}$ for all $\alpha \in \text{acc}(\kappa)$ with $i(\alpha) \leq i$. For each $i < \lambda$, recursively define $\rho_0^{\vec{C}_i} : [\kappa]^2 \rightarrow {}^{<\omega}\kappa$ as in [25] by letting, for all $\alpha < \beta < \kappa$,

$$\rho_0^{\vec{C}_i}(\alpha, \beta) = \langle \text{otp}(C_\beta^i \cap \alpha) \rangle \frown \rho_0^{\vec{C}_i}(\alpha, \min(C_\beta^i \setminus \alpha)),$$

subject to the boundary condition $\rho_0^{\vec{C}_i}(\alpha, \alpha) = \emptyset$ for all $\alpha < \kappa$. Given $i < \lambda$, we set

$$\mathbb{T}_i = \mathbb{T}(\rho_0^{\vec{C}_i}) = (\{\rho_0^{\vec{C}_i}(\cdot, \beta) \upharpoonright \alpha \mid \alpha \leq \beta < \kappa\}, \subset).$$

Claim 1. *If $i < \lambda$, then the tree \mathbb{T}_i has no cofinal branches.*

Proof of the Claim. Suppose \mathbb{T}_i has a cofinal branch. By [25, (1.7)], there is a club D in κ and a $\xi < \kappa$ with the property that, for all $\alpha < \kappa$, there is $\beta(\alpha) \geq \alpha$ with

$$D \cap \alpha = C_{\beta(\alpha)}^i \cap [\xi, \alpha].$$

Given $\alpha \in \text{acc}(D)$, we have $\beta(\alpha) \in \text{acc}(\kappa)$ and there is a $j(\alpha) < \lambda$ with the property that $C_{\beta(\alpha)}^i = C_{\beta(\alpha), j(\alpha)}$. Since \vec{C} is a $\square^{\text{ind}}(\kappa, \lambda)$ -sequence and $\alpha \in \text{acc}(C_{\beta(\alpha)}^i) = \text{acc}(C_{\beta(\alpha), j(\alpha)})$ for all $\alpha \in \text{acc}(D)$, we can conclude that

$$D \cap \alpha = C_{\beta(\alpha)}^i \cap [\xi, \alpha] = C_{\beta(\alpha), j(\alpha)} \cap [\xi, \alpha] = C_{\alpha, j(\alpha)} \cap [\xi, \alpha].$$

Fix an unbounded subset E of $\text{acc}(D)$ and $j < \lambda$ with $j(\alpha) = j$ for all $\alpha \in E$. Given $\alpha, \beta \in E$ with $\xi < \alpha < \beta$, we have $\alpha \in \text{acc}(C_{\beta, j})$ and therefore $C_{\alpha, j} = C_{\beta, j} \cap \alpha$. If

⁵Following [25, Section 1], given an uncountable regular cardinal κ , we say that a sequence $\langle C_\alpha \mid \alpha < \kappa \rangle$ is a C -sequence if for all $\alpha < \kappa$, the set C_α is a closed unbounded subset of α and $C_{\alpha+1} = \{\alpha\}$ holds.

we define $C^* = \bigcup\{C_{\alpha,j} \mid \xi < \alpha \in E\}$, then C^* is a club in κ and, for all $\alpha \in \text{acc}(C^*)$, we have $C_{\alpha,j} = C^* \cap \alpha$, contradicting the fact that \vec{C} is a $\square^{\text{ind}}(\kappa, \lambda)$ -sequence. \square

Claim 2. *If $i < \lambda$ and $\alpha < \kappa$, then the α -th level of \mathbb{T}_i has cardinality less than κ .*

Proof of the Claim. Let $\mathcal{D}_\alpha^i = \{C_\beta^i \cap \alpha \mid \alpha \leq \beta < \kappa\}$. By [25, (1.3)], the α -th level of \mathbb{T}_i has cardinality at most $|\mathcal{D}_\alpha^i| + \aleph_0$. If $D \in \mathcal{D}_\alpha^i$, then D is the union of a finite subset of α and a set of the form $C_{\gamma,j}$, where $\gamma \leq \alpha$ and $j < \lambda$. There are only $|\alpha|$ -many finite subsets of α and only $\max\{\lambda, |\alpha|\}$ -many sets of the form $C_{\gamma,j}$, where $\gamma \leq \alpha$ and $j < \lambda$. In combination, this shows that $|\mathcal{D}_\alpha^i| \leq \max\{\lambda, |\alpha|\} < \kappa$. \square

Now, define \mathbb{T} to be the unique tree with the following properties:

- (1) The underlying set of \mathbb{T} is the collection of all pairs $\langle i, t \rangle$ such that $i < \lambda$, $t \in \mathbb{T}_i$ and $\text{ht}_{\mathbb{T}_i}(t)$ is a limit ordinal.
- (2) Given nodes $\langle i, t \rangle$ and $\langle j, u \rangle$ in \mathbb{T} , we have $\langle i, t \rangle \leq_{\mathbb{T}} \langle j, u \rangle$ if and only if $i = j$ and $t \leq_{\mathbb{T}_i} u$.

Then Claims 1 and 2 directly imply that \mathbb{T} is a κ -Aronszajn tree. Given $\alpha < \kappa$, we define

$$c_\alpha : \lambda \longrightarrow \mathbb{T}; \quad i \longmapsto \langle i, \rho_0^{\vec{C}_i}(\cdot, \omega \cdot \alpha) \rangle.$$

Claim 3. *The sequence $\langle c_\alpha \mid \alpha < \kappa \rangle$ is a λ -ascent path in \mathbb{T} .*

Proof of the Claim. Note that, for all $i < \lambda$ and all $\alpha < \beta < \kappa$, if $C_{\omega \cdot \alpha}^i = C_{\omega \cdot \beta}^i \cap (\omega \cdot \alpha)$, then

$$\rho_0^{\vec{C}_i}(\cdot, \omega \cdot \alpha) = \rho_0^{\vec{C}_i}(\cdot, \omega \cdot \beta) \upharpoonright \omega \cdot \alpha.$$

Let $j_{\alpha,\beta} < \lambda$ be least such that $\omega \cdot \alpha \in \text{acc}(C_{\omega \cdot \beta, j_{\alpha,\beta}}^i)$. Then, for all $j_{\alpha,\beta} \leq i < \lambda$, we have $C_{\omega \cdot \alpha}^i = C_{\omega \cdot \alpha, i}$, $C_{\omega \cdot \beta}^i = C_{\omega \cdot \beta, i}$, and $C_{\omega \cdot \alpha, i} = C_{\omega \cdot \beta, i} \cap \omega \cdot \alpha$. It follows that, for all $\alpha < \beta < \kappa$ and all $j_{\alpha,\beta} \leq i < \lambda$, we have $c_\alpha(i) <_{\mathbb{T}} c_\beta(i)$. \square

This completes the proof of the theorem. \square

The statement of Theorem 1.1 now follows directly from Theorems 3.4 and 3.5.

4. FORCING PRELIMINARIES

In this section, we review some forcing posets designed to add and thread⁶ square sequences. We also recall constructions to make weak compactness or the tree property indestructible under mild forcing.

4.1. Forcing square sequences. We first recall the notion of strategic closure.

Definition 4.1. Let \mathbb{P} be a partial order (with maximal element $\mathbb{1}_{\mathbb{P}}$) and let β be an ordinal.

- (1) $\mathcal{D}_\beta(\mathbb{P})$ is the two-player game of perfect information in which Players I and II alternate playing conditions from \mathbb{P} to attempt to construct a $\leq_{\mathbb{P}}$ -decreasing sequence $\langle p_\alpha \mid \alpha < \beta \rangle$. Player I plays at all odd stages, and Player II plays at all even (including limit) stages. Player II is required to play $p_0 = \mathbb{1}_{\mathbb{P}}$. If, during the course of play, a limit ordinal $\alpha < \beta$ is

⁶Given an uncountable regular cardinal κ , a cardinal $1 < \lambda \leq \kappa$ and a sequence $\vec{C} = \langle C_\alpha \mid \alpha < \kappa \rangle$ that satisfies the first two properties listed in Definition 3.1, a club D in κ is a *thread through \vec{C}* if D witnesses a failure of the third property listed in Definition 3.1, i.e. if $D \cap \alpha \in C_\alpha$ holds for all $\alpha \in \text{acc}(D)$.

reached such that $\langle p_\xi \mid \xi < \alpha \rangle$ has no lower bound in \mathbb{P} , then Player I wins. Otherwise, Player II wins.

(2) \mathbb{P} is β -strategically closed if Player II has a winning strategy in $\mathfrak{D}_\beta(\mathbb{P})$.

Definition 4.2. Given cardinals $1 < \lambda \leq \kappa$ with κ regular and uncountable, we let $\mathbb{S}(\kappa, < \lambda)$ denote the partial order defined by the following clauses:

- (1) A condition in $\mathbb{S}(\kappa, < \lambda)$ is a sequence $p = \langle \mathcal{C}_\alpha^p \mid \alpha \leq \gamma^p \rangle$ with $\gamma^p \in \text{acc}(\kappa)$ such that the following statements hold for all $\alpha, \beta \in \text{acc}(\gamma^p + 1)$:
 - (a) \mathcal{C}_α^p is a collection of club subsets of α with $0 < |\mathcal{C}_\alpha^p| < \lambda$.
 - (b) If $C \in \mathcal{C}_\beta^p$ with $\alpha \in \text{acc}(C)$, then $C \cap \alpha \in \mathcal{C}_\alpha^p$.
- (2) The ordering of $\mathbb{S}(\kappa, < \lambda)$ is given by end-extension, i.e., $q \leq_{\mathbb{S}(\kappa, < \lambda)} p$ holds if and only if $\gamma^q \geq \gamma^p$ and, for all $\alpha \leq \gamma^p$, $\mathcal{C}_\alpha^q = \mathcal{C}_\alpha^p$.

In the following, we will usually write $\mathbb{S}(\kappa, \lambda)$ instead of $\mathbb{S}(\kappa, < \lambda^+)$. The proof of the following lemma is standard and follows, for example, from the proofs of [14, Proposition 33 and Lemma 35].

Lemma 4.3. *Let $1 < \lambda \leq \kappa$ be cardinals with κ regular and uncountable.*

- (1) $\mathbb{S}(\kappa, < \lambda)$ is ω_1 -closed.⁷
- (2) $\mathbb{S}(\kappa, < \lambda)$ is κ -strategically closed.
- (3) If G is $\mathbb{S}(\kappa, < \lambda)$ -generic over V , then $\bigcup G$ is a $\square(\kappa, < \lambda)$ -sequence in $V[G]$.

We next consider a forcing poset meant to add a thread to a $\square(\kappa, < \lambda)$ -sequence.

Definition 4.4. Let $1 < \lambda \leq \kappa$ be cardinals with κ regular and uncountable, and let $\vec{C} = \langle \mathcal{C}_\alpha \mid \alpha < \kappa \rangle$ be a $\square(\kappa, < \lambda)$ -sequence. We let $\mathbb{T}(\vec{C})$ denote the partial order whose underlying set is $\bigcup \{ \mathcal{C}_\alpha \mid \alpha \in \text{acc}(\kappa) \}$ and whose ordering is given by end-extension, i.e. $D \leq_{\mathbb{T}(\vec{C})} C$ holds if and only if $C = D \cap \text{sup}(C)$.

In what follows, if \mathbb{P} is a partial order and θ is a cardinal, then we let \mathbb{P}^θ denote the full-support product of θ copies of \mathbb{P} .

Lemma 4.5. *Let $1 < \lambda \leq \kappa$ be cardinals with κ regular and uncountable, let $\mathbb{S} = \mathbb{S}(\kappa, < \lambda)$, let \dot{C} be the canonical \mathbb{S} -name for the $\square(\kappa, < \lambda)$ -sequence added by \mathbb{S} and let $\dot{\mathbb{T}}$ be the canonical \mathbb{S} -name for $\mathbb{T}(\dot{C})$. For all $0 < \theta < \lambda$, the partial order $\mathbb{S} * \dot{\mathbb{T}}^\theta$ has a dense κ -directed closed subset.*

Proof. Fix $\theta < \lambda$, and let \mathbb{U} denote the set of all $\langle p, \dot{f} \rangle \in \mathbb{S} * \dot{\mathbb{T}}^\theta$ such that, for all $\xi < \theta$, there is a $D_\xi \in \mathcal{C}_{\gamma^p}^p$ with $p \Vdash_{\mathbb{S}} \dot{f}(\xi) = \check{D}_\xi$. We claim that \mathbb{U} is as required.

Let us first show that \mathbb{U} is dense in $\mathbb{S} * \dot{\mathbb{T}}^\theta$. To this end, fix $(p_0, \dot{f}_0) \in \mathbb{S} * \dot{\mathbb{T}}^\theta$. Since \mathbb{S} is κ -strategically closed and hence does not add any new sequences from V of length less than κ , we can assume, extending p_0 if necessary, that, for all $\xi < \theta$, there is a set $D_{\xi,0} \in V$ such that $p_0 \Vdash_{\mathbb{S}} \dot{f}_0(\xi) = \check{D}_{\xi,0}$. We may also assume that, for all $\xi < \theta$, we have $\text{sup}(D_{\xi,0}) < \gamma^{p_0}$. We now define a condition $p \leq_{\mathbb{S}} p_0$ such that $\gamma^p = \gamma^{p_0} + \omega$. It suffices to define $\mathcal{C}_{\gamma^p}^p$. For each $\xi < \theta$, let $D_\xi = D_{\xi,0} \cup \{ \text{sup}(D_{\xi,0}) \} \cup \{ \gamma^{p_0} + n \mid n < \omega \}$, and let $\mathcal{C}_{\gamma^p}^p = \{ D_\xi \mid \xi < \theta \}$. Finally, let \dot{f} be an \mathbb{S} -name for an element of $\dot{\mathbb{T}}^\theta$ such that, for all $\xi < \theta$, $p \Vdash_{\mathbb{S}} \dot{f}(\xi) = \check{D}_\xi$. It is easily verified that (p, \dot{f}) is an element of \mathbb{U} extending (p_0, \dot{f}_0) .

Let us now show that \mathbb{U} is κ -directed closed. Since \mathbb{U} is *tree-like*, i.e., if f_0 extends both f_1 and f_2 in \mathbb{U} , then f_1 and f_2 are themselves comparable by $\leq_{\mathbb{U}}$, it suffices

⁷As defined in [13, Chapter VIII, Definition 6.12].

to show that \mathbb{U} is κ -closed. Fix a limit ordinal $\zeta < \kappa$, and let $\langle (p_\eta, \dot{f}_\eta) \mid \eta < \zeta \rangle$ be a strictly decreasing sequence of conditions in \mathbb{U} . Let $\gamma = \sup\{\gamma^{p_\eta} \mid \eta < \zeta\}$. For each $\xi < \theta$ and $\eta < \zeta$, let $D_{\xi, \eta}$ be the element of $\mathcal{C}_{\gamma^{p_\eta}}^{p_\eta}$ such that $p_\eta \Vdash_{\mathbb{S}} \text{“}\dot{f}_\eta(\xi) = \check{D}_{\xi, \eta}\text{”}$. For each $\xi < \theta$, let $D_\xi = \bigcup_{\eta < \zeta} D_{\xi, \eta}$, and note that D_ξ is a club in γ . Define a condition $p \in \mathbb{S}$ with $\gamma^p = \gamma$ as follows. If $\alpha \in \text{acc}(\gamma)$, then find $\eta < \zeta$ such that $\alpha \leq \gamma^{p_\eta}$, and let $C_\alpha^p = C_\alpha^{p_\eta}$. Let $C_\gamma^p = \{D_\xi \mid \xi < \theta\}$. Finally, let \dot{f} be an \mathbb{S} -name for an element of $\dot{\mathbb{T}}^\theta$ such that, for all $\xi < \theta$, $p \Vdash_{\mathbb{S}} \text{“}\dot{f}(\xi) = \check{D}_\xi\text{”}$. It is again easily verified that (p, \dot{f}) is an element of \mathbb{U} and extends (p_η, \dot{f}_η) for all $\eta < \zeta$. \square

Remark 4.6. The proof of the above Lemma also makes it clear that, if G is \mathbb{S} -generic over V and $\alpha < \kappa$, then the set of all $t \in \dot{\mathbb{T}}^G$ such that $\alpha < \text{sup}(t)$ is a dense subset of $\dot{\mathbb{T}}^G$ in $V[G]$. From this observation and the Lemma, we can conclude that, in $V[G]$, forcing with $\dot{\mathbb{T}}^G$ preserves all cofinalities and adds a thread through the generically-added $\square(\kappa, < \lambda)$ -sequence.

There is also a natural forcing notion to add a $\square^{\text{ind}}(\kappa, \lambda)$ -sequence by initial segments.

Definition 4.7. Given infinite, regular cardinals $\lambda < \kappa$, we let $\mathbb{P}(\kappa, \lambda)$ denote the partial order defined by the following clauses:

- (1) A condition in $\mathbb{P}(\kappa, \lambda)$ is a matrix

$$p = \langle C_{\alpha, i}^p \mid \alpha \leq \gamma^p, i(\alpha)^p \leq i < \lambda \rangle$$

with $\gamma^p \in \text{acc}(\kappa)$ and $i(\alpha)^p < \lambda$ for all $\alpha \leq \gamma^p$ such that the following statements hold for all $\alpha, \beta \in \text{acc}(\gamma^p + 1)$:

- (a) If $i(\alpha)^p \leq i < \lambda$, then $C_{\alpha, i}^p$ is a club subset of α .
 - (b) If $i(\alpha)^p \leq i < j < \lambda$, then $C_{\alpha, i}^p \subseteq C_{\alpha, j}^p$.
 - (c) If $\alpha \in \text{acc}(C_{\beta, i}^p)$ for some $i(\beta)^p \leq i < \lambda$, then we have $i \geq i(\alpha)^p$ and $C_{\alpha, i}^p = C_{\beta, i}^p \cap \alpha$.
 - (d) If $\alpha < \beta$, then there is $i < \lambda$ with $\alpha \in \text{acc}(C_{\beta, i}^p)$.
- (2) The ordering of $\mathbb{P}(\kappa, \lambda)$ is given by end-extension, i.e., $q \leq_{\mathbb{P}(\kappa, \lambda)} p$ holds if and only if $\gamma^q \geq \gamma^p$ and the following statements hold for all $\alpha \in \text{acc}(\gamma^p + 1)$:
- (a) $i(\alpha)^q = i(\alpha)^p$.
 - (b) If $i(\alpha)^p \leq i < \lambda$, then $C_{\alpha, i}^q = C_{\alpha, i}^p$.

Lemma 4.8. *Let $\lambda < \kappa$ be infinite, regular cardinals.*

- (1) $\mathbb{P}(\kappa, \lambda)$ is λ -directed closed.
- (2) $\mathbb{P}(\kappa, \lambda)$ is κ -strategically closed.
- (3) *If G is $\mathbb{P}(\kappa, \lambda)$ -generic over V , then $\bigcup G$ is a $\square^{\text{ind}}(\kappa, \lambda)$ -sequence in $V[G]$ and for every regular cardinal $\mu < \kappa$ and every $i^* < \lambda$, the set*

$$\{\alpha \in E_\mu^\kappa \mid \exists p \in G [\alpha \leq \gamma^p \wedge i(\alpha)^p = i^*]\}$$

is stationary in κ in $V[G]$.

Proof. Clauses (1) and (2) are exactly [16, Propositions 7.2 and 7.3], respectively, so we omit their proofs. Let us sketch a proof of Clause (3). Working in V , let $\mathbb{P} = \mathbb{P}(\kappa, \lambda)$ and fix a regular cardinal $\mu < \kappa$, an ordinal $i^* < \lambda$, a condition $p^* \in \mathbb{P}$, and a \mathbb{P} -name \dot{C} forced by p^* to be a club in κ . We will find a condition $p \leq_{\mathbb{P}} p^*$ such that $\text{cof}(\gamma^p) = \mu$, $p \Vdash_{\mathbb{P}} \text{“}\dot{\gamma}^p \in \dot{C}\text{”}$, and $i(\gamma^p)^p = i^*$. Since p^* was arbitrary, this will suffice to prove the Lemma.

Let $\langle p_\alpha \mid \alpha < \mu + 1 \rangle$ be a run of the game $\mathcal{D}_{\mu+1}(\mathbb{P})$ satisfying the following conditions.

- $p_1 \leq_{\mathbb{P}} p^*$.
- Player II plays according to the winning strategy isolated in the proof of [16, Proposition 7.2]. In particular, the set $\{\gamma^{p_\alpha} \mid \alpha < \mu, \alpha \text{ is even}\}$ is a club in γ^{p_μ} and, for every non-zero, even ordinal $\alpha \leq \mu$, we have $i(\gamma^{p_\alpha})^{p_\alpha} = 0$.
- For every even ordinal $\alpha < \mu$, Player I chooses $p_{\alpha+1}$ so that there is β_α for which $\gamma^{p_\alpha} \leq \beta_\alpha < \gamma^{p_{\alpha+1}}$ and $p_{\alpha+1} \Vdash_{\mathbb{P}} \check{\beta}_\alpha \in \check{C}$.

Let $\gamma = \gamma^{p_\mu}$, and note that $\text{cof}(\gamma) = \mu$. We will modify p_μ to obtain p as desired, with $\gamma^p = \gamma$. For all $\alpha < \gamma$, let $i(\alpha)^p = i(\alpha)^{p_\mu}$, and, for all $i(\alpha)^p \leq i < \lambda$, let $C_{\alpha,i}^p = C_{\alpha,i}^{p_\mu}$. Then, let $i(\gamma)^p = i^*$ and, for all $i^* \leq i < \lambda$, let $C_{\gamma,i}^p = C_{\gamma,i}^{p_\mu}$. It is easily seen that p is still a condition in \mathbb{P} and extends p_α for all $\alpha < \mu$. Since $\gamma = \sup\{\beta_\alpha \mid \alpha < \mu, \alpha \text{ is even}\}$ and \check{C} is forced by p_1 to be a club in κ , we have $p \Vdash_{\mathbb{P}} \check{\gamma} \in \check{C}$. But we also clearly have $i(\gamma)^p = i^*$, as desired. This finishes the proof. \square

We finally introduce a forcing notion to add a thread through a $\square^{\text{ind}}(\kappa, \lambda)$ -sequence.

Definition 4.9. Let $\vec{C} = \langle C_{\alpha,i} \mid \alpha < \kappa, i(\alpha) \leq i < \lambda \rangle$ be a $\square^{\text{ind}}(\kappa, \lambda)$ -sequence. Given $i < \lambda$, let $\mathbb{T}_i(\vec{C})$ denote the partial order whose underlying set is

$$\{C_{\alpha,i} \mid \alpha \in \text{acc}(\kappa), i(\alpha) \leq i\}$$

and whose ordering is given by end-extension.

The following result is proven in [10, Section 3].

Lemma 4.10. *Let $\lambda < \kappa$ be infinite, regular cardinals, let \check{C} be the canonical $\mathbb{P}(\kappa, \lambda)$ -name for the generic $\square^{\text{ind}}(\lambda, \kappa)$ -sequence and, for $i < \kappa$, let $\dot{\mathbb{T}}_i$ be a $\mathbb{P}(\kappa, \lambda)$ -name for $\mathbb{T}_i(\check{C})$.*

- (1) *For all $i < \lambda$, $\mathbb{P}(\kappa, \lambda) * \dot{\mathbb{T}}_i$ has a dense κ -directed closed subset.*
- (2) *Let G be $\mathbb{P}(\kappa, \lambda)$ -generic over V and let $\bigcup G = \langle C_{\alpha,i} \mid \alpha < \kappa, i(\alpha) \leq i < \lambda \rangle$ be the generic $\square^{\text{ind}}(\lambda, \kappa)$ -sequence. Given $i < j < \lambda$, the map*

$$\pi_{i,j} : \dot{\mathbb{T}}_i^G \longrightarrow \dot{\mathbb{T}}_j^G; C_{\alpha,i} \longmapsto C_{\alpha,j}$$

is a forcing projection in $V[G]$.

Lemma 4.11. *Let $\lambda < \kappa$ be infinite, regular cardinals, let G be $\mathbb{P}(\kappa, \lambda)$ -generic over V , and let \vec{C} be the generic $\square^{\text{ind}}(\lambda, \kappa)$ -sequence in $V[G]$. Assume that, in $V[G]$, for each $i < \lambda$, \dot{x}_i is a $\mathbb{T}_i(\vec{C})$ -name for an element of $V[G]$. Given $t \in \mathbb{T}_0(\vec{C})$, there is a condition $s \leq_{\mathbb{T}_0(\vec{C})} t$ such that $\pi_{0,i}(s)$ decides the value of \dot{x}_i for all $i < \lambda$.*

Proof. Work in $V[G]$. For all $i < \lambda$, let \mathbb{D}_i be the set of $s \in \mathbb{T}_0(\vec{C})$ such that $\pi_{0,i}(s)$ decides the value of \dot{x}_i . Since $\pi_{0,i}$ is a projection, the set \mathbb{D}_i is dense and open in $\mathbb{T}_0(\vec{C})$. Since $\mathbb{P}(\kappa, \lambda) * \dot{\mathbb{T}}_0$ has a dense κ -directed closed subset in V , it follows that the partial order $\mathbb{T}_0(\vec{C})$ is κ -distributive in $V[G]$. We can thus find $s \in \bigcap \{\mathbb{D}_i \mid i < \lambda\}$ with $s \leq_{\mathbb{T}_0(\vec{C})} t$. Then the condition s is as desired. \square

4.2. Indestructibility. In this subsection, we outline techniques for arranging so that weak compactness and the tree property necessarily hold at a cardinal κ after forcing with the partial order $\text{Add}(\kappa, 1)$ that adds a Cohen subset to κ . These methods are well-known, so we will just sketch the constructions or refer the reader elsewhere for details.

A classical result of Silver shows that weak compactness of a weakly compact cardinal can be made indestructible under forcing with $\text{Add}(\kappa, 1)$. The proof of this result is described in [12, Section 3]. A modern presentation of these arguments can be found in [6, Example 16.2].

Lemma 4.12 (Silver). *Let κ be a weakly compact cardinal. Then there is a forcing extension $V[G]$ such that κ is weakly compact in $V[G, H]$ whenever H is $\text{Add}(\kappa, 1)$ -generic over $V[G]$.*

Note that the conclusion of the above result implies that κ is also weakly compact in $V[G]$, because forcing with κ -closed partial orders does not add branches to κ -Aronszajn trees.

Definition 4.13. Given an uncountable, regular cardinal κ , we let $\text{TP}^+(\kappa)$ denote the conjunction of $\text{TP}(\kappa)$ and the statement that for every regular cardinal λ with $\lambda^+ < \kappa$, if \vec{b} is a λ -ascent path through a tree \mathbb{T} of height κ , then there is a true cofinal branch through \vec{b} .

It is easy to see that, if κ is a weakly compact cardinal, then $\text{TP}^+(\kappa)$ holds. The proof of the following result is a slight modification of the presentation of Mitchell's consistency proof of the tree property at \aleph_2 in [6, Section 23].

Lemma 4.14. *Let κ be a weakly compact cardinal and let $\mu < \kappa$ be an infinite, regular cardinal with $\mu = \mu^{<\mu}$. Then there is a forcing extension $V[G]$ of the ground model V such that the following statements hold:*

- (1) V and $V[G]$ have the same cofinalities below $(\mu^+)^V$.
- (2) $(\mu^+)^V = (\mu^+)^{V[G]}$, $\kappa = (2^\mu)^{V[G]} = (\mu^{++})^{V[G]}$ and $\kappa = (\kappa^{<\kappa})^{V[G]}$.
- (3) If H is $\text{Add}(\kappa, 1)$ -generic over $V[G]$, then $\text{TP}^+(\kappa)$ holds in $V[G, H]$.

Sketch of the proof. Given $\alpha \leq \kappa$, let $\dot{\mathbb{R}}_\alpha$ denote the canonical $\text{Add}(\mu, \alpha)$ -name with the property that, whenever G is $\text{Add}(\mu, \alpha)$ -generic over V , then $\dot{\mathbb{R}}_\alpha^G = \text{Add}(\mu^+, 1)^{V[G]}$. By induction on $\alpha \leq \kappa$, we define a sequence $\langle \mathbb{Q}(\alpha) \mid \alpha \leq \kappa \rangle$ of partial orders with the property that, if $\alpha \leq \kappa$ is inaccessible and G is $\mathbb{Q}(\alpha)$ -generic over V , then V and $V[G]$ have the same cofinalities below $(\mu^+)^V$, $(\mu^+)^V = (\mu^+)^{V[G]}$, $\alpha = (2^\mu)^{V[G]} = (\mu^{++})^{V[G]}$, and $\alpha = (\alpha^{<\alpha})^{V[G]}$.

Fix $\beta \leq \kappa$ and assume that the partial order $\mathbb{Q}(\alpha)$ with the above properties has been defined for all $\alpha < \beta$. Given $\alpha < \beta$ inaccessible, let $\dot{\mathbb{S}}_\alpha$ denote the canonical $\mathbb{Q}(\alpha)$ -name with the property that, whenever G is $\mathbb{Q}(\alpha)$ -generic over V , then $\dot{\mathbb{S}}_\alpha^G = \text{Add}(\alpha, 1)^{V[G]}$. We define the underlying set of the partial order $\mathbb{Q}(\beta)$ to consist of triples $\langle p, f, g \rangle$ that satisfy the following statements:

- (a) p is a condition in $\text{Add}(\mu, \beta)$.
- (b) f is a partial function on β of cardinality at most μ with the property that for all $\alpha \in \text{dom}(f)$, α is a successor ordinal and $f(\alpha)$ is an $\text{Add}(\mu, \alpha)$ -name for a condition in $\dot{\mathbb{R}}_\alpha$.

- (c) g is a partial function on β of cardinality at most μ with the property that for all $\alpha \in \text{dom}(g)$, α is an inaccessible cardinal and $g(\alpha)$ is a $\mathbb{Q}(\alpha)$ -name for a condition in \dot{S}_α .

Given triples $\langle p_0, f_0, g_0 \rangle$ and $\langle p_1, f_1, g_1 \rangle$ satisfying the above statements, we define $\langle p_1, f_1, g_1 \rangle \leq_{\mathbb{Q}(\beta)} \langle p_0, f_0, g_0 \rangle$ to hold if and only if the following statements hold:

- (i) $p_1 \leq_{\text{Add}(\mu, \beta)} p_0$.
- (ii) $\text{dom}(f_1) \supseteq \text{dom}(f_0)$ and $\text{dom}(g_1) \supseteq \text{dom}(g_0)$.
- (iii) If $\alpha \in \text{dom}(f_0)$, then

$$p_1 \upharpoonright \alpha \Vdash_{\text{Add}(\mu, \alpha)} "f_1(\alpha) \leq_{\dot{S}_\alpha} f_0(\alpha)".$$

- (iv) If $\alpha \in \text{dom}(g_0)$, then

$$\langle p_1 \upharpoonright \alpha, f_1 \upharpoonright \alpha, g_1 \upharpoonright \alpha \rangle \Vdash_{\mathbb{Q}(\alpha)} "g_1(\alpha) \leq_{\dot{S}_\alpha} g_0(\alpha)".$$

By representing $\mathbb{Q}(\beta)$ as a projection of a product $\text{Add}(\mu, \beta) \times \mathbb{P}$ for some μ^+ -closed partial order \mathbb{P} , it is possible to show that $\mathbb{Q}(\beta)$ satisfies the assumptions listed above.

Set $\mathbb{Q} = \mathbb{Q}(\kappa)$. Given $\alpha < \kappa$ inaccessible, the canonical map

$$\pi_\alpha : \mathbb{Q} \longrightarrow \mathbb{Q}(\alpha) * \dot{S}_\alpha; \langle p, f, g \rangle \longmapsto \langle \langle p \upharpoonright \alpha, f \upharpoonright \alpha, g \upharpoonright \alpha \rangle, g(\alpha) \rangle$$

is a projection. Let G be \mathbb{Q} -generic over V , let H be $\text{Add}(\kappa, 1)$ -generic over $V[G]$, and let $G_\alpha * H_\alpha$ be the filter on $\mathbb{P}(\alpha) * \dot{S}_\alpha$ induced by G via π_α . Suppose for sake of contradiction that \mathbb{T} is a κ -Aronszajn tree (with κ as an underlying set) in $V[G, H]$. A routine application of weak compactness yields an inaccessible cardinal $\alpha < \kappa$ such that $\mathbb{T} \upharpoonright \alpha \in V[G_\alpha, H_\alpha]$ and $\mathbb{T} \upharpoonright \alpha$ is an α -Aronszajn tree in $V[G_\alpha, H_\alpha]$. However, standard arguments show that μ^+ -approximation holds between $V[G_\alpha, H_\alpha]$ and $V[G]$ (see [6, Definition 21.2]). It follows that α -Aronszajn trees in $V[G_\alpha, H_\alpha]$ cannot gain cofinal branches in $V[G, H]$. However, $\mathbb{T} \upharpoonright \alpha$ does have a cofinal branch in $V[G, H]$, namely the set of predecessors of any element of $\mathbb{T}(\alpha)$, which is a contradiction.

Similarly, if $\lambda \leq \mu$ and \vec{b} is a λ -ascent path through a κ -tree \mathbb{T} in $V[G, H]$ with no true cofinal branch, then there is an inaccessible $\alpha < \kappa$ such that $\mathbb{T} \upharpoonright \alpha, \vec{b} \upharpoonright \alpha \in V[G_\alpha, H_\alpha]$ and there is no true cofinal branch through $\vec{b} \upharpoonright \alpha$ in $V[G_\alpha, H_\alpha]$. As above, an appeal to μ^+ -approximation yields a contradiction. \square

5. CONSISTENCY RESULTS FOR TREES

Building on the results of the previous section, we prove consistency results that will provide upper bounds for the consistency strength of two interactions between ascent paths and special trees listed in Table 1.

Theorem 5.1. *Let κ be an uncountable cardinal with the property that $\kappa = \kappa^{<\kappa}$ and $\mathbb{1}_{\text{Add}(\kappa, 1)} \Vdash \text{TP}^+(\check{\kappa})$. Then the following statements hold in a cofinality-preserving forcing extension of the ground model:*

- (1) *There are κ -Aronszajn trees.*
- (2) *There are no special κ -trees.*
- (3) *For all λ with $\lambda^+ < \kappa$, there are no κ -Aronszajn trees with λ -ascent paths.*

Proof. Let $\mathbb{P} = \mathbb{S}(\kappa, 2)$ be the forcing from Definition 4.2 that adds a $\square(\kappa, 2)$ -sequence. Let \dot{C} be the canonical \mathbb{P} -name for the generically-added $\square(\kappa, 2)$ -sequence, and let \dot{Q} be the canonical \mathbb{P} -name for the partial order $\mathbb{T}(\dot{C})$ defined in Definition 4.4. By Lemma 4.5, if $\theta \in \{1, 2\}$, then the partial order $\mathbb{P} * \dot{Q}^\theta$ has a dense κ -directed closed subset. Moreover, since $\kappa = \kappa^{<\kappa}$ holds, the dense subset constructed in the proof of Lemma 4.5 has size κ and is hence forcing equivalent to $\text{Add}(\kappa, 1)$.

Let G be \mathbb{P} -generic over V , and set $\mathbb{Q} = \dot{Q}^G$. We claim that the above statements hold in $V[G]$. We first note that, since $\square(\kappa, 2)$ holds in $V[G]$, there are κ -Aronszajn trees in this model. To verify clause (2), note that our assumptions and the above computations imply that, if H is \mathbb{Q} -generic over $V[G]$, then κ is a regular cardinal in $V[G, H]$ and every κ -Aronszajn tree in $V[G]$ has a cofinal branch in $V[G, H]$. This observation directly implies that there are no special κ -trees in $V[G]$.

To verify clause (3), suppose for sake of a contradiction that, in $V[G]$, there is a κ -Aronszajn tree \mathbb{T} , a cardinal λ with $\lambda^+ < \kappa$, and a λ -ascent path $\vec{b} = \langle b_\alpha \mid \alpha < \kappa \rangle$ through \mathbb{T} . Given $\alpha < \beta < \kappa$, let $i_{\alpha, \beta} < \lambda$ be least such that, for all $i_{\alpha, \beta} \leq j < \lambda$, we have $b_\alpha(j) <_{\mathbb{T}} b_\beta(j)$. Since \mathbb{Q} forces $\text{TP}^+(\kappa)$ to hold, there are \mathbb{Q} -names \dot{I} and \dot{B} such that

$$\mathbb{1}_{\mathbb{Q}} \Vdash \text{“The pair } \langle \dot{I}, \dot{B} \rangle \text{ is a true cofinal branch through } \vec{b}\text{”}$$

holds in $V[G]$. Let $H_0 \times H_1$ be \mathbb{Q}^2 -generic over $V[G]$, and, given $\varepsilon < 2$, set $I_\varepsilon = \dot{I}^{H_\varepsilon}$ and $B_\varepsilon = \dot{B}^{H_\varepsilon}$. Note that, since $\mathbb{P} * \dot{Q}^2$ has a dense κ -directed closed subset in V , κ remains a regular cardinal in $V[G, H_0, H_1]$.

Work now in $V[G, H_0, H_1]$. For each $\alpha < \kappa$ and $\varepsilon < 2$, set $\alpha_\varepsilon = \min(B_\varepsilon \setminus (\alpha + 1))$. Using the regularity of κ and the fact that $\lambda < \kappa$, we find $\max(I_0, I_1) \leq I_* < \lambda$ and unbounded subsets $B_0^* \subseteq B_0$ and $B_1^* \subseteq B_1$ with $i_{\alpha, \alpha_{1-\varepsilon}} \leq I_*$ for all $\varepsilon < 2$ and all $\alpha \in B_\varepsilon^*$. Given $\varepsilon < 2$, set $A_\varepsilon = \{\alpha < \kappa \mid i_{\alpha, \alpha_\varepsilon} \leq I_*\}$.

Claim 1. *For $\varepsilon < 2$, the pair $\langle I_*, A_\varepsilon \rangle$ is a true cofinal branch through \vec{b} .*

Proof of the Claim. Fix $\alpha, \beta \in A_\varepsilon$ with $\alpha < \beta$. By definition of A_ε and the fact that $I_* \geq I_\varepsilon$, it follows that $b_\alpha(i) <_{\mathbb{T}} b_{\alpha_\varepsilon}(i) \leq_{\mathbb{T}} b_{\beta_\varepsilon}(i)$ and $b_\beta(i) <_{\mathbb{T}} b_{\beta_\varepsilon}(i)$ for all $I_* \leq i < \lambda$. Since $\leq_{\mathbb{T}}$ is a tree order, it follows that $b_\alpha(i) <_{\mathbb{T}} b_\beta(i)$ for all $I_* \leq i < \lambda$ and therefore $\langle I_*, A_\varepsilon \rangle$ satisfies Clause (3a) of Definition 2.5.

Next, fix $\beta \in A_\varepsilon$ and $\alpha < \beta$ such that $b_\alpha(i) <_{\mathbb{T}} b_\beta(i)$ for all $I_* \leq i < \lambda$. As above, it follows that $b_\alpha(i) <_{\mathbb{T}} b_\beta(i) <_{\mathbb{T}} b_{\beta_\varepsilon}(i)$ and $b_{\alpha_\varepsilon}(i) \leq_{\mathbb{T}} b_{\beta_\varepsilon}(i)$ for all $I_* \leq i < \lambda$. Thus, again by the fact that $\leq_{\mathbb{T}}$ is a tree order, this implies that $b_\alpha(i) <_{\mathbb{T}} b_{\alpha_\varepsilon}(i)$ for all $I_* \leq i < \lambda$ and hence $\alpha \in A_\varepsilon$. This allows us to conclude that the pair $\langle I_*, A_\varepsilon \rangle$ also satisfies Clause (3b) of Definition 2.5. \square

Claim 2. $A_0 = A_1$.

Proof of the Claim. We show $A_0 \subseteq A_1$. The proof of the reverse inclusion is symmetric. Thus, fix $\alpha \in A_0$ and set $\beta = \min(B_0^* \setminus \alpha_1)$. Given $I_* \leq i < \lambda$, we then have

$$b_\alpha(i) <_{\mathbb{T}} b_{\alpha_0}(i) \leq_{\mathbb{T}} b_\beta(i) <_{\mathbb{T}} b_{\beta_1}(i),$$

where the last relation holds because $\beta \in B_0^*$. But we also have $b_{\alpha_1}(i) <_{\mathbb{T}} b_{\beta_1}(i)$ for all $I_* \leq i < \lambda$, because $\alpha_1, \beta_1 \in B_1$ and $I_* \geq I_1$. But then, again using the fact that $\leq_{\mathbb{T}}$ is a tree order, we can conclude that $b_\alpha(i) <_{\mathbb{T}} b_{\alpha_1}(i)$ holds for all $I_* \leq i < \lambda$. This shows that α is an element of A_1 . \square

Given $\varepsilon < 2$, the set $A_0 = A_\varepsilon$ is definable from I_* and B_ε . Hence the pair $\langle I_*, A_0 \rangle$ is a member of $V[G][H_\varepsilon]$ for all $\varepsilon < 2$. By the Product Lemma, it follows that $\langle I_*, A_0 \rangle$ is contained in $V[G]$. But then $\{b_\alpha(I_*) \mid \alpha \in A_0\}$ is a cofinal branch through \mathbb{T} , contradicting the assumption that \mathbb{T} is a κ -Aronszajn tree in $V[G]$. \square

Next, we show that, consistently, there are κ -Aronszajn trees and all such trees contain ascent paths of small width.

Proof of Theorem 1.3. Let $\lambda < \kappa$ be infinite, regular cardinals such that $\kappa = \kappa^{<\kappa}$ and $\mathbb{1}_{\text{Add}(\kappa,1)} \Vdash \text{TP}(\check{\kappa})$. Define $\mathbb{P} = \mathbb{P}(\kappa, \lambda)$ to be the forcing notion from Definition 4.7 that adds a $\square^{\text{ind}}(\kappa, \lambda)$ -sequence. Let \dot{C} be the canonical \mathbb{P} -name for the generically-added $\square^{\text{ind}}(\kappa, \lambda)$ -sequence, and, for all $i < \lambda$, let \dot{Q}_i be a \mathbb{P} -name for the partial order $\mathbb{T}_i(\dot{C})$ defined in Definition 4.9. By Lemma 4.10, if $i < \lambda$, then the partial order $\mathbb{P} * \dot{Q}_i$ has a dense, κ -directed closed subset and the assumption that $\kappa = \kappa^{<\kappa}$ implies that this dense subset is forcing equivalent to $\text{Add}(\kappa, 1)$. Therefore, by our assumptions, we know that $\mathbb{1}_{\mathbb{P} * \dot{Q}_i} \Vdash \text{TP}(\check{\kappa})$ for all $i < \lambda$.

Let G be \mathbb{P} -generic over V and, for all $i < \lambda$, set $\mathbb{Q}_i = \dot{Q}_i^G$. Given $i < j < \lambda$, let $\pi_{i,j} : \mathbb{Q}_i \rightarrow \mathbb{Q}_j$ be the projection given by Lemma 4.10. Let

$$\dot{C}^G = \langle C_{\alpha,i} \mid \alpha < \kappa, i(\alpha) \leq i < \lambda \rangle$$

be the realization of \dot{C} . We claim that $V[G]$ is the desired forcing extension. Since $\square(\kappa, \lambda)$ holds in $V[G]$, there are κ -Aronszajn trees in this model. We thus verify requirement (2). To this end, work in $V[G]$ and fix a κ -Aronszajn tree \mathbb{T} . For all $i < \lambda$, we have $\mathbb{1}_{\mathbb{Q}_i} \Vdash \text{TP}(\check{\kappa})$ and hence we can fix a \mathbb{Q}_i -name \dot{B}_i for a cofinal branch in \mathbb{T} . We may assume that \dot{B}_i is forced to be $\leq_{\mathbb{T}}$ -downward closed, i.e., that \dot{B}_i is forced to meet every level of \mathbb{T} . For each $\alpha < \kappa$, use Lemma 4.11 to find $q_\alpha \in \mathbb{Q}_0$ such that, for all $i < \lambda$, there is a node $x_{\alpha,i} \in \mathbb{T}(\alpha)$ with the property that

$$\pi_{0,i}(q_\alpha) \Vdash_{\mathbb{Q}_i} \text{“} \check{x}_{\alpha,i} \in \dot{B}_i \cap \check{\mathbb{T}}(\check{\alpha}) \text{”}.$$

Define a sequence of functions $\vec{b} = \langle b_\alpha : \lambda \rightarrow \mathbb{T}(\alpha) \mid \alpha < \kappa \rangle$ by setting $b_\alpha(i) = x_{\alpha,i}$ for all $\alpha < \kappa$ and $i < \lambda$.

We claim that \vec{b} is a λ -ascent path through \mathbb{T} . To see this, fix $\alpha < \beta < \kappa$. Pick $\gamma_\alpha, \gamma_\beta \in \text{acc}(\kappa)$ such that $q_\alpha = C_{\gamma_\alpha,0}$ and $q_\beta = C_{\gamma_\beta,0}$. Without loss of generality, we may assume that $\gamma_\alpha < \gamma_\beta$; the other cases are treated similarly. Fix an $i < \lambda$ such that $\gamma_\alpha \in \text{acc}(C_{\gamma_\beta,i})$ and hence $\pi_{0,j}(q_\beta) \leq_{\mathbb{Q}_j} \pi_{0,j}(q_\alpha)$ for all $i \leq j < \lambda$. This shows that $\pi_{0,j}(q_\beta) \Vdash_{\mathbb{Q}_j} \text{“} x_{\alpha,j}, x_{\beta,j} \in \dot{B}_j \text{”}$ holds for all $i \leq j < \lambda$. Since \dot{B}_i is a name for a branch through \mathbb{T} , this implies that

$$b_\alpha(j) = x_{\alpha,j} <_{\mathbb{T}} x_{\beta,j} = b_\beta(j)$$

holds for all $i \leq j < \lambda$. Therefore, \vec{b} is a λ -ascent path through \mathbb{T} . \square

With the help of a result from [18], it is easy to see that the conclusion of Theorem 1.3 implies that κ is a weakly compact cardinal in L .

Lemma 5.2. *Let κ be a regular cardinal and let λ be an infinite cardinal with $\lambda^+ < \kappa$. If every κ -Aronszajn tree contains a λ -ascending path, then $\square(\kappa)$ fails.*

Proof. Assume, towards a contradiction, that there is a $\square(\kappa)$ -sequence \vec{C} and let $\mathbb{T} = \mathbb{T}(\rho_0^{\vec{C}})$ denote the tree of full codes of walks through \vec{C} defined in [25, Section

1] (as in the proof of Theorem 3.5). Then the results of [25] imply that \mathbb{T} is a κ -Aronszajn tree and [18, Corollary 4.7] shows that \mathbb{T} does not contain a λ -ascending path, contradicting our assumption. \square

It has long been known that, for regular cardinals $\kappa > \aleph_1$, the principle $\square(\kappa)$ does not imply the existence of a special κ -Aronszajn tree. For example, this is the case in L if κ is a Mahlo cardinal that is not weakly compact, and it will remain true in the forcing extension of L by $\text{Col}(\aleph_1, < \kappa)$, in which $\kappa = \aleph_2$. We now show that $\square(\kappa)$ does not even imply the existence of a κ -Aronszajn tree \mathbb{T} such that there is a stationary subset of κ that is non-stationary with respect to \mathbb{T} . In particular, this shows that the various trees that will be constructed from the principle $\square(\kappa)$ in the proofs of Theorem 1.4 and 1.6 in Section 7 (see Definition 7.1 and Lemma 7.2) cannot be assumed to be κ -Aronszajn trees.

Theorem 5.3. *Let $\kappa > \aleph_1$ be a regular cardinal with the property that $\kappa = \kappa^{< \kappa}$ and $\mathbb{1}_{\text{Add}(\kappa, 1)} \Vdash \text{TP}(\check{\kappa})$. Then the following statements hold in a cofinality-preserving forcing extension of the ground model:*

- (1) $\square(\kappa)$ holds.
- (2) If \mathbb{T} is a κ -Aronszajn tree and S is a stationary subset of κ , then S is stationary with respect to \mathbb{T} .

Proof. Let $\mathbb{S} = \mathbb{S}(\kappa, 1)$ be the forcing from Definition 4.2 that adds a $\square(\kappa)$ -sequence. Let \dot{C} be the canonical \mathbb{S} -name for the generically-added square sequence, and let $\dot{\mathbb{R}}$ be an \mathbb{S} -name for the threading forcing $\mathbb{T}(\dot{C})$ defined in Definition 4.4. By Lemma 4.5 and our assumptions, the partial order $\mathbb{S} * \dot{\mathbb{R}}$ has a dense κ -directed closed subset.

Let G be \mathbb{S} -generic over V and set $\mathbb{R} = \dot{\mathbb{R}}^G$. By [10, Lemma 3.1 and Corollary 3.1], there is a forcing iteration $\langle \mathbb{P}_\eta, \dot{Q}_\xi \mid \eta \leq \kappa^+, \xi < \kappa^+ \rangle$ in $V[G]$ with supports of size less than κ , such that, letting $\mathbb{P} = \mathbb{P}_{\kappa^+}$, the following statements hold:

- (a) If $\eta \leq \kappa^+$ and $\dot{\mathbb{P}}_\eta$ is the canonical \mathbb{S} -name for \mathbb{P}_η in V , then $\mathbb{S} * (\dot{\mathbb{P}}_\eta \times \dot{\mathbb{R}})$ has a κ -directed closed dense subset in V . Moreover, if $\eta < \kappa^+$, then this subset can be assumed to have size κ .
- (b) \mathbb{P} satisfies the κ^+ -chain condition in $V[G]$.
- (c) If H is \mathbb{P} -generic over $V[G]$, then $\square(\kappa)$ holds in $V[G, H]$ and, for every stationary subset E of κ in $V[G, H]$, there is a condition r in \mathbb{R} such that $r \Vdash_{\mathbb{R}} \check{E} \text{ is stationary in } \check{\kappa}$ holds in $V[G, H]$.

Let H be \mathbb{P} -generic over $V[G]$. For $\eta < \kappa^+$, let H_η be the \mathbb{P}_η -generic filter induced by H . We claim that $V[G, H]$ is the desired model. Thus, work in $V[G, H]$ and suppose, for the sake of contradiction, that \mathbb{T} is a κ -Aronszajn and E is a stationary subset of κ that is non-stationary with respect to \mathbb{T} . By the properties of \mathbb{P} , we can find $r \in \mathbb{R}$ with $r \Vdash_{\mathbb{R}} \check{E} \text{ is a stationary subset of } \check{\kappa}$.

Since the tree \mathbb{T} , the subset E , and the maps witnessing that E is non-stationary with respect to \mathbb{T} can all be coded by subsets of V of cardinality κ in $V[G, H]$, the fact that \mathbb{P} satisfies the κ^+ -chain condition in $V[G]$ implies that there is an $\eta < \kappa^+$ such that $E, \mathbb{T} \in V[G, H_\eta]$ and E is non-stationary with respect to \mathbb{T} in $V[G, H_\eta]$.

Let K be \mathbb{R} -generic over $V[G, H]$ with $r \in K$. Since the partial order $\mathbb{S} * (\dot{\mathbb{P}}_\eta \times \dot{\mathbb{R}})$ has a dense κ -directed closed subset of size κ in V , our assumptions imply that the tree property holds at κ in $V[G, H_\eta, K]$. However, E remains stationary in $V[G, H, K]$ and thus, *a fortiori*, in $V[G, H_\eta, K]$. Moreover, the maps witnessing that E is non-stationary with respect to \mathbb{T} obviously persist in $V[G, H_\eta, K]$, so E

remains non-stationary with respect to \mathbb{T} in $V[G, H_\eta, K]$, and so, by Fact 2.4, \mathbb{T} is a κ -Aronszajn tree in $V[G, H_\eta, K]$, contradicting the fact that the tree property holds at κ . \square

6. PROVABLE IMPLICATIONS

In this section, we piece things together to provide a complete explanation of Table 1 from the end of the Introduction, thus completing the picture of the interaction between special trees and trees with ascent paths at successors of regular cardinals. Throughout this section, we will work under the assumption that there are \aleph_2 -Aronszajn trees.

6.1. Inconsistencies. We first note that, by Lemma 2.6, an \aleph_2 -Aronszajn tree with an \aleph_0 -ascent path cannot be special (see also [18, Corollary 1.7] and [27, Proposition 2.3]). This immediately implies that $\forall\mathbb{T}.S(\mathbb{T})$ is incompatible with $\exists\mathbb{T}.A(\mathbb{T})$ and that $\forall\mathbb{T}.A(\mathbb{T})$ is incompatible with $\exists\mathbb{T}.S(\mathbb{T})$, so the three boxes in the upper left of the table are inconsistent.

6.2. Lower bounds. We now deal with lower bounds for the boxes in the bottom row and right column of the table. First, by Theorem 1.1, $\square(\aleph_2)$ implies the existence of an \aleph_2 -Aronszajn tree with an \aleph_0 -ascent path. By results of Jensen and Todorćević (see [11, Section 6] and [25, (1.10)]), a failure of $\square(\aleph_2)$ implies that \aleph_2 is weakly compact in L . Therefore, the consistency of $\forall\mathbb{T}.\neg A(\mathbb{T})$ implies the consistency of a weakly compact cardinal, which takes care of all three boxes in the right column of the table.

By Lemma 5.2, the assumption that $\forall\mathbb{T}.A(\mathbb{T})$ holds implies a failure of $\square(\kappa)$ and, as above, we can conclude that the consistency of $\forall\mathbb{T}.A(\mathbb{T})$ implies the consistency of a weakly compact cardinal.

Finally, Jensen's principle \square_{\aleph_1} implies the existence of a special \aleph_2 -Aronszajn tree and, by a result of Jensen in [11], the failure of \square_{\aleph_1} implies that \aleph_2 is Mahlo in L . In particular, the consistency of $\forall\mathbb{T}.\neg S(\mathbb{T})$ implies the consistency of a Mahlo cardinal, thus finishing our derivation of lower bounds.

6.3. Upper bounds. We finally deal with upper bounds. First, Theorem 1.3 shows that the consistency of $\forall\mathbb{T}.A(\mathbb{T})$ follows from the consistency of a weakly compact cardinal. Since $\forall\mathbb{T}.A(\mathbb{T})$ implies $\forall\mathbb{T}.\neg S(\mathbb{T})$, it follows that the consistency of the conjunction of these two statements follows from the consistency of a weakly compact cardinal.

Next, in [17], Laver and Shelah prove that the consistency of a weakly compact cardinal implies the consistency of Souslin's Hypothesis at \aleph_2 . A straightforward and well-known strengthening of their argument yields that the consistency of a weakly compact cardinal in fact implies the consistency of $\forall\mathbb{T}.S(\mathbb{T})$. Since $\forall\mathbb{T}.S(\mathbb{T})$ implies $\forall\mathbb{T}.\neg A(\mathbb{T})$, it follows that the consistency of the conjunction of these two statements follows from the consistency of a weakly compact cardinal. Moreover, it immediately follows that the conjunction of $\exists\mathbb{T}.S(\mathbb{T})$ and $\forall\mathbb{T}.\neg A(\mathbb{T})$ follows from the consistency of a weakly compact cardinal.

Since \square_{\aleph_1} implies the existence both of a special \aleph_2 -Aronszajn tree and an \aleph_2 -Aronszajn tree with an \aleph_0 -ascent path, it follows that the consistency of the conjunction of $\exists\mathbb{T}.S(\mathbb{T})$ and $\exists\mathbb{T}.A(\mathbb{T})$ does not require large cardinals.

Suppose κ is the least Mahlo cardinal in L , and force over L with the partial order constructed by Mitchell in [19] for κ . Then there are no special \aleph_2 -Aronszajn

trees in the extension. Since κ is not weakly compact in L , $\square(\aleph_2)$ holds in the extension, so, by Corollary 1.2, there is an \aleph_2 -Aronszajn tree with an \aleph_0 -ascent path. It follows that the consistency of the conjunction of $\forall\mathbb{T}.\neg S(\mathbb{T})$ and $\exists\mathbb{T}.A(\mathbb{T})$ follows from the consistency of a Mahlo cardinal.

Finally, Theorem 5.1 shows that the consistency of the conjunction of $\forall\mathbb{T}.\neg S(\mathbb{T})$ and $\forall\mathbb{T}.\neg A(\mathbb{T})$ follows from the consistency of a weakly compact cardinal, thus completing the table.

7. CHAIN CONDITIONS

The first part of this section is devoted to the proofs of Theorem 1.4 and 1.6. In the second part, we will use ideas from Section 5 to construct a model of set theory in which the class of all partial orders satisfying the κ -chain condition exhibits an interesting product behavior.

Definition 7.1. Suppose that $\vec{C} = \langle C_{\alpha,i} \mid \alpha < \kappa, i(\alpha) \leq i < \lambda \rangle$ is a $\square^{\text{ind}}(\kappa, \lambda)$ -sequence, and $i < \lambda$.

- (1) We define $S_i^{\vec{C}} = \{\alpha \in \text{acc}(\kappa) \mid i(\alpha) = i\}$ and $S_{\leq i}^{\vec{C}} = \{\alpha \in \text{acc}(\kappa) \mid i(\alpha) \leq i\}$.
The sets $S_{< i}^{\vec{C}}$, $S_{> i}^{\vec{C}}$, etc. are defined analogously.
- (2) We let $\mathbb{T}_i^{\vec{C}}$ denote the tree with underlying set $S_{\leq i}^{\vec{C}}$ and

$$\alpha <_{\mathbb{T}_i^{\vec{C}}} \beta \iff \alpha \in \text{acc}(C_{\beta,i})$$

for all $\alpha, \beta \in S_{\leq i}^{\vec{C}}$.

Lemma 7.2. Let $\vec{C} = \langle C_{\alpha,i} \mid \alpha < \kappa, i(\alpha) \leq i < \lambda \rangle$ be a $\square^{\text{ind}}(\kappa, \lambda)$ -sequence and let $i < \lambda$. If the tree $\mathbb{T}_i^{\vec{C}}$ has height κ , then the set $S_{> i}^{\vec{C}}$ is non-stationary with respect to $\mathbb{T}_i^{\vec{C}}$.

Proof. Set $\mathbb{T} = \mathbb{T}_i^{\vec{C}}$ and $S = S_{> i}^{\vec{C}}$. Let A denote the set of all $\alpha \in \mathbb{T} \upharpoonright S$ with $\text{acc}(C_{\alpha,i}) \cap \text{ht}_{\mathbb{T}}(\alpha) \neq \emptyset$. Given $\alpha \in A$, define $s(\alpha) = \sup(\text{acc}(C_{\alpha,i}) \cap \text{ht}_{\mathbb{T}}(\alpha))$. Since $\alpha \geq \text{ht}_{\mathbb{T}}(\alpha) \in S$, we have $\alpha \notin S$ and therefore $\alpha > \text{ht}_{\mathbb{T}}(\alpha)$ holds for all $\alpha \in A$. This shows that $s(\alpha) \leq \text{ht}_{\mathbb{T}}(\alpha) < \alpha$, $s(\alpha) \in \text{acc}(C_{\alpha,i})$, $i(s(\alpha)) \leq i$ and $s(\alpha) \in \mathbb{T}$. Moreover, we have $\text{acc}(C_{\alpha,i}) \cap S = \emptyset$ and hence $s(\alpha) < \text{ht}_{\mathbb{T}}(\alpha)$ for all $\alpha \in A$.

Given $\beta \in \mathbb{T}$, define A_β to be the set of all $<_{\mathbb{T}}$ -minimal elements γ in $s^{-1}\{\beta\}$. Let B denote the set of all $\alpha \in A$ with $\alpha \notin A_{s(\alpha)}$, and let $r : \mathbb{T} \upharpoonright S \rightarrow \mathbb{T}$ denote the unique function with the following properties:

- (1) If $\alpha \in B$, then $r(\alpha)$ is the unique element γ of $A_{s(\alpha)}$ with $\gamma <_{\mathbb{T}} \alpha$.
- (2) If $\alpha \in A \setminus B$, then $r(\alpha) = s(\alpha)$.
- (3) If $\alpha \in (\mathbb{T} \upharpoonright S) \setminus A$ is not minimal in \mathbb{T} , then we define $r(\alpha) = \min(\text{acc}(C_{\alpha,i}))$.
- (4) If $\alpha \in \mathbb{T} \upharpoonright S$ is minimal in \mathbb{T} , then we define $r(\alpha) = \alpha$.

By the above remarks, the function r is regressive on $\mathbb{T} \upharpoonright S$. Fix $\gamma \in \mathbb{T}$ and let $c_\gamma : r^{-1}\{\gamma\} \rightarrow \omega \times \kappa$ denote the unique function with the following properties:

- (a) If $\alpha \in B$, then $c_\gamma(\alpha) = \langle 0, \text{ht}_{\mathbb{T}}(\alpha) \rangle$.
- (b) If $\alpha \in A \setminus B$, then $c_\gamma(\alpha) = \langle 1, 0 \rangle$.
- (c) If $\alpha \in (\mathbb{T} \upharpoonright S) \setminus A$ is not minimal in \mathbb{T} , then $c_\gamma(\alpha) = \langle 2, \text{ht}_{\mathbb{T}}(\alpha) \rangle$.
- (d) If $\alpha \in \mathbb{T} \upharpoonright S$ is minimal in \mathbb{T} , then $c_\gamma(\alpha) = \langle 3, 0 \rangle$.

Then c_γ is injective on $<_{\mathbb{T}}$ -chains in $r^{-1}\{\gamma\}$. If $\alpha \in \text{dom}(c_\gamma) \setminus A$ is not minimal in \mathbb{T} , then we have $\text{acc}(C_{\alpha,i}) \neq \emptyset$, $\text{acc}(C_{\alpha,i}) \cap \text{ht}_{\mathbb{T}}(\alpha) = \emptyset$ and therefore

$$\text{ht}_{\mathbb{T}}(\alpha) \leq \min(\text{acc}(C_{\alpha,i})) = r(\alpha) = \gamma.$$

Next, pick $\alpha \in B \cap \text{dom}(c_\gamma)$. Then we have $\text{ht}_{\mathbb{T}}(\alpha) < \alpha$, $\gamma \in A_{s(\alpha)} \subseteq A \subseteq \mathbb{T} \upharpoonright S$ and $\gamma <_{\mathbb{T}} \alpha$. This implies that $\text{ht}_{\mathbb{T}}(\gamma) < \min\{\text{ht}_{\mathbb{T}}(\alpha), \gamma\}$, $C_{\gamma,i} = C_{\alpha,i} \cap \gamma$ and

$$\max(\text{acc}(C_{\alpha,i}) \cap \text{ht}_{\mathbb{T}}(\alpha)) = \max(\text{acc}(C_{\gamma,i}) \cap \text{ht}_{\mathbb{T}}(\gamma)) = \max(\text{acc}(C_{\alpha,i}) \cap \text{ht}_{\mathbb{T}}(\gamma)).$$

In particular, we have $\text{acc}(C_{\alpha,i}) \cap [\text{ht}_{\mathbb{T}}(\gamma), \text{ht}_{\mathbb{T}}(\alpha)] = \emptyset$ and therefore $\text{ht}_{\mathbb{T}}(\alpha) \leq \gamma$, because otherwise we would have $\gamma \in \text{acc}(C_{\alpha,i}) \cap [\text{ht}_{\mathbb{T}}(\gamma), \text{ht}_{\mathbb{T}}(\alpha)]$. These computations show that the range of c_γ has cardinality strictly less than κ . \square

The following type of partial order will be crucial in our construction of Knaster partial orders with interesting product behavior.

Definition 7.3. Given a tree \mathbb{T} , we let $\mathbb{P}(\mathbb{T})$ denote the partial order consisting of finite partial functions $f : \mathbb{T} \xrightarrow{\text{part}} \omega$ that are injective on $<_{\mathbb{T}}$ -chains and that are ordered by reverse inclusion.

Remember that a tree \mathbb{T} is *extensional at limit levels* if $\text{pred}_{\mathbb{T}}(s) \neq \text{pred}_{\mathbb{T}}(t)$ holds for every limit ordinal α and all $s, t \in \mathbb{T}(\alpha)$ with $s \neq t$.

Lemma 7.4. *Let κ be an uncountable regular cardinal, let $\mu < \kappa$ be a (possibly finite) cardinal with $\nu^\mu < \kappa$ for all $\nu < \kappa$, let S be a subset of $E_{>\mu}^\kappa$ that is stationary in κ , and let $\langle \mathbb{T}^\gamma \mid \gamma < \mu \rangle$ be a sequence of trees of height at most κ that are extensional at limit levels. Assume that S is non-stationary with respect to \mathbb{T}^γ for every $\gamma < \lambda$ with $\text{ht}(\mathbb{T}^\gamma) = \kappa$. Then the full support product $\prod_{\gamma < \mu} \mathbb{P}(\mathbb{T}^\gamma)$ is κ -Knaster.*

Proof. Since two conditions in a partial order of the form $\mathbb{P}(\mathbb{T})$ are compatible if and only if their union is a condition, it suffices to prove the statement for trees of cardinality κ , because we can always consider trees of this form that are given by the downward closures of the unions of the domains of κ -sequences of conditions. Note that every such tree is isomorphic to a tree \mathbb{T} with the property that the underlying set of \mathbb{T} is a subset of $\kappa \times \kappa$, we have $\mathbb{T}(\alpha) \subseteq \{\alpha\} \times \kappa$ for all $\alpha < \kappa$ and $\langle \alpha_0, \beta_0 \rangle <_{\mathbb{T}} \langle \alpha_1, \beta_1 \rangle$ implies $\beta_0 < \beta_1$ for all nodes $\langle \alpha_0, \beta_0 \rangle$ and $\langle \alpha_1, \beta_1 \rangle$ in \mathbb{T} .

Assume that all trees in the above sequence are of this form and fix a sequence $\langle p_\alpha \mid \alpha < \kappa \rangle$ of conditions in $\prod_{\gamma < \mu} \mathbb{P}(\mathbb{T}^\gamma)$. If $\gamma < \mu$ and $\text{ht}(\mathbb{T}^\gamma) = \kappa$, then we also fix functions $r_\gamma : \mathbb{T}^\gamma \upharpoonright S \rightarrow \mathbb{T}^\gamma$ and $\langle c_t^\gamma : r_\gamma^{-1}\{t\} \rightarrow \kappa_t^\gamma \mid t \in \mathbb{T}^\gamma \rangle$ witnessing the non-stationarity of S with respect to \mathbb{T}^γ . In the other case, if $\gamma < \mu$ and $\text{ht}(\mathbb{T}^\gamma) < \kappa$, then we let $r_\gamma : \mathbb{T}^\gamma \upharpoonright S \rightarrow \mathbb{T}^\gamma$ denote the unique regressive function with $\text{ran}(r_\gamma) \subseteq \mathbb{T}^\gamma(0)$ and we set $c_{r_\gamma(t)}^\gamma(t) = \text{ht}_{\mathbb{T}^\gamma}(t)$ for all $t \in \mathbb{T}^\gamma \upharpoonright S$. Now, define \mathbb{D}_γ to be the set of all conditions p in $\mathbb{P}(\mathbb{T}^\gamma)$ with the property that for all $t, u \in \text{dom}(p)$ with $\text{ht}_{\mathbb{T}^\gamma}(t) < \text{ht}_{\mathbb{T}^\gamma}(u)$, there is $s \in \text{dom}(p)$ with $\text{ht}_{\mathbb{T}^\gamma}(s) = \text{ht}_{\mathbb{T}^\gamma}(t)$ and $s <_{\mathbb{T}^\gamma} u$. Then it is easy to see that \mathbb{D}_γ is a dense subset of $\mathbb{P}(\mathbb{T}^\gamma)$ and we can pick a sequence $\langle q_\alpha \mid \alpha < \kappa \rangle$ of conditions in $\prod_{\gamma < \mu} \mathbb{P}(\mathbb{T}^\gamma)$ with $q_\alpha(\gamma) \leq_{\mathbb{P}(\mathbb{T}^\gamma)} p_\alpha(\gamma)$ and $q_\alpha(\gamma) \in \mathbb{D}_\gamma$ for all $\alpha < \kappa$ and $\gamma < \mu$. By our assumptions on the trees \mathbb{T}^γ , there is a club C of limit ordinals in κ with the property that $\text{dom}(q_\alpha(\gamma)) \subseteq \beta \times \beta$ holds for all $\alpha, \beta \in C$ with $\alpha < \beta$ and all $\gamma < \mu$. For all $\alpha \in C$ and $\gamma < \mu$, fix an injective enumeration $\langle t_k^{\alpha,\gamma} \mid k < n_{\alpha,\gamma} \rangle$ of the finite set $\{t \in \mathbb{T}^\gamma(\alpha) \mid \exists u \in \text{dom}(q_\alpha(\gamma)) t <_{\mathbb{T}^\gamma} u\}$. Since the

trees \mathbb{T}^γ are extensional at limit levels and S is a subset of $E_{>\mu}^\kappa$, there is a regressive function $\rho : C \cap S \rightarrow \kappa$ and a matrix

$$\langle \iota_{\alpha,\gamma} : n_{\alpha,\gamma} \longrightarrow \mathbb{T}^\gamma(\rho(\alpha)) \mid \alpha \in C \cap S, \gamma < \mu \rangle$$

of injections with $\text{dom}(q_\alpha(\gamma)) \cap \mathbb{T}_{<\alpha}^\gamma \subseteq \mathbb{T}_{<\rho(\alpha)}^\gamma$, $\text{dom}(q_\alpha(\gamma)) \cap (\alpha \times \alpha) \subseteq \rho(\alpha) \times \rho(\alpha)$ and $r_\gamma(t_k^{\alpha,\gamma}) <_{\mathbb{T}^\gamma} \iota_{\alpha,\gamma}(k) <_{\mathbb{T}^\gamma} t_k^{\alpha,\gamma}$ for all $\alpha \in C \cap S$, $\gamma < \mu$ and $k < n_\alpha$.

In this situation, the assumption that $\nu^\mu < \kappa$ holds for all $\nu < \kappa$ yields a stationary subset E of $C \cap S$, an ordinal $\xi < \kappa$, a sequence $\langle n_\gamma \mid \gamma < \mu \rangle$ of natural numbers, a subset $K \subseteq \mu \times \omega$, a subset $H \subseteq \mu \times \xi$ and a sequence $\langle D_\gamma \mid \gamma < \mu \rangle$ of finite subsets of $\kappa \times \kappa$ such that the following statements hold for all $\alpha, \beta \in E$ and $\langle \gamma, k \rangle \in K$:

- (1) $n_\gamma = n_{\alpha,\gamma}$ and $\rho(\alpha) = \xi$.
- (2) $K = \{ \langle \gamma, k \rangle \mid \gamma < \mu, k < n_\gamma, \iota_{\alpha,\gamma}(k) \in \alpha \times \alpha \}$.
- (3) $H = \{ \langle \gamma, \text{ht}_{\mathbb{T}^\gamma}(t) \rangle \mid \gamma < \mu, t \in \text{dom}(q_\alpha(\gamma)) \cap \mathbb{T}_{<\alpha}^\gamma \}$.
- (4) $D_\gamma = \text{dom}(q_\alpha(\gamma)) \cap (\alpha \times \alpha)$ and $q_\alpha(\gamma) \upharpoonright D_\gamma = q_\beta(\gamma) \upharpoonright D_\gamma$.
- (5) $\iota_{\alpha,\gamma}(k) = \iota_{\beta,\gamma}(k)$, $r_\gamma(t_k^{\alpha,\gamma}) = r_\gamma(t_k^{\beta,\gamma})$ and $c_{r_\gamma(t_k^{\alpha,\gamma})}^\gamma(t_k^{\alpha,\gamma}) = c_{r_\gamma(t_k^{\beta,\gamma})}^\gamma(t_k^{\beta,\gamma})$.

Now, pick $\alpha, \beta \in E$ with $\alpha < \beta$ and assume for a contradiction that the conditions q_α and q_β are incompatible in $\prod_{\gamma < \mu} \mathbb{P}(\mathbb{T}^\gamma)$. Then there is a $\gamma < \mu$ such that $q_\alpha(\gamma) \cup q_\beta(\gamma)$ is not a condition in $\mathbb{P}(\mathbb{T}^\gamma)$ and hence there are $<_{\mathbb{T}^\gamma}$ -comparable nodes $t, u \in \mathbb{T}^\gamma$ such that $t \in \text{dom}(q_\alpha(\gamma)) \setminus \text{dom}(q_\beta(\gamma))$, $u \in \text{dom}(q_\beta(\gamma)) \setminus \text{dom}(q_\alpha(\gamma))$ and $q_\alpha(\gamma)(t) = q_\beta(\gamma)(u)$. But then $t <_{\mathbb{T}^\gamma} u$, because otherwise $u <_{\mathbb{T}^\gamma} t \in \beta \times \beta$ would imply that $u \in \text{dom}(q_\beta(\gamma)) \cap (\beta \times \beta) = D_\gamma \subseteq \text{dom}(q_\alpha(\gamma))$. Next, assume that $\text{ht}_{\mathbb{T}^\gamma}(t) < \alpha$. Then $\langle \gamma, \text{ht}_{\mathbb{T}^\gamma}(t) \rangle \in H$ and there is a $t_0 \in \text{dom}(q_\beta(\gamma))$ with $\text{ht}_{\mathbb{T}^\gamma}(t_0) = \text{ht}_{\mathbb{T}^\gamma}(t) < \text{ht}_{\mathbb{T}^\gamma}(u)$. Since $q_\beta(\gamma) \in \mathbb{D}_\gamma$, we can find $s \in \text{dom}(q_\beta(\gamma))$ with $\text{ht}_{\mathbb{T}^\gamma}(s) = \text{ht}_{\mathbb{T}^\gamma}(t_0)$ and $s <_{\mathbb{T}^\gamma} u$. But then $s = t \in \text{dom}(q_\beta(\gamma))$, a contradiction. This shows that $\text{ht}_{\mathbb{T}^\gamma}(t) \geq \alpha > \xi$ and hence there is a $k < n_\gamma$ with $t_k^{\alpha,\gamma} \leq_{\mathbb{T}^\gamma} t$. But then $\text{ht}_{\mathbb{T}^\gamma}(u) > \text{ht}_{\mathbb{T}^\gamma}(t) > \xi = \rho(\beta)$ implies that $\text{ht}_{\mathbb{T}^\gamma}(u) \geq \beta$ and hence there is an $l < n_\gamma$ with $t_l^{\beta,\gamma} \leq_{\mathbb{T}^\gamma} u$. Since $t <_{\mathbb{T}^\gamma} u$ and $\text{ht}_{\mathbb{T}^\gamma}(\iota_{\alpha,\gamma}(k)) = \xi = \text{ht}_{\mathbb{T}^\gamma}(\iota_{\beta,\gamma}(l))$, we know that $\iota_{\alpha,\gamma}(k) = \iota_{\beta,\gamma}(l)$ and therefore $\iota_{\beta,\gamma}(l) <_{\mathbb{T}^\gamma} t \in \beta \times \beta$ implies that $\langle \gamma, l \rangle \in K$. This shows that $\iota_{\alpha,\gamma}(k) = \iota_{\beta,\gamma}(l) = \iota_{\alpha,\gamma}(l)$ and the injectivity of $\iota_{\alpha,\gamma}$ implies that $k = l$ and $\langle \gamma, k \rangle \in K$. In this situation, the above choices ensure that $r_\gamma(t_k^{\alpha,\gamma}) = r_\gamma(t_k^{\beta,\gamma})$ and $c_{r_\gamma(t_k^{\alpha,\gamma})}^\gamma(t_k^{\alpha,\gamma}) = c_{r_\gamma(t_k^{\beta,\gamma})}^\gamma(t_k^{\beta,\gamma})$. Since $t_k^{\alpha,\gamma} \neq t_k^{\beta,\gamma}$, this implies that $\text{ht}(\mathbb{T}^\gamma) = \kappa$ and hence the nodes $t_k^{\alpha,\gamma}$ and $t_k^{\beta,\kappa}$ are incompatible in \mathbb{T}^γ . But this yields a contradiction, because we have $t_k^{\alpha,\gamma} \leq_{\mathbb{T}^\gamma} t <_{\mathbb{T}^\gamma} u$ and $t_k^{\beta,\gamma} \leq_{\mathbb{T}^\gamma} u$.

The above computations show that the sequence $\langle p_\alpha \mid \alpha \in E \rangle$ consists of pairwise compatible conditions in $\prod_{\gamma < \mu} \mathbb{P}(\mathbb{T}^\gamma)$. \square

We now introduce the κ -Knaster partial order that is used in the proofs of Theorem 1.4 and 1.6.

Definition 7.5. Suppose that \vec{C} is a $\square^{\text{ind}}(\kappa, \lambda)$ -sequence. We let $\mathbb{P}_{\vec{C}}$ denote the lottery sum of the sequence $\langle \mathbb{P}(\mathbb{T}_i^{\vec{C}}) \mid i < \lambda \rangle$ of partial orders, i.e., conditions in $\mathbb{P}_{\vec{C}}$ are pairs $\langle p, i \rangle$ with $i < \lambda$ and $p \in \mathbb{P}(\mathbb{T}_i^{\vec{C}})$ and, given $\langle p, i \rangle, \langle q, j \rangle \in \mathbb{P}_{\vec{C}}$, we have $\langle p, i \rangle \leq_{\mathbb{P}_{\vec{C}}} \langle q, j \rangle$ if either $i = j$ and $p \leq_{\mathbb{P}(\mathbb{T}_i^{\vec{C}})} q$ or $q = \mathbb{1}_{\mathbb{P}(\mathbb{T}_j^{\vec{C}})}$.

Lemma 7.6. Let $\vec{C} = \langle C_{\alpha,i} \mid \alpha < \kappa, i(\alpha) \leq i < \lambda \rangle$ be a $\square^{\text{ind}}(\kappa, \lambda)$ -sequence with the property that the set $E_{\geq \lambda}^\kappa \cap S_{\geq i}^{\vec{C}}$ is stationary in κ for all $i < \lambda$.

- (1) If $\mu < \lambda$ is a cardinal with $\nu^\mu < \kappa$ for all $\nu < \kappa$, then the full support product $\mathbb{P}_{\mathcal{C}}^\mu$ is κ -Knaster.
- (2) The full support product $\mathbb{P}_{\mathcal{C}}^\lambda$ does not satisfy the κ -chain condition.

Proof. (1) Let $\langle p_\alpha \mid \alpha < \kappa \rangle$ be a sequence of conditions in $\mathbb{P}_{\mathcal{C}}^\mu$. Since $\lambda^\mu < \kappa$ holds, we may assume that there is a function $f : \mu \rightarrow \lambda$ and a sequence $\langle q_\alpha \mid \alpha < \kappa \rangle$ of conditions in the full support product $\prod_{\gamma < \mu} \mathbb{P}(\mathbb{T}_{f(\gamma)}^{\vec{C}})$ with the property that $p_\alpha(\gamma) = \langle q_\alpha(\gamma), f(\gamma) \rangle$ holds for all $\alpha < \kappa$ and $\gamma < \mu$. Set $i_* = \text{lub}(\text{ran}(f)) < \lambda$. In this situation, Lemma 7.2 shows that for every $\gamma < \mu$ with the property that the tree $\mathbb{T}_{f(\gamma)}^{\vec{C}}$ has height κ , the set $S_{\geq i_*}^{\vec{C}} \subseteq S_{> f(\gamma)}^{\vec{C}}$ is non-stationary with respect to $\mathbb{T}_{f(\gamma)}^{\vec{C}}$. Since our assumptions imply that the set $E_{\geq \lambda}^\kappa \cap S_{\geq i_*}^{\vec{C}}$ is stationary in κ , we can apply Lemma 7.4 to conclude that the product $\prod_{\gamma < \mu} \mathbb{P}(\mathbb{T}_{f(\gamma)}^{\vec{C}})$ is κ -Knaster. Hence there is an unbounded subset U of κ such that the sequence $\langle q_\alpha \mid \alpha \in U \rangle$ consists of pairwise compatible conditions in $\prod_{\gamma < \mu} \mathbb{P}(\mathbb{T}_{f(\gamma)}^{\vec{C}})$ and this implies that the sequence $\langle p_\alpha \mid \alpha \in U \rangle$ consists of pairwise compatible conditions in $\mathbb{P}_{\mathcal{C}}^\mu$.

(2) Given $\alpha \in \text{acc}(\kappa)$ and $i < \lambda$, the function $\{\langle \alpha, 0 \rangle\}$ is a condition in the partial order $\mathbb{P}(\mathbb{T}_{\max\{i, i(\alpha)\}}^{\vec{C}})$. This shows that for every $\alpha \in \text{acc}(\kappa)$, there is a unique condition p_α in $\mathbb{P}_{\mathcal{C}}^\lambda$ with $p_\alpha(i) = \{\{\langle \alpha, 0 \rangle\}, \max\{i, i(\alpha)\}\}$ for all $i < \lambda$. Fix $\alpha, \beta \in \text{acc}(\kappa)$ with $\alpha < \beta$. Then there is $i(\beta) \leq i < \lambda$ such that $\alpha \in \text{acc}(C_{\beta, i})$. This implies that $i \geq i(\alpha)$, $C_{\alpha, i} = C_{\beta, i} \cap \alpha$ and $\alpha <_{\mathbb{T}_{\vec{C}}} \beta$. We can conclude that the conditions $p_\alpha(i) = \{\{\langle \alpha, 0 \rangle\}, i\}$ and $p_\beta(i) = \{\{\langle \beta, 0 \rangle\}, i\}$ are incompatible in $\mathbb{P}_{\mathcal{C}}$ and therefore the condition p_α and p_β are incompatible in $\mathbb{P}_{\mathcal{C}}^\lambda$. These computations show that the sequence $\langle p_\alpha \mid \alpha \in \text{acc}(\kappa) \rangle$ enumerates an antichain in $\mathbb{P}_{\mathcal{C}}^\lambda$. \square

The statement of Theorem 1.4 now follows directly from an application of Theorem 3.4 with $S = E_{\geq \lambda}^\kappa$ and Lemma 7.6. Moreover, by combining the above with the results of [4], we can show that $\square(\kappa)$ implies the existence of a κ -Knaster partial order that is not κ -stationarily layered.

Proof of Theorem 1.6. Assume that κ is an uncountable regular cardinal with the property that every κ -Knaster partial order is κ -stationarily layered. Then [4, Theorem 1.11] shows that κ is a Mahlo cardinal with the property that every stationary subset of κ reflects. Assume, towards a contradiction, that $\square(\kappa)$ holds. In this situation, we can apply Theorem 3.4 to obtain a $\square^{\text{ind}}(\kappa, \aleph_0)$ -sequence $\vec{C} = \langle C_{\alpha, i} \mid \alpha < \kappa, i(\alpha) \leq i < \omega \rangle$ with the property that there exists a $\square(\kappa)$ -sequence $\langle D_\alpha \mid \alpha < \kappa \rangle$ such that $\text{acc}(D_\alpha) \subseteq \text{acc}(C_{\alpha, i(\alpha)})$ holds for all $\alpha \in \text{acc}(\kappa)$.

Claim. *The set $\{\alpha \in \text{acc}(\kappa) \mid \text{otp}(D_\alpha) < \alpha\}$ is not stationary in κ .*

Proof of the Claim. Assume, for a contradiction, that the set is stationary in κ . Then a pressing down argument yields $\xi < \kappa$ such that the set

$$E = \{\alpha \in \text{acc}(\kappa) \mid \text{otp}(D_\alpha) = \xi\}$$

is stationary in κ . By the above remarks, there is an $\alpha < \kappa$ such that $\text{cof}(\alpha) > \omega$ and the set $E \cap \alpha$ is stationary in α . But then we can find $\beta, \gamma \in \text{acc}(D_\alpha) \cap E$ with $\gamma < \beta$. This implies that $D_\gamma = D_\beta \cap \gamma$ and hence $\xi = \text{otp}(D_\gamma) < \text{otp}(D_\beta) = \xi$, a contradiction. \square

By the above claim, we can find a club C in κ consisting of strong limit cardinals such that $\text{otp}(D_\alpha) = \alpha$ holds for all $\alpha \in C$. Let $\mathbb{P}_{\vec{C}}$ denote the partial order defined in Definition 7.5. Then a combination of Theorem 3.4 with Lemma 7.6 implies that $\mathbb{P}_{\vec{C}}$ is κ -Knaster and, by our assumption, this shows that $\mathbb{P}_{\vec{C}}$ is κ -stationarily layered. Pick a sufficiently large regular cardinal $\theta > \kappa$. Then [4, Lemma 2.3] shows that there is an elementary substructure M of $\mathbb{H}(\theta)$ of cardinality less than κ and $\alpha \in C$ such that $\alpha = \kappa \cap M$, $\vec{C} \in M$, and $\mathbb{P}_{\vec{C}} \cap M$ is a regular suborder of $\mathbb{P}_{\vec{C}}$. Set $p_0 = \{\langle \alpha, 0 \rangle\}$. Then $p = \langle p_0, i(\alpha) \rangle$ is a condition in $\mathbb{P}_{\vec{C}}$ and there is a *reduct* q of p in $\mathbb{P}_{\vec{C}} \cap M$, i.e., $q \in M$ is a condition in $\mathbb{P}_{\vec{C}}$ with the property that for every $r \in \mathbb{P}_{\vec{C}} \cap M$ with $r \leq_{\mathbb{P}_{\vec{C}}} q$, the conditions p and r are compatible in $\mathbb{P}_{\vec{C}}$ (see [4, Proposition 2.1]). Then there is a condition q_0 in $\mathbb{P}(\mathbb{T}_{i(\alpha)}^{\vec{C}}) \cap M$ with $q = \langle q_0, i(\alpha) \rangle$. Since the conditions p_0 and q_0 are compatible in $\mathbb{P}(\mathbb{T}_{i(\alpha)}^{\vec{C}})$, we know that $q_0(\beta) \neq 0$ holds for all $\beta \in \text{dom}(q_0)$ with $\beta <_{\mathbb{T}_{i(\alpha)}^{\vec{C}}} \alpha$. Since $\alpha = \text{otp}(D_\alpha)$ is a cardinal and $\text{dom}(q_0)$ is a finite subset of α , there is a $\gamma \in \text{acc}(D_\alpha)$ with $\text{dom}(q_0) \subseteq \gamma$. Then $\gamma \in \text{acc}(C_{\alpha, i(\alpha)})$, $i(\gamma) \leq i(\alpha)$ and $\gamma <_{\mathbb{T}_{i(\alpha)}^{\vec{C}}} \alpha$. Moreover, the above remarks show that $r = \langle q_0 \cup \{\langle \gamma, 0 \rangle\}, i(\alpha) \rangle \in M$ is a condition in $\mathbb{P}_{\vec{C}}$ that strengthens q . But this implies that the conditions p and r are compatible in $\mathbb{P}_{\vec{C}}$, a contradiction. \square

The proof of the following result is similar to that of Theorem 1.3 presented in Section 5.

Theorem 7.7. *Let κ be an inaccessible cardinal with $\mathbb{1}_{\text{Add}(\kappa, 1)} \Vdash \text{“}\kappa \text{ is weakly compact”}$. If $\lambda < \kappa$ is an infinite, regular cardinal, then the following statements hold in a cofinality-preserving forcing extension of the ground model:*

- (1) *There is a κ -Knaster partial order \mathbb{P} such that \mathbb{P}^μ is κ -Knaster for all $\mu < \lambda$, but \mathbb{P}^λ does not satisfy the κ -chain condition.*
- (2) *If \mathbb{R} is a partial order with the property that \mathbb{R}^λ satisfies the κ -chain condition, then \mathbb{R}^θ satisfies the κ -chain condition for all $\theta < \kappa$.*

Proof. Let $\mathbb{P} = \mathbb{P}(\kappa, \lambda)$ be the forcing notion from Definition 4.7 that adds a $\square^{\text{ind}}(\kappa, \lambda)$ -sequence, let \dot{C} be a \mathbb{P} -name for the generically-added $\square^{\text{ind}}(\kappa, \lambda)$ -sequence, and, for all $i < \lambda$, let \dot{Q}_i be a \mathbb{P} -name for the partial order $\mathbb{T}_i(\dot{C})$ defined in Definition 4.9. If $i < \lambda$, then our assumptions imply that κ is weakly compact in all $(\mathbb{P} * \dot{Q}_i)$ -generic extensions of V .

Let G be \mathbb{P} -generic over V , and let $\dot{C}^G = \vec{C} = \langle C_{\alpha, i} \mid \alpha < \kappa, i(\alpha) \leq i < \lambda \rangle$ be the realization of \dot{C} . Given $i < j < \lambda$, set $\mathbb{Q}_i = \dot{Q}_i^G$ and let $\pi_{i, j} : \mathbb{Q}_i \rightarrow \mathbb{Q}_j$ be the projection map given by $\pi_{i, j}(C_{\alpha, i}) = C_{\alpha, j}$. We claim that $V[G]$ is the desired forcing extension. By Lemma 4.8, the set $E_{\geq \lambda}^\kappa \cap S_{\geq i}^{\vec{C}}$ is a stationary subset of κ in $V[G]$ for all $i < \lambda$. In this situation, Lemma 7.6 shows that the partial order $\mathbb{P}_{\vec{C}}$ from Definition 7.5 witnesses that the above statement (1) holds.

Let us now show that requirement (2) holds in $V[G]$. To this end, work in $V[G]$ and fix a partial order \mathbb{R} such that \mathbb{R}^λ satisfies the κ -chain condition.

Claim. *There is an $i < \lambda$ and a condition q in \mathbb{Q}_i such that*

$$q \Vdash_{\mathbb{Q}_i} \text{“}\check{\mathbb{R}} \text{ satisfies the } \check{\kappa}\text{-chain condition”}.$$

Proof of the Claim. Suppose not. Given $i < \lambda$, this assumption yields a sequence $\langle \dot{r}_{i, \eta} \mid \eta < \kappa \rangle$ of \mathbb{Q}_i -names for elements of \mathbb{R} such that

$$\mathbb{1}_{\mathbb{Q}_i} \Vdash \text{“The conditions } \dot{r}_{i, \eta} \text{ and } \dot{r}_{i, \xi} \text{ are incompatible in } \check{\mathbb{R}} \text{”}$$

for all $\eta < \xi < \kappa$. For each $\eta < \kappa$, use Lemma 4.11 to find $q_\eta \in \mathbb{Q}_0$ and a sequence $\langle r_{i,\eta} \mid i < \lambda \rangle$ of conditions in \mathbb{R} such that $\pi_{0,i}(q_\eta) \Vdash_{\mathbb{Q}_i} \text{“}\dot{r}_{i,\eta} = \check{r}_{i,\eta}\text{”}$ for all $i < \lambda$. Given $\eta < \kappa$, pick $\alpha_\eta \in \text{acc}(\kappa)$ with $q_\eta = C_{\alpha_\eta,0}$.

For each $\eta < \lambda$, let s_η denote the unique condition in \mathbb{R}^λ with $s_\eta(i) = r_{i,\eta}$ for all $i < \lambda$. Fix $\eta, \xi < \lambda$ with $\alpha_\eta < \alpha_\xi$ and $i < \lambda$ with $\alpha_\eta \in \text{acc}(C_{\alpha_\xi,j})$ for all $i \leq j < \lambda$. Given $i \leq j < \lambda$, we then have $\pi_{0,i}(q_\xi) \leq_{\mathbb{Q}_i} \pi_{0,i}(q_\eta)$ and therefore

$$\pi_{0,i}(q_\xi) \Vdash_{\mathbb{Q}_i} \text{“}\dot{r}_{i,\eta} = \check{r}_{i,\eta} \text{ and } \dot{r}_{i,\xi} = \check{r}_{i,\xi}\text{”}.$$

In particular, the conditions $s_\eta(i)$ and $s_\xi(i)$ are incompatible in \mathbb{R} , and therefore the conditions s_η and s_ξ are incompatible in \mathbb{R}^λ . But this shows that $\{s_\eta \mid \eta < \kappa\}$ is an antichain in \mathbb{R}^λ of size κ , contradicting our assumption that \mathbb{R}^λ satisfies the κ -chain condition. \square

Fix i and q as given in the claim, and $\theta < \kappa$. Let H be \mathbb{Q}_i -generic over $V[G]$ with $q \in H$. In $V[G, H]$, κ is weakly compact and \mathbb{R} satisfies the κ -chain condition. In particular, arguments discussed in Section 1 show that \mathbb{R}^θ is κ -Knaster in $V[G, H]$. Since \mathbb{Q}_i is $<\kappa$ -distributive in $V[G]$, we have $(\mathbb{R}^\theta)^{V[G]} = (\mathbb{R}^\theta)^{V[G,H]}$. Moreover, since the property of satisfying the κ -chain condition is easily seen to be downward absolute, we can conclude that \mathbb{R}^θ satisfies the κ -chain condition in $V[G]$. \square

REFERENCES

1. James E. Baumgartner, *Iterated forcing*, Surveys in set theory, London Math. Soc. Lecture Note Ser., vol. 87, Cambridge Univ. Press, Cambridge, 1983, pp. 1–59. MR 823775 (87c:03099)
2. Ari Meir Brodsky and Assaf Rinot, *Reduced powers of Souslin trees*, Forum Math. Sigma **5** (2017), e2, 82. MR 3596614
3. Sean Cox, *Layered posets and Kunen’s universal collapse*, to appear in the *Notre Dame Journal of Formal Logic*.
4. Sean Cox and Philipp Lücke, *Characterizing large cardinals in terms of layered posets*, Ann. Pure Appl. Logic **168** (2017), no. 5, 1112–1131. MR 3620068
5. James Cummings, *Souslin trees which are hard to specialise*, Proc. Amer. Math. Soc. **125** (1997), no. 8, 2435–2441. MR 1376756 (97j:03095)
6. ———, *Iterated forcing and elementary embeddings*, Handbook of set theory. Vols. 1, 2, 3, Springer, Dordrecht, 2010, pp. 775–883. MR 2768691
7. James Cummings, Matthew Foreman, and Menachem Magidor, *Squares, scales and stationary reflection*, J. Math. Log. **1** (2001), no. 1, 35–98. MR 1838355
8. James Cummings and Ernest Schimmerling, *Indexed squares*, Israel J. Math. **131** (2002), 61–99. MR 1942302
9. Keith J. Devlin, *Reduced powers of \aleph_2 -trees*, Fund. Math. **118** (1983), no. 2, 129–134. MR 732661 (85i:03156)
10. Yair Hayut and Chris Lambie-Hanson, *Simultaneous stationary reflection and square sequences*, J. Math. Log. **17** (2017), no. 2, 1750010, 27. MR 3730566
11. R. Björn Jensen, *The fine structure of the constructible hierarchy*, Ann. Math. Logic **4** (1972), 229–308; erratum, *ibid.* **4** (1972), 443, With a section by Jack Silver. MR 0309729
12. Kenneth Kunen, *Saturated ideals*, J. Symbolic Logic **43** (1978), no. 1, 65–76. MR 495118 (80a:03068)
13. ———, *Set theory*, Studies in Logic and the Foundations of Mathematics, vol. 102, North-Holland Publishing Co., Amsterdam-New York, 1980, An introduction to independence proofs. MR 597342
14. Chris Lambie-Hanson, *Squares and covering matrices*, Ann. Pure Appl. Logic **165** (2014), no. 2, 673–694. MR 3129734
15. ———, *Aronszajn trees, square principles, and stationary reflection*, MLQ Math. Log. Q. **63** (2017), no. 3-4, 265–281. MR 3724382
16. ———, *Squares and narrow systems*, J. Symb. Log. **82** (2017), no. 3, 834–859. MR 3694332

17. Richard Laver and Saharon Shelah, *The \aleph_2 -Souslin hypothesis*, Trans. Amer. Math. Soc. **264** (1981), no. 2, 411–417. MR 603771
18. Philipp Lücke, *Ascending paths and forcings that specialize higher Aronszajn trees*, Fund. Math. **239** (2017), no. 1, 51–84. MR 3667758
19. William Mitchell, *Aronszajn trees and the independence of the transfer property*, Ann. Math. Logic **5** (1972/73), 21–46. MR 0313057
20. Assaf Rinot, *Chain conditions of products, and weakly compact cardinals*, Bull. Symb. Log. **20** (2014), no. 3, 293–314. MR 3271280
21. Assaf Shani, *Fresh subsets of ultrapowers*, Arch. Math. Logic **55** (2016), no. 5-6, 835–845. MR 3523658
22. Saharon Shelah and Lee Stanley, *Weakly compact cardinals and nonspecial Aronszajn trees*, Proc. Amer. Math. Soc. **104** (1988), no. 3, 887–897. MR 964870
23. Stevo Todorčević, *Stationary sets, trees and continuums*, Publ. Inst. Math. (Beograd) (N.S.) **29(43)** (1981), 249–262. MR 657114
24. ———, *Partition relations for partially ordered sets*, Acta Math. **155** (1985), no. 1-2, 1–25. MR 793235 (87d:03126)
25. ———, *Partitioning pairs of countable ordinals*, Acta Math. **159** (1987), no. 3-4, 261–294. MR 908147
26. ———, *Special square sequences*, Proc. Amer. Math. Soc. **105** (1989), no. 1, 199–205. MR 929410 (89e:03082)
27. Stevo Todorčević and Víctor Torres Pérez, *Conjectures of Rado and Chang and special Aronszajn trees*, Math. Log. Q. **58** (2012), no. 4-5, 342–347. MR 2965421

DEPARTMENT OF MATHEMATICS AND APPLIED MATHEMATICS, VIRGINIA COMMONWEALTH UNIVERSITY, 1015 FLOYD AVENUE, RICHMOND, VIRGINIA 23284, USA

URL: <http://people.vcu.edu/~cblambiehanso/>

MATHEMATISCHES INSTITUT, UNIVERSITÄT BONN, ENDENICHER ALLEE 60, 53115 BONN, GERMANY

URL: <http://www.math.uni-bonn.de/people/pluecke/>