

# WEAK COMPACTNESS CARDINALS FOR STRONG LOGICS AND SUBTLETY PROPERTIES OF THE CLASS OF ORDINALS

PHILIPP LÜCKE

ABSTRACT. Motivated by recent work of Boney, Dimopoulos, Gitman and Magidor, we characterize the existence of weak compactness cardinals for all abstract logics through combinatorial properties of the class of ordinals. This analysis is then used to show that, in contrast to the existence of strong compactness cardinals, the existence of weak compactness cardinals for abstract logics does not imply the existence of a strongly inaccessible cardinal. More precisely, it is proven that the existence of a proper class of subtle cardinals is consistent with the axioms of ZFC if and only if it is not possible to derive the existence of strongly inaccessible cardinals from the existence of weak compactness cardinals for all abstract logics. Complementing this result, it is shown that the existence of weak compactness cardinals for all abstract logics implies that unboundedly many ordinals are strongly inaccessible in the inner model HOD of all hereditarily ordinal definable sets.

## 1. INTRODUCTION

The theory of strong axioms of infinity and the study of extensions of first-order logic are deeply connected through results showing that the ability to generalize fundamental structural features of first-order logic to stronger logics is equivalent to the existence of large cardinals.<sup>1</sup> A classical example of such a result is a theorem of Magidor in [14] that shows that the existence of a *strong compactness cardinal* for second-order logic  $\mathcal{L}^2$  (i.e., the existence of a cardinal  $\kappa$  with the property that every unsatisfiable  $\mathcal{L}^2$ -theory contains an unsatisfiable subtheory of cardinality less than  $\kappa$ ) is equivalent to the existence of an extendible cardinal. In [16], Makowsky proved an analog of this result that provides a large cardinal characterization for the existence of strong compactness cardinals for all abstract logics.<sup>2</sup> Recall that *Vopěnka's Principle* is the scheme of axioms stating that for every proper class<sup>3</sup> of graphs, there are two distinct members of the class with a homomorphism between them. The validity of this combinatorial principle is known to be characterizable through the existence of certain large cardinals (see [1, Section 4] and [21, Section 6]). Makowsky's result can then be stated as follows:

---

2020 *Mathematics Subject Classification.* (Primary) 03B16; (Secondary) 03C55, 03E45, 03E55.

*Key words and phrases.* Abstract logics, large cardinals, weak compactness cardinals, strict Löwenheim–Skolem–Tarski numbers, ordinal definability.

The author would like to thank Will Boney and Victoria Gitman for discussions that motivated much of the content of this paper. In addition, he would like to thank Toshimichi Usuba for helpful comments on earlier versions of the presented results. The author gratefully acknowledges supported by the Deutsche Forschungsgemeinschaft (Project number 522490605).

<sup>1</sup>Throughout this paper, we refer to properties of cardinal numbers that imply weak inaccessibility as large cardinal properties.

<sup>2</sup>The definition of abstract logics and all related model-theoretic notions can be found in Section 5.

<sup>3</sup>Throughout this paper, we work in ZFC. Therefore, all classes are definable by first-order formulas in the language of set theory, possibly using sets as parameters.

**Theorem 1.1** ([16, Theorem 2]). *The following schemes are equivalent over ZFC:*

- (1) *Vopěnka’s Principle.*
- (2) *Every abstract logic has a strong compactness cardinal.*

The work presented in this paper is motivated by recent results of Boney, Dimopoulos, Gitman and Magidor in [5] that deal with the existence of *weak compactness cardinals* for abstract logics, i.e., the existence of cardinals  $\kappa$  with the property that every unsatisfiable theory of cardinality  $\kappa$  contains an unsatisfiable subtheory of cardinality less than  $\kappa$ . These results show that the existence of weak compactness cardinals for strong logics is connected to the combinatorial concept of *subtleness*, introduced by Jensen and Kunen in [11]. Remember that an infinite cardinal  $\kappa$  is *subtle* if for every sequence  $\langle A_\alpha \mid \alpha < \kappa \rangle$  with  $A_\alpha \subseteq \alpha$  for all  $\alpha < \kappa$  and every closed unbounded subset  $C$  of  $\kappa$ , there are  $\alpha, \beta \in C$  with  $\alpha < \beta$  and  $A_\beta \cap \alpha = A_\alpha$ . A short argument shows that subtle cardinals are inaccessible limits of totally indescribable cardinals. This large cardinal notion has a canonical class-version, i.e., one defines “*Ord is subtle*” to be the schemes of sentences stating that for every class sequence  $\langle A_\alpha \mid \alpha \in \text{Ord} \rangle$  with  $A_\alpha \subseteq \alpha$  for all  $\alpha \in \text{Ord}$  and every closed unbounded class  $C$  of ordinals, there exist  $\alpha, \beta \in C$  with  $\alpha < \beta$  and  $A_\beta \cap \alpha = A_\alpha$ . The results of [5] now show that, in the presence of a definable well-ordering of  $V$ ,<sup>4</sup> the validity of this principle is connected to the existence of weak compactness cardinals for abstract logics:

**Theorem 1.2** ([5, Theorem 5.3]). *The following schemes are equivalent over ZFC together with the existence of a definable well-ordering of  $V$ :*

- (1) *Ord is subtle.*
- (2) *Every abstract logic has a stationary class of weak compactness cardinals.*

This result immediately raises the following questions:

- Is it also possible to characterize the mere existence of weak compactness cardinals for every abstract logic (instead of the existence of stationary classes of such cardinals) through combinatorial properties of the class of ordinals?
- Is it possible to establish such characterizations in the absence of definable well-orderings of  $V$ ?

The results of this paper will provide affirmative answers to both of these questions. Moreover, this analysis will be based on arguments showing that the validity of the involved combinatorial principles for the class of ordinals is equivalent to the existence of certain large cardinals.

First, in order to prove results that do not rely on the existence of definable well-orderings, we will make use of the following variation of the above class-principle that was introduced by Bagaria and the author in their work on *principles of structural reflection*:

**Definition 1.3** ([2]). *We let “Ord is essentially subtle” denote the scheme of sentences stating that for every class sequence  $\langle E_\alpha \mid \alpha \in \text{Ord} \rangle$  with  $\emptyset \neq E_\alpha \subseteq \mathcal{P}(\alpha)$  for all  $\alpha \in \text{Ord}$  and every closed unbounded class  $C$  of ordinals, there exist  $\alpha, \beta \in C$  and  $A \in E_\beta$  with  $\alpha < \beta$  and  $A \cap \alpha \in E_\alpha$ .*

Obviously, if Ord is essentially subtle, then Ord is subtle. Moreover, in the presence of a definable well-ordering of  $V$ , these principles are equivalent. In particular, since the assumption that Ord is subtle is easily seen to be downwards absolute from  $V$  to the constructible universe  $L$ , it follows that these two principles are equiconsistent over ZFC. In contrast, by combining a result in [5] with Theorem 1.4 below, it is possible to show that, over ZFC, the assumption that Ord is subtle does not

<sup>4</sup>Note that the existence of such a well-ordering is equivalent to the statement that  $V = \text{HOD}_x$  for some set  $x$  and can therefore be formulated as a single sentence in the language of set theory.

imply that  $\text{Ord}$  is essentially subtle (see Corollary 6.3 below). The following direct strengthening of Theorem 1.2, proven in Section 6 below, now provides the desired ZFC-characterization of the existence of stationary classes of weak compactness cardinals for all abstract logics:

**Theorem 1.4.** *The following schemes of sentences are equivalent over ZFC:*

- (1) *Ord is essentially subtle.*
- (2) *Every abstract logic has a stationary class of weak compactness cardinals.*

Next, we present a characterization of the existence of weak compactness cardinals for all abstract logics in terms of combinatorial properties of the class  $\text{Ord}$ . This characterization is based on the notion of *faintness*, introduced by Matet in [17]. Matet defined a cardinal  $\kappa$  to be *faint* if for every sequence  $\langle A_\alpha \mid \alpha < \kappa \rangle$  with  $A_\alpha \subseteq \alpha$  for all  $\alpha < \kappa$  and every  $\xi < \kappa$ , there are  $\xi < \alpha < \beta < \kappa$  with  $A_\beta \cap \alpha = A_\alpha$ . The results of [17] then show that a cardinal is faint if and only if it is either subtle or a limit of subtle cardinals. Following earlier patterns, we introduce the following class-version of faintness:

**Definition 1.5.** *We let “Ord is essentially faint” denote the scheme of sentences stating that for every class sequence  $\langle E_\alpha \mid \alpha \in \text{Ord} \rangle$  with  $\emptyset \neq E_\alpha \subseteq \mathcal{P}(\alpha)$  for all  $\alpha \in \text{Ord}$  and every ordinal  $\xi$ , there are ordinals  $\xi < \alpha < \beta$  and  $A \in E_\beta$  with  $A \cap \alpha \in E_\alpha$ .*

By definition, if  $\text{Ord}$  is essentially subtle, then  $\text{Ord}$  is essentially faint. Moreover, it is easy to see that the existence of a proper class of subtle cardinals implies that  $\text{Ord}$  is essentially faint, because, given a class sequence  $\langle E_\alpha \mid \alpha \in \text{Ord} \rangle$  with  $\emptyset \neq E_\alpha \subseteq \mathcal{P}(\alpha)$  for all  $\alpha \in \text{Ord}$  and an ordinal  $\xi$ , we can pick a subtle cardinal  $\kappa > \xi$  and use the Axiom of Choice to find a sequence  $\langle A_\alpha \mid \alpha < \kappa \rangle$  with  $A_\alpha \in E_\alpha$  for all  $\alpha < \kappa$ . The following result, again proven in Section 6, shows that this principle exactly determines the combinatorial property of  $\text{Ord}$  needed for the desired characterization:

**Theorem 1.6.** *The following schemes of sentences are equivalent over ZFC:*

- (1) *Ord is essentially faint.*
- (2) *Every abstract logic has a weak compactness cardinal.*

As already mentioned earlier, the proofs of the above theorems rely heavily on characterizations of the involved combinatorial class-principles through large cardinal assumptions. This analysis, carried out in Sections 2 and 3 below, can also be used to show that the ability to generalize other structural properties of first-order logic to all abstract logics can also be characterized through the validity of the studied combinatorial principles for  $\text{Ord}$ . More specifically, motivated by an unpublished result of Stavi that provides an analog to Theorem 1.1 for the existence of *Löwenheim–Skolem–Tarski numbers* for abstract logics (see Theorem 5.4 below), we will prove analogs of Theorems 1.4 and 1.6 that characterize the existence of *strict Löwenheim–Skolem–Tarski numbers*, studied by Bagaria and Väänänen in [3], through subtlety properties of  $\text{Ord}$  (see Theorem 5.5).

The results presented above naturally raise questions about the relationship between the principle “Ord is essentially subtle” and the principle “Ord is essentially faint”. Trivially, the former principle implies the latter. However, it is not obvious whether these principles are equivalent. Somewhat surprisingly, the next result shows that the inequivalence of the two principles has consistency strength strictly greater than the consistency strength of the individual principles. Moreover, this result uses the inequivalence of the two principles to derive the existence of certain large cardinals not only in the constructible universe  $L$ , but also in the class  $\text{HOD}$  of all hereditarily ordinal definable sets.

**Theorem 1.7.** *If Ord is essentially faint and only boundedly many cardinals are subtle in HOD, then Ord is essentially subtle.*

In particular, should it be the case that the axioms of ZFC prove that there are only set-many subtle cardinals, then these axioms also prove the equivalence of the two principles. Moreover, since the existence of a subtle cardinal clearly implies the existence of a set-sized model of ZFC in which Ord is essentially subtle, the above result directly yields the following equiconsistency:

**Corollary 1.8.** *The following theories are equiconsistent:*

- (1) ZFC + “Ord is essentially subtle”.
- (2) ZFC + “Ord is essentially faint”.

□

Motivated by Theorem 1.7, we now consider the possibility of separating the principle “Ord is essentially faint” from the principle “Ord is essentially subtle”. Our results will show that this question is closely related to questions about the existence of sentences in the language of set theory that imply all sentences appearing in the given schemes of sentences. To motivate these results, we will first observe that, over ZFC, the principle “Ord is essentially subtle” cannot be derived from a single consistent sentence. In Section 2, we will show that this observation directly follows from the results of [2, Section 9] that show that for every natural number  $n$ , the assumption that Ord is essentially subtle implies the existence of class-many strongly inaccessible cardinals  $\kappa$  with the property that  $V_\kappa$  is a  $\Sigma_n$ -elementary submodel of  $V$ .

**Proposition 1.9.** *If  $\phi$  is a sentence in the language of set theory with the property that ZFC +  $\phi$  is consistent, then*

$$\text{ZFC} + \phi \not\vdash \text{“Ord is essentially subtle”}.$$

In contrast, assuming the consistency of sufficiently strong large cardinal assumptions, it is easy to find consistent sentences that provably imply that Ord is essentially faint. As observed above, the statement that there is a proper class of subtle cardinals is an example of a sentence with this property. Our further analysis will yield additional examples of such sentences. In particular, we will produce two incompatible sentences with the given property (see Theorem 1.14 below). The following theorem now shows that the existence of such a consistent sentence is actually equivalent to the possibility of separating the two principles. Moreover, it shows that Theorem 1.7 already provides the correct consistency strength for the inequality of these principles.

**Theorem 1.10.** *The following statements are equivalent:*

- (1) ZFC + “Ord is essentially faint”  $\not\vdash$  “Ord is essentially subtle”.
- (2) *There exists a sentence  $\phi$  in the language of set theory such that the theory ZFC +  $\phi$  is consistent and*

$$\text{ZFC} + \phi \vdash \text{“Ord is essentially faint”}.$$

- (3) *The theory*

$$\text{ZFC} + \text{“There is a proper class of subtle cardinals”}$$

*is consistent.*

Next, we consider the possibility of axiomatizing the validity of the given principles by single sentences. Proposition 1.9 already shows that this is not possible for the principle stating that Ord is essentially subtle. A short argument in Section 4 will prove the following result that provides the same conclusion for the principle “Ord is essentially faint”.

**Corollary 1.11.** *If the theory*

$$\text{ZFC} + \text{“Ord is essentially faint”}$$

*is consistent, then no sentence  $\phi$  in the language of set theory satisfies the following statements:*

- (1)  $\text{ZFC} + \text{“Ord is essentially faint”} \vdash \phi$ .
- (2)  $\text{ZFC} + \phi \vdash \text{“Ord is essentially faint”}$ .

We now consider the question whether the validity of the scheme “Ord is essentially subtle” can be finitely axiomatized over the theory  $\text{ZFC} + \text{“Ord is essentially faint”}$ . The following observation, proven in Section 4, motivates the formulation of the subsequent theorem:

**Proposition 1.12.** *There exists a theory T in the language of set theory such that the following statements hold:*

- (1)  $\text{ZFC} + \text{“Ord is essentially subtle”} \vdash \text{T}$ .
- (2)  $\text{ZFC} + \text{“Ord is essentially faint”} + \text{T} \vdash \text{“Ord is essentially subtle”}$ .
- (3) *For every sentence  $\phi$  in T, we have*

$$\text{ZFC} + \neg\phi \vdash \text{“Ord is essentially faint”}.$$

It is now natural to ask whether the theory T appearing in the above proposition can be replaced by a single sentence. The next result, also proven in Section 4, shows that the answer to this question depends on the consistency of the class-version of the large cardinal property of being a subtle limit of subtle cardinals:

**Theorem 1.13.** *The following statements are equivalent:*

- (1) *There is no sentence  $\phi$  in the language of set theory satisfying the following statements:*
  - (a)  $\text{ZFC} + \text{“Ord is essentially subtle”} \vdash \phi$ .
  - (b)  $\text{ZFC} + \text{“Ord is essentially faint”} + \phi \vdash \text{“Ord is essentially subtle”}$ .
  - (c)  $\text{ZFC} + \neg\phi \vdash \text{“Ord is essentially faint”}$ .
- (2) *There is no sentence  $\phi$  in the language of set theory satisfying the above statements (1a) and (1b).*
- (3) *The theory*

$$\text{ZFC} + \text{“There is a proper class of subtle cardinals”} + \text{“Ord is essentially subtle”}$$

*is consistent.*

The final topic that will be treated in this paper is the relationship between the existence of weak compactness for abstract logics and the existence of strongly inaccessible cardinals. By combining Theorem 1.1 with the results of [1, Section 4], it is easy to see that the existence of strong compactness cardinals for all abstract logics implies the existence of a proper class of strongly inaccessible cardinals. Moreover, a combination of Theorem 1.4 with results in [2, Section 9] (see Theorem 2.4 below) shows that the same conclusion follows from the assumption that every abstract logic has a stationary class of weak compactness cardinals. In contrast, it turns out that the existence of weak compactness cardinals for abstract logics is compatible with the non-existence of strongly inaccessible cardinals. The corresponding arguments will also allow us to show that there can be incompatible sentences witnessing (2) in Theorem 1.10.

**Theorem 1.14.** *The following statements are equivalent:*

(1) *The theory*

ZFC + “There is a proper class of subtle cardinals”

*is consistent.*

(2) *The theory*

ZFC + “Ord is essentially faint” + “There are no strongly inaccessible cardinals”

*is consistent.*

(3) *There is a sentence  $\phi$  in the language of set theory such that ZFC +  $\phi$  is consistent and*

*ZFC +  $\phi \vdash$  “Ord is essentially faint” + “There are no strongly inaccessible cardinals”.*

Notice that, in combination with Theorem 1.6, the above theorem shows that the existence of a weak compactness cardinal for second-order logic does not imply the existence of a strongly inaccessible cardinal and this shows that *weakly sub-extendible* cardinals, introduced by Fuchino and Sakai in their analysis of weak compactness cardinals for  $\mathcal{L}^2$  in [8], can exist in models of set theory that do not contain strongly inaccessible cardinals.

Finally, while Theorem 1.14 shows that the existence of weak compactness cardinals for all abstract logics does not imply the existence of strongly inaccessible cardinals, our analysis allows us to conclude that the given assumption at least implies the existence of strongly inaccessible cardinals in the inner model HOD:

**Theorem 1.15.** *If Ord is essentially faint, then unboundedly many ordinals are strongly inaccessible cardinals in HOD.*

Note that the proof of this result will actually show that strongly inaccessible cardinals with strong combinatorial properties exist in HOD (see Corollary 4.5 below).

## 2. $C^{(n)}$ -STRONGLY UNFOLDABLE CARDINALS

In this section, we review the large cardinal characterization of the principle “Ord is essentially subtle” derived in [2, Section 9]. The starting point of these results is the notion of *strongly unfoldable cardinal* introduced by Villaveces in [22]. Villaveces defined an inaccessible cardinal  $\kappa$  to be *strongly unfoldable* if for every ordinal  $\lambda$  and every transitive  $\text{ZF}^-$ -model  $M$  of cardinality  $\kappa$  with  $\kappa \in M$  and  ${}^{<\kappa}M \subseteq M$ , there is a transitive set  $N$  with  $V_\lambda \subseteq N$  and an elementary embedding  $j : M \rightarrow N$  with  $\text{crit}(j) = \kappa$  and  $j(\kappa) \geq \lambda$ . In [2], Bagaria and the author introduced natural strengthenings of this definition that demand the existence of embeddings into models with stronger correctness properties. Following [1], for every natural number  $n$ , we let  $C^{(n)}$  denote the class of all  $\Sigma_n$ -correct ordinals, i.e., the class of all ordinals  $\alpha$  with the property that  $V_\alpha$  is a  $\Sigma_n$ -elementary submodel of the set-theoretic universe  $V$ , denoted by  $V_\alpha \prec_{\Sigma_n} V$ . It is then easy to show that for each natural number  $n$ , the class  $C^{(n)}$  is a closed unbounded class of ordinals that is definable by a  $\Pi_n$ -formula without parameters. Moreover, short arguments show that  $C^{(0)}$  is the class of all ordinals and  $C^{(1)}$  is the class of all cardinals  $\kappa$  satisfying  $H(\kappa) = V_\kappa$ . In particular, there is a sentence  $\varphi$  in the language of set theory with the property that  $C^{(1)}$  is equal to the class of all ordinals  $\alpha$  with the property that  $\varphi$  holds in  $V_\alpha$ .

**Definition 2.1** ([2]). *Given a natural number  $n$ , an inaccessible cardinal  $\kappa$  is  $C^{(n)}$ -strongly unfoldable if for every ordinal  $\lambda \in C^{(n)}$  greater than  $\kappa$  and every transitive  $\text{ZF}^-$ -model  $M$  of cardinality  $\kappa$  with  $\kappa \in M$  and  ${}^{<\kappa}M \subseteq M$ , there is a transitive set  $N$  with  $V_\lambda \subseteq N$  and an elementary embedding  $j : M \rightarrow N$  with  $\text{crit}(j) = \kappa$ ,  $j(\kappa) > \lambda$  and  $V_\lambda \prec_{\Sigma_n} V_{j(\kappa)}^N$ .*

The following basic properties of these large cardinal notions were established in [2, Section 9]:

- Proposition 2.2** ([2]). (1) *A cardinal is strongly unfoldable if and only if it is  $C^{(n)}$ -strongly unfoldable for some natural number  $n < 2$ .*  
(2) *Given natural numbers  $m < n$ , every  $C^{(n)}$ -strongly unfoldable cardinal is  $C^{(m)}$ -strongly unfoldable.*  
(3) *For every natural number  $n$ , all  $C^{(n)}$ -strongly unfoldable cardinals are elements of  $C^{(n+1)}$ .*

In [19], Rathjen defined a cardinal  $\kappa$  to be *shrewd* if for every formula  $\Phi(v_0, v_1)$  in the language of set theory, every ordinal  $\gamma > \kappa$  and every  $A \subseteq V_\kappa$  with the property that  $\Phi(A, \kappa)$  holds in  $V_\gamma$ , there exist ordinals  $\alpha < \beta < \kappa$  such that  $\Phi(A \cap V_\alpha, \alpha)$  holds in  $V_\beta$ . The results of [12] then show that the notions of shrewdness and strong unfoldability coincide. The proof of this equivalence relies on an embedding characterization of shrewdness in [13] that closely resembles Magidor's classical characterization of supercompactness in [14]. The following result establishes analogous equivalences for the notion of  $C^{(n)}$ -strong unfoldability (see [2, Theorem 9.4]):

**Lemma 2.3** ([2]). *Given a natural number  $n > 0$ , the following statements are equivalent for every cardinal  $\kappa$ :*

- (1) *The cardinal  $\kappa$  is  $C^{(n)}$ -strongly unfoldable.*
- (2) *For every  $\mathcal{L}_\in$ -formula  $\varphi(v_0, v_1)$ , every ordinal  $\gamma \in C^{(n)}$  greater than  $\kappa$ , and every subset  $A$  of  $V_\kappa$  with the property that  $\varphi(\kappa, A)$  holds in  $V_\gamma$ , there exist ordinals  $\alpha < \beta < \kappa$  with  $\beta \in C^{(n)}$  and the property that  $\varphi(\alpha, A \cap V_\alpha)$  holds in  $V_\beta$ .*
- (3) *For every ordinal  $\gamma \in C^{(n)}$  greater than  $\kappa$  and every  $z \in V_\gamma$  there exists an ordinal  $\bar{\gamma} \in C^{(n)} \cap \kappa$ , a cardinal  $\bar{\kappa} < \bar{\gamma}$ , an elementary submodel  $X$  of  $V_{\bar{\gamma}}$  with  $V_{\bar{\kappa}} \cup \{\bar{\kappa}\} \subseteq X$ , and an elementary embedding  $j : X \rightarrow V_\gamma$  with  $j \upharpoonright \bar{\kappa} = \text{id}_{\bar{\kappa}}$ ,  $j(\bar{\kappa}) = \kappa$  and  $z \in \text{ran}(j)$ .*

Analogous to the characterization of Vopěnka's Principle through the existence of  $C^{(n)}$ -extendible cardinals provided by the results of [1, Section 4], it turns out that the existence  $C^{(n)}$ -strongly unfoldable cardinals for all natural numbers  $n$  characterizes the validity of the principle “Ord is essentially subtle” (see [2, Theorem 1.17]).

**Theorem 2.4** ([2]). *The following schemes of axioms are equivalent over ZFC:*

- (1) *Ord is essentially subtle.*
- (2) *For every natural number  $n$ , there exists a  $C^{(n)}$ -strongly unfoldable cardinal.*
- (3) *For every natural number  $n$ , there exists a proper class of  $C^{(n)}$ -strongly unfoldable cardinals.*

The above results now allow us to show that no consistent sentence can imply that Ord is essentially subtle. Note that an almost identical argument shows that Vopěnka's Principle is not the consequence of a single consistent sentence.

*Proof of Proposition 1.9.* Assume, towards a contradiction, that there is a sentence  $\phi$  in the language of set theory such that the theory  $\text{ZFC} + \phi$  is consistent and proves that Ord is essentially subtle. Work in a model of  $\text{ZFC} + \phi$ . By combining Theorem 2.4 with Proposition 2.2.(3), we can now find an inaccessible cardinal  $\kappa$  with the property that  $\phi$  holds in  $V_\kappa$ . Let  $\kappa$  be minimal with this property. Since  $V_\kappa$  is a model of  $\text{ZFC} + \phi$ , we can repeat this argument in  $V_\kappa$  and find an inaccessible cardinal  $\mu < \kappa$  such that  $\phi$  holds in  $V_\mu$ , contradicting the minimality of  $\kappa$ .  $\square$

### 3. $C^{(n)}$ -WEAKLY SHREWD CARDINALS

In this section, we will formulate and prove an analog of Theorem 2.4 for the principle “Ord is essentially faint”. The starting point of this analysis is the notion of *weak shrewdness* introduced

in [13] as a natural weakening of shrewdness that does not imply the strong inaccessibility of the given cardinal. In [13], a cardinal  $\kappa$  is defined to be *weakly shrewd* if for every formula  $\varphi(v_0, v_1)$  in the language of set theory, every cardinal  $\theta > \kappa$  and every subset  $z$  of  $\kappa$  with the property that  $\varphi(z, \kappa)$  holds in  $H(\theta)$ , there exist cardinals  $\bar{\kappa} < \bar{\theta}$  such that  $\bar{\kappa} < \kappa$  and  $\varphi(z \cap \bar{\kappa}, \bar{\kappa})$  holds in  $H(\bar{\theta})$ . The results of [13, Section 3] then show that every shrewd cardinal is weakly shrewd and every weakly shrewd cardinal is weakly Mahlo. Moreover, these results show that, starting from a subtle cardinal, it is possible to construct a model of set theory that contains a weakly shrewd cardinal that is not strongly inaccessible and therefore also not shrewd. Moreover, it is shown that weakly shrewd cardinals can even exist below the cardinality of the continuum. The results of the paper will show that these consistency results make use of the correct large cardinal hypothesis (see Lemma 4.3 below). Motivated by Lemma 2.3, we strengthen the notion of weak shrewdness in the same way as the transition to  $C^{(n)}$ -strong unfoldability strengthens the notions of shrewdness.

**Definition 3.1.** *Given a natural number  $n$ , a cardinal  $\kappa$  is  $C^{(n)}$ -weakly shrewd if for every formula  $\varphi(v_0, v_1)$  in the language of set theory, every cardinal  $\kappa < \theta \in C^{(n)}$  and every subset  $z$  of  $\kappa$  with the property that  $\varphi(z, \kappa)$  holds in  $H(\theta)$ , there exist a cardinal  $\bar{\theta} \in C^{(n)}$  and a cardinal  $\bar{\kappa} < \min(\kappa, \bar{\theta})$  such that  $\varphi(z \cap \bar{\kappa}, \bar{\kappa})$  holds in  $H(\bar{\theta})$ .*

We now derive some basic properties of this newly defined notion:

**Proposition 3.2.** (1) *Given natural numbers  $m < n$ , every  $C^{(n)}$ -weakly shrewd cardinal is  $C^{(m)}$ -weakly shrewd.*

(2) *A cardinal is weakly shrewd if and only if it is  $C^{(n)}$ -weakly shrewd for some  $n < 2$ .*

*Proof.* (1) Let  $\kappa$  be a  $C^{(n)}$ -weakly shrewd cardinal. Fix a formula  $\varphi(v_0, v_1)$  in the language of set theory, a cardinal  $\kappa < \theta \in C^{(m)}$  and  $z \subseteq \kappa$  such that  $\varphi(z, \kappa)$  holds in  $H(\theta)$ . Pick a cardinal  $\theta < \vartheta \in C^{(n)}$ . Then, in  $H(\vartheta)$ , there exists a cardinal  $\kappa < \nu \in C^{(m)}$  with the property that  $\varphi(z, \kappa)$  holds in  $H(\nu)$ . By our assumptions on  $\kappa$ , we can now find a cardinal  $\bar{\vartheta} \in C^{(n)}$  and a cardinal  $\bar{\kappa} < \min(\kappa, \bar{\vartheta})$  such that, in  $H(\bar{\vartheta})$ , there exists a cardinal  $\bar{\kappa} < \bar{\nu} \in C^{(m)}$  with the property that  $\varphi(z \cap \bar{\kappa}, \bar{\kappa})$  holds in  $H(\bar{\nu})$ . Since  $\bar{\vartheta} \in C^{(n)}$  and  $n > 0$ , we then know that, in  $V$ , the ordinal  $\bar{\theta}$  is a cardinal in  $C^{(m)}$  with the property that  $\varphi(z \cap \bar{\kappa}, \bar{\kappa})$  holds in  $H(\bar{\nu})$ .

(2) By definition and (1), it suffices to show that every weakly shrewd cardinal is  $C^{(1)}$ -weakly shrewd. Let  $\kappa$  be a weakly shrewd cardinal, let  $\varphi(v_0, v_1)$  be a formula in the language of set theory, let  $\kappa < \theta \in C^{(1)}$  be a cardinal and let  $z$  be a subset of  $\kappa$  with the property that  $\varphi(z, \kappa)$  holds in  $H(\theta)$ . Since  $\theta$  is an element of  $C^{(1)}$ , we then know that, in  $H(\theta)$ , for every ordinal  $\alpha$ , the class  $V_\alpha$  is a set. Our assumption then allows us to find a cardinal  $\bar{\theta}$  and a cardinal  $\bar{\kappa} < \min(\kappa, \bar{\theta})$  with the property that, in  $H(\bar{\theta})$ , the statement  $\varphi(z \cap \bar{\kappa}, \bar{\kappa})$  holds and for every ordinal  $\alpha$ , the class  $V_\alpha$  is a set. This completes the proof, because this conclusion ensures that  $\bar{\theta}$  is an element of  $C^{(1)}$ .  $\square$

In combination with [13, Proposition 3.3], the above proposition shows that all weakly  $C^{(n)}$ -shrewd cardinals are weakly Mahlo. In particular, if a  $C^{(n)}$ -weakly shrewd cardinal is an element of  $C^{(1)}$ , then it is strongly inaccessible. Motivated by [13, Lemma 3.1], we now prove an analog of Lemma 2.3 for  $C^{(n)}$ -weak shrewdness.

**Lemma 3.3.** *Given a natural number  $n > 0$ , the following statements are equivalent for every cardinal  $\kappa$ :*

(1) *The cardinal  $\kappa$  is  $C^{(n)}$ -weakly shrewd.*



- (2) For every cardinal  $\kappa < \theta \in C^{(n)}$  and every  $y \in H(\theta)$ , there exists a cardinal  $\bar{\theta} \in C^{(n)}$ , a cardinal  $\bar{\kappa} < \min(\kappa, \bar{\theta})$ , an elementary submodel  $X$  of  $H(\bar{\theta})$  with  $\bar{\kappa} + 1 \subseteq X$  and an elementary embedding  $j : X \rightarrow H(\theta)$  with  $j \upharpoonright \bar{\kappa} = \text{id}_{\bar{\kappa}}$ ,  $j(\bar{\kappa}) = \kappa$  and  $y \in \text{ran}(j)$ .

*Proof.* First, assume that (1) holds,  $\kappa < \theta \in C^{(n)}$  is a cardinal and  $y$  is an element of  $H(\theta)$ . Pick a cardinal  $\theta < \vartheta \in C^{(n)}$  and an elementary submodel  $Y$  of  $H(\theta)$  of cardinality  $\kappa$  with the property that  $\kappa \cup \{\kappa, z\} \subseteq Y$ . In addition, fix a bijection  $b : \kappa \rightarrow Y$  satisfying  $b(0) = \kappa$ ,  $b(1) = \langle \kappa, z \rangle$  and  $b(\omega \cdot (1 + \alpha)) = \alpha$  for all  $\alpha < \kappa$ . Finally, define  $z$  to be the set of all elements of  $\kappa$  of the form  $\langle \ell, \alpha_0, \dots, \alpha_{n-1} \rangle$ <sup>5</sup> with the property that  $\ell < \omega$  is the Gödel number of a set-theoretic formula with  $n$  free variables,  $\alpha_0, \dots, \alpha_{n-1} < \kappa$  and  $\text{Sat}(Y, \ell, \langle b(\alpha_0), \dots, b(\alpha_{n-1}) \rangle)$  holds, where  $\text{Sat}$  denotes the formalized satisfaction relation (see, for example, [7, Section I.9]).

By our assumption and the fact that  $\text{Sat}$  is uniformly  $\Delta_1$ -definable over models of  $\text{ZF}^-$ , we can now find a cardinal  $\bar{\vartheta} \in C^{(n)}$  and a cardinal  $\bar{\kappa} < \min(\kappa, \bar{\vartheta})$  with the property that, in  $H(\bar{\vartheta})$ , there exists a cardinal  $\bar{\kappa} < \bar{\theta} \in C^{(n)}$ , an element  $\bar{z}$  of  $H(\bar{\theta})$ , an elementary submodel  $X$  of  $H(\bar{\theta})$  of cardinality  $\bar{\kappa}$  with  $\bar{\kappa} \cup \{\bar{\kappa}, \bar{z}\} \subseteq X$  and a bijection  $a : \bar{\kappa} \rightarrow X$  with  $a(0) = \bar{\kappa}$ ,  $a(\omega \cdot (1 + \alpha)) = \alpha$  for all  $\alpha < \bar{\kappa}$  and the property that  $z \cap \bar{\kappa}$  consists of all elements of  $\kappa$  of the form  $\langle \ell, \alpha_0, \dots, \alpha_{n-1} \rangle$  such that  $\ell < \omega$  is the Gödel number of a set-theoretic formula with  $n$  free variables,  $\alpha_0, \dots, \alpha_{n-1} < \bar{\kappa}$  and  $\text{Sat}(X, \ell, \langle a(\alpha_0), \dots, a(\alpha_{n-1}) \rangle)$  holds. Since  $\bar{\vartheta} \in C^{(n)}$ , we now know that, in  $V$ , the ordinal  $\bar{\theta}$  is a cardinal in  $C^{(n)}$  and the set  $X$  is an elementary submodel of  $H(\bar{\theta})$ . If we now define

$$j = b \circ a^{-1} : X \rightarrow H(\theta),$$

then it is easy to check that  $j$  is an elementary embedding with  $j \upharpoonright \bar{\kappa} = \text{id}_{\bar{\kappa}}$ ,  $j(\bar{\kappa}) = \kappa$  and  $y \in \text{ran}(j)$ .

Now, assume that (2) holds,  $\varphi(v_0, v_1)$  is a formula in the language of set theory,  $\kappa < \theta \in C^{(n)}$  is a cardinal and  $z$  is a subset of  $\kappa$  with the property that  $\varphi(z, \kappa)$  holds in  $H(\theta)$ . Our assumptions then allows us to find a cardinal  $\bar{\theta} \in C^{(n)}$ , a cardinal  $\bar{\kappa} < \min(\kappa, \bar{\theta})$ , an elementary submodel  $X$  of  $H(\bar{\theta})$  with  $\bar{\kappa} + 1 \subseteq X$  and an elementary embedding  $j : X \rightarrow H(\theta)$  with  $j \upharpoonright \bar{\kappa} = \text{id}_{\bar{\kappa}}$ ,  $j(\bar{\kappa}) = \kappa$  and  $z \in \text{ran}(j)$ . We then know that  $z \cap \bar{\kappa} \in X$  and  $j(z \cap \bar{\kappa}) = z$ . But, this shows that  $\varphi(z \cap \bar{\kappa}, \bar{\kappa})$  holds in both  $X$  and  $H(\bar{\theta})$ .  $\square$

Next, we observe that the existence of subtle cardinals entails the existence of many  $C^{(n)}$ -weakly shrewd cardinals:

**Lemma 3.4.** *Given a natural number  $n > 0$ , every subtle cardinal is a stationary limit of strongly inaccessible  $C^{(n)}$ -weakly shrewd cardinals.*

*Proof.* Assume, towards a contradiction, that there is a subtle cardinal  $\delta$  and a closed unbounded subset  $C$  of  $\delta$  that consists of uncountable cardinals that are not  $C^{(n)}$ -weakly shrewd. Given  $\gamma \in C$ , we can use Lemma 3.3 to find  $\ell_\gamma < \omega$  and  $z_\gamma \subseteq \gamma$  such that  $\ell_\gamma$  is the Gödel number of a set-theoretic formula  $\varphi(v_0, v_1)$  with the property that there exists a cardinal  $\gamma < \theta \in C^{(n)}$  such that  $\varphi(z, \gamma)$  holds in  $H(\theta)$  and for every cardinal  $\bar{\theta} \in C^{(n)}$ , there is no cardinal  $\bar{\gamma} < \min(\gamma, \bar{\theta})$  with the property that  $\varphi(z \cap \bar{\gamma}, \bar{\gamma})$  holds in  $H(\bar{\theta})$ . Now, let  $\langle E_\gamma \mid \gamma \in C \rangle$  denote the unique sequence with

$$E_\gamma = \langle 0, \ell_\gamma \rangle \cup \{ \langle 1, \beta \rangle \mid \beta \in z_\gamma \}$$

for all  $\gamma \in C$ . Using the subtlety of  $\delta$ , we can now find  $\beta < \gamma$  in  $C$  with  $E_\gamma \cap \beta = E_\beta$ . We then know that  $\ell_\beta = \ell_\gamma$  and  $z_\gamma \cap \beta = z_\beta$ . But, this contradicts the choice of  $\ell_\gamma$  and  $z_\gamma$ , because, if  $\varphi(v_0, v_1)$  is the formula coded by  $\ell_\gamma$ , then  $\beta$  is a cardinal below  $\gamma$  with the property that there exists  $\beta < \theta \in C^{(n)}$  such that  $\varphi(z_\gamma \cap \beta, \beta)$  holds in  $H(\theta)$ .

<sup>5</sup>We use  $\langle \cdot, \dots, \cdot \rangle$  to denote (iterated applications) of the Gödel pairing function.

Since subtle cardinals are regular limit points of the class  $C^{(1)}$ , the above computations show that subtle cardinals are stationary limits of  $C^{(n)}$ -weakly shrewd cardinals that are elements of  $C^{(1)}$ . This yields the statement of the lemma, because, as observed earlier,  $C^{(n)}$ -weakly shrewd cardinals are inaccessible if and only if they are elements of  $C^{(1)}$ .  $\square$

We close this section by proving the desired characterization of the principle “Ord is essentially faint” through the existence of weakly  $C^{(n)}$ -shrewd cardinals:

**Theorem 3.5.** *The following schemes of axioms are equivalent over ZFC:*

- (1) *Ord is essentially faint.*
- (2) *For every natural number  $n$ , there exists a proper class of  $C^{(n)}$ -weakly shrewd cardinals.*

*Proof.* First, assume, towards a contradiction, that (1) holds,  $n > 0$  is a natural number and there exists an ordinal  $\xi$  with the property that there are no weakly  $C^{(n)}$ -shrewd cardinals above  $\xi$ . Then there is a unique class function  $E$  with domain Ord such that the following statements hold:

- $E(\alpha) = \mathcal{P}(\alpha)$  for all ordinal  $\alpha \leq \xi$ .
- $E(\alpha + 1) = \{\{<0, \alpha>\}\}$  for all ordinals  $\alpha \geq \xi$ .
- If  $\alpha > \xi$  is a singular limit ordinal, then  $E(\alpha)$  consists of all subsets of  $\alpha$  of the form

$$\{<1, \text{cof}(\alpha)>\} \cup \{<2, \eta, c(\eta)> \mid \eta < \text{cof}(\alpha)\},$$

where  $c : \text{cof}(\alpha) \rightarrow \alpha$  is a strictly increasing cofinal function.

- If  $\alpha > \xi$  is a regular cardinal, then  $E(\alpha)$  consists of all subsets of  $\alpha$  of the form

$$\{<3, \ell>\} \cup \{<4, \eta> \mid \eta \in z\},$$

where  $\ell$  is the Gödel number of a set-theoretic formula  $\varphi(v_0, v_1)$  and  $z$  is a subset of  $\alpha$  with the property that there exists a cardinal  $\alpha < \theta \in C^{(n)}$  such that  $\varphi(z, \alpha)$  holds in  $H(\theta)$  and for every cardinal  $\bar{\theta} \in C^{(n)}$ , there is no cardinal  $\bar{\alpha} < \min(\alpha, \bar{\theta})$  with the property that  $\varphi(z \cap \bar{\alpha}, \bar{\alpha})$  holds in  $H(\bar{\theta})$ .

Our assumption then ensures that  $E(\alpha) \neq \emptyset$  holds for all  $\alpha \in \text{Ord}$ . Therefore, we can find ordinals  $\xi < \alpha < \beta$  and  $A \in E(\beta)$  with  $A \cap \alpha \in E(\alpha)$ . The definition of  $E$  then ensures that  $\alpha$  and  $\beta$  are both regular cardinals. Fix a set-theoretic formula  $\varphi(v_0, v_1)$  and a subset  $z$  of  $\beta$  such that  $A = \{<3, \ell>\} \cup \{<4, \eta> \mid \eta \in z\}$ , where  $\ell$  is the Gödel number of  $\varphi(v_0, v_1)$ . Since

$$A \cap \alpha = \{<3, \ell>\} \cup \{<4, \eta> \mid \eta \in z \cap \alpha\} \in E(\alpha),$$

we now know that there exists a cardinal  $\alpha < \theta \in C^{(n)}$  with the property that  $\varphi(z \cap \alpha, \alpha)$  holds in  $H(\theta)$ . Since  $\alpha < \min(\beta, \theta)$ , this contradicts the fact that  $A$  is an element of  $E(\beta)$ .

Now, assume that (2) holds and  $E$  is a class function with domain Ord and the property that  $\emptyset \neq E(\alpha) \subseteq \mathcal{P}(\alpha)$  holds for all  $\alpha \in \text{Ord}$ . Fix an ordinal  $\xi$ . Pick a natural number  $n > 0$  such that there exists a  $\Sigma_n$ -formula  $\varphi(v_0, v_1, v_2)$  and a parameter  $y$  defining  $E$ . By our assumption, there is a  $C^{(n)}$ -weakly shrewd cardinal  $\kappa > \xi$  with  $y \in H(\kappa)$ . Pick  $A \in E(\kappa)$  and a cardinal  $\kappa < \theta \in C^{(n)}$ . Then Lemma 3.3 yields a cardinal  $\bar{\theta} \in C^{(n)}$ , a cardinal  $\bar{\kappa} < \min(\kappa, \bar{\theta})$ , an elementary submodel  $X$  of  $H(\bar{\theta})$  with  $\bar{\kappa} + 1 \subseteq X$  and an elementary embedding  $j : X \rightarrow H(\theta)$  with  $j \upharpoonright \bar{\kappa} = \text{id}_{\bar{\kappa}}$ ,  $j(\bar{\kappa}) = \kappa$  and  $A, y, \xi \in \text{ran}(j)$ . We then know that  $\xi < \bar{\kappa}$ ,  $y \in H(\bar{\kappa}) \cap X$  with  $j(y) = y$  and  $A \cap \bar{\kappa} \in X$  with  $j(A \cap \bar{\kappa}) = A$ . Moreover, the fact that  $\theta \in C^{(n)}$  ensures that, in  $H(\theta)$ , there exists a set  $\mathcal{A}$  such that  $A \in \mathcal{A}$  and  $\varphi(\kappa, \mathcal{A}, y)$  holds. This implies that, in  $H(\bar{\theta})$ , there exists a set  $\bar{\mathcal{A}}$  such that  $A \cap \bar{\kappa} \in \bar{\mathcal{A}}$  and  $\varphi(\bar{\kappa}, \bar{\mathcal{A}}, y)$  holds. Since  $\bar{\theta} \in C^{(n)}$ , we then know that  $\varphi(\bar{\kappa}, \bar{\mathcal{A}}, y)$  also holds in  $V$  and we can conclude that  $A \cap \bar{\kappa} \in \bar{\mathcal{A}} = E(\bar{\kappa})$ .  $\square$

## 4. SEPARATING THE SCHEMES

The goal of this section is to identify scenarios in which the principle “Ord is essentially faint” holds while the principle “Ord is essentially subtle” fails. In the light of Theorems 2.4 and 3.5, this task is closely connected to the analysis of  $C^{(n)}$ -weakly shrewd cardinals that are not  $C^{(n)}$ -strongly unfoldable. For the case  $n = 1$ , this analysis is provided by [13, Lemma 3.5], which shows that a weakly shrewd cardinal is strongly unfoldable if and only if it is an element of  $C^{(2)}$ . The following lemma provides a direct analog of this result for  $C^{(n)}$ -weakly shrewd cardinals:

**Lemma 4.1.** *Given a natural number  $n > 0$ , the following statements are equivalent for every  $C^{(n)}$ -weakly shrewd cardinal:*

- (1) *The cardinal  $\kappa$  is  $C^{(n)}$ -strongly unfoldable.*
- (2) *The cardinal  $\kappa$  is an element of  $C^{(n+1)}$ .*
- (3) *There is no ordinal  $\varepsilon \geq \kappa$  with the property that the set  $\{\varepsilon\}$  is definable by a  $\Sigma_{n+1}$ -formula with parameters in  $H(\kappa)$ .*

*Proof.* First, assume that (1) fails. Then Lemma 2.3 shows that there is a set-theoretic formula  $\varphi(v_0, v_1)$  and a subset  $z$  of  $\kappa$  with the property that there exists a cardinal  $\kappa < \vartheta \in C^{(n)}$  such that  $\varphi(z, \kappa)$  holds in  $H(\vartheta)$  and there is no cardinal  $\bar{\theta} \in C^{(n)}$  such that  $\varphi(z \cap \bar{\kappa}, \bar{\kappa})$  holds in  $H(\bar{\theta})$  for some cardinal  $\bar{\kappa} < \bar{\theta}$ . Since  $\kappa$  is weakly  $C^{(n)}$ -shrewd, we can find a cardinal  $\bar{\theta} \in C^{(n)}$  and a cardinal  $\bar{\kappa} < \min(\kappa, \bar{\theta})$  such that  $\varphi(z \cap \bar{\kappa}, \bar{\kappa})$  holds in  $H(\bar{\theta})$ . Now, define  $\varepsilon$  to be the least element of  $C^{(n)}$  above  $\bar{\kappa}$  with the property that  $\varphi(z \cap \bar{\kappa}, \bar{\kappa})$  holds in  $H(\varepsilon)$ . We then know that  $\varepsilon \geq \kappa$  and it is easy to see that the set  $\{\varepsilon\}$  is definable by a  $\Sigma_{n+1}$ -formula with parameters  $\bar{\kappa}, z \cap \bar{\kappa} \in H(\kappa)$ . This shows that (3) fails and hence we know that (3) implies (1). This completes the proof of the lemma, because (2) obviously implies (3) and Proposition 2.2.(3) shows that (1) implies (2).  $\square$

**Corollary 4.2.** *Given a natural number  $n > 0$ , every subtle cardinal that is not a limit of subtle cardinals is a stationary limit of strongly inaccessible  $C^{(n)}$ -weakly shrewd cardinals that are not  $C^{(n)}$ -strongly unfoldable.*

*Proof.* Let  $\delta$  be a subtle cardinal that is not a limit of subtle cardinals and let  $\gamma < \delta$  be an ordinal with the property that the interval  $(\gamma, \delta)$  does not contain subtle cardinals. Then the set  $\{\delta\}$  can be defined by a  $\Sigma_2$ -formula with parameter  $\gamma$ . In this situation, Lemma 4.1 shows that no  $C^{(n)}$ -weakly shrewd cardinal in the interval  $(\gamma, \delta)$  is  $C^{(n)}$ -strongly unfoldable. An application of Lemma 3.4 now yields the statement of the corollary.  $\square$

The next lemma provides us with a tool to derive non-trivial consistency strength from the existence of  $C^{(n)}$ -weakly shrewd cardinals that are not  $C^{(n)}$ -strongly unfoldable. An intriguing aspect of this result is the fact that it establishes the existence of a subtle cardinal not only in some specific canonical inner model (e.g., in the constructible universe  $L$ ), but in all canonical inner models (e.g., all submodels of the class HOD).

**Lemma 4.3.** *Let  $n > 0$  be a natural number and let  $\kappa$  be a  $C^{(n)}$ -weakly shrewd cardinal that is not  $C^{(n)}$ -strongly unfoldable. If  $\delta$  is the least ordinal  $\varepsilon \geq \kappa$  with the property that the set  $\{\varepsilon\}$  is definable by a  $\Sigma_{n+1}$ -formula with parameters in  $H(\kappa)$ , then the following statements hold:*

- (1)  *$\delta$  is a cardinal greater than  $\kappa$  with  $\text{cof}(\delta) \in \kappa \cup \{\delta\}$ .*
- (2)  *$\delta$  is a subtle cardinal in HOD.*

*Proof.* Fix a  $\Sigma_{n+1}$ -formula  $\varphi(v_0, v_1)$  and an element  $y$  with the property that  $\delta$  is the unique set  $x$  such that  $\varphi(x, y)$  holds. Pick a cardinal  $\kappa < \theta \in C^{(n+1)}$  and use Lemma 3.3 find a cardinal

$\bar{\theta} \in C^{(n)}$ , a cardinal  $\bar{\kappa} < \min(\kappa, \bar{\theta})$ , an elementary submodel  $X$  of  $H(\bar{\theta})$  with  $\bar{\kappa} + 1 \subseteq X$  and an elementary embedding  $j : X \rightarrow H(\theta)$  with  $j \upharpoonright \bar{\kappa} = \text{id}_{\bar{\kappa}}$ ,  $j(\bar{\kappa}) = \kappa$  and  $y, \delta \in \text{ran}(j)$ . Then  $y \in H(\bar{\kappa})$  with  $j(y) = y$ . Pick  $\varepsilon \in X$  with  $j(\varepsilon) = \delta$ . Since  $\varphi(\delta, y)$  holds in  $H(\theta)$ , we then know that  $\varphi(\varepsilon, y)$  holds in  $H(\bar{\theta})$  and the fact  $\bar{\theta}$  is an element of  $C^{(n)}$  implies that  $\varphi(\varepsilon, y)$  also holds in  $V$ . This shows that  $\delta = \varepsilon$  and hence  $\delta > \kappa$ . Next, note that the set  $\{|\delta|\}$  is  $\Sigma_2$ -definable from the parameter  $\delta$  and, since  $|\delta| \geq \kappa$ , the minimality of  $\delta$  implies that  $\delta$  is a cardinal. Finally, since the set  $\{\text{cof}(\delta)\}$  is  $\Sigma_2$ -definable from the parameter  $\delta$ , the minimality of  $\delta$  also ensures that either  $\delta$  is regular or  $\text{cof}(\delta) < \kappa$  holds.

Now, assume that  $\delta$  is not a subtle cardinal in HOD. Let  $\langle \vec{A}, C \rangle$  denote the least pair in the canonical well-ordering of HOD with the property that  $\vec{A} = \langle A_\gamma \mid \gamma < \delta \rangle$  is a sequence of length  $\delta$  with  $A_\gamma \subseteq \gamma$  for all  $\gamma < \delta$  and  $C$  is a closed unbounded subset of  $\delta$  with the property that  $A_\gamma \cap \beta \neq A_\beta$  holds for all  $\beta < \gamma$  in  $C$ . Since the class of all proper initial segments of the canonical well-ordering of HOD is definable in  $V$  by a  $\Sigma_2$ -formula without parameters (see [10, Lemma 13.25]), we then know that, in  $V$ , the sets  $\{\vec{A}\}$  and  $\{C\}$  are both definable by  $\Sigma_2$ -formulas with parameter  $\delta$ , and hence these sets are also definable by  $\Sigma_{n+1}$ -formulas with parameter  $y$ . We then know that  $C \cap \kappa \neq \emptyset$ , because otherwise  $\kappa \leq \min(C) < \delta$  is an ordinal with the property that the set  $\{\min(C)\}$  is definable by a  $\Sigma_{n+1}$ -formula with parameters in  $H(\kappa)$ .

**Claim.**  $\kappa \in C$ .

*Proof of the Claim.* Assume, towards a contradiction, that  $\kappa$  is not an element of  $C$  and set  $\alpha = \max(C \cap \kappa) < \kappa$  and  $\beta = \min(C \setminus \kappa) = \min(C \setminus (\alpha + 1)) < \delta$ . Then  $\beta$  is an ordinal greater than  $\kappa$  with the property that the set  $\{\beta\}$  is definable by a  $\Sigma_{n+1}$ -formula with parameters  $\alpha, z \in H(\kappa)$  and hence it follows that  $\beta \geq \delta$ , a contradiction.  $\square$

Pick a cardinal  $\delta < \theta \in C^{(n+1)}$  and use Lemma 3.3 to find a cardinal  $\bar{\theta} \in C^{(n)}$ , a cardinal  $\bar{\kappa} < \min(\kappa, \bar{\theta})$ , an elementary submodel  $X$  of  $H(\bar{\theta})$  with  $\bar{\kappa} + 1 \subseteq X$  and an elementary embedding  $j : X \rightarrow H(\theta)$  with  $j \upharpoonright \bar{\kappa} = \text{id}_{\bar{\kappa}}$ ,  $j(\bar{\kappa}) = \kappa$  and  $y \in \text{ran}(j)$ . We then again know that  $y \in H(\bar{\kappa})$  with  $j(y) = y$  and, by the correctness properties of  $H(\theta)$  and  $H(\bar{\theta})$ , the fact that the sets  $\{\vec{A}\}$ ,  $\{C\}$  and  $\{\delta\}$  are all definable by  $\Sigma_{n+1}$ -formulas with parameter  $y$  implies that  $\vec{A}, C, \delta \in X$  with  $j(\vec{A}) = \vec{A}$ ,  $j(C) = C$  and  $j(\delta) = \delta$ . In this situation, elementarity allows us to conclude that  $\bar{\kappa} \in C$  and  $A_{\bar{\kappa}} \in X$  with  $j(A_{\bar{\kappa}}) = A_\kappa$  and  $A_\kappa \cap \bar{\kappa} = j(A_{\bar{\kappa}}) \cap \bar{\kappa} = A_{\bar{\kappa}}$ , contradicting the choice of  $\vec{A}$  and  $C$ .  $\square$

As a first application of the above lemma, we observe that, by combining it with Corollary 4.2, we can determine the exact consistency strength of the existence of a  $C^{(n)}$ -weakly shrewd cardinal that is not  $C^{(n)}$ -strongly unfoldable. The following corollary directly strengthens [13, Theorem 1.9.(i)]. Note that, in combination with Theorem 2.4, this result shows that, over ZFC, the existence of a  $C^{(n)}$ -weakly shrewd cardinal that is not  $C^{(n)}$ -strongly unfoldable has strictly greater consistency strength than the existence of a  $C^{(n)}$ -weakly shrewd cardinal.

**Corollary 4.4.** *Given a natural number  $n > 0$ , the following statements are equiconsistent over ZFC:*

- (1) *There exists a  $C^{(n)}$ -weakly shrewd cardinal that is not  $C^{(n)}$ -strongly unfoldable.*
- (2) *There exists a subtle cardinal.*  $\square$

Next, we use Lemma 4.3 to show that the principles “Ord is essentially subtle” and “Ord is essentially faint” are equivalent if only boundedly many cardinals are subtle in HOD.

*Proof of Theorem 1.7.* Assume, towards a contradiction that Ord is essentially faint and not essentially subtle. By Theorem 2.4, the fact that Ord is not essentially subtle implies that for some natural number  $n > 0$ , there are no  $C^{(n)}$ -strongly unfoldable cardinals. Since Ord is essentially faint, Theorem 3.5 now shows that there is a proper class of  $C^{(n)}$ -weakly shrewd cardinals that are not  $C^{(n)}$ -strongly unfoldable. In this situation, we can apply Lemma 4.3 to conclude that a proper class of cardinals is subtle in HOD.  $\square$

We now continue by using Theorem 1.7 to show that the principle “Ord is essentially faint” is not finitely axiomatizable over ZFC.

*Proof of Corollary 1.11.* Assume, towards a contradiction, that the theory

$$\text{ZFC} + \text{“Ord is essentially faint”}$$

is consistent and there is a sentence  $\phi$  in the language of set theory with the property that the statements (1) and (2) listed in the corollary hold. By combining Theorem 1.7 with Theorem 2.4, it follows that the theory

$$\text{ZFC} + \text{“Ord is essentially subtle”} + \text{“There are no subtle cardinals in HOD”}$$

is also consistent. Define  $\psi$  to be the conjunction of  $\phi$  with the statement that only boundedly many cardinals are subtle in HOD. Then (1) together with the above observations ensures that  $\text{ZFC} + \psi$  is consistent. Moreover, by (2), we can apply Theorem 1.7 to conclude that

$$\text{ZFC} + \psi \vdash \text{“Ord is essentially subtle”}$$

holds, contradicting Proposition 1.9.  $\square$

Another application of Theorem 1.7 yields the following strengthening of Theorem 1.15:

**Corollary 4.5.** *Given a natural number  $n$ , if Ord is essentially faint, then unboundedly many ordinals are strongly inaccessible  $C^{(n)}$ -weakly shrewd cardinals in HOD.*

*Proof.* If Ord is essentially subtle, then Ord is essentially subtle in HOD and an application of Theorem 2.4 in HOD yields the desired conclusion. Hence, we may assume that Ord is not essentially subtle. Then Theorem 1.7 shows that, in HOD, there is a proper class of subtle cardinals. An application of Lemma 3.4 now allows us to conclude that unboundedly many ordinals are inaccessible  $C^{(n)}$ -weakly shrewd cardinals in HOD.  $\square$

We now continue by using the developed techniques to show that the consistency strength of the inequality of the principles “Ord is essentially subtle” and “Ord is essentially faint” is equal to the existence of a proper class of subtle cardinals.

*Proof of Theorem 1.10.* First, assume that

$$\text{ZFC} + \text{“Ord is essentially faint”} \not\vdash \text{“Ord is essentially subtle”}$$

holds. Then Theorem 2.4 yields a natural number  $n > 0$  with the property that the theory

$$\text{ZFC} + \text{“Ord is essentially faint”} + \text{“There are no } C^{(n)}\text{-strongly unfoldable cardinals”}$$

is consistent and we can work in a model of this theory. An application of Theorem 1.7 now directly shows that, in HOD, there is a proper class of subtle cardinals. These computations show that (1) implies (3) in the statement of the theorem.

Next, let  $\phi$  denote the set-theoretic sentence stating that there is a proper class of subtle cardinals. A combination of Lemma 3.4 with Theorem 3.5 then shows that the theory  $\text{ZFC} + \phi$  proves that  $\text{Ord}$  is essentially faint. This shows that (3) implies (2) in the statement of the theorem.

Finally, assume that there is a sentence  $\phi$  such that the theory  $\text{ZFC} + \phi$  is consistent and

$$\text{ZFC} + \phi \vdash \text{“Ord is essentially faint”}.$$

Then Proposition 1.9 ensures that

$$\text{ZFC} + \phi \not\vdash \text{“Ord is essentially subtle”}$$

and hence we can conclude that

$$\text{ZFC} + \text{“Ord is essentially faint”} \not\vdash \text{“Ord is essentially subtle”}.$$

This shows that (2) implies (1) in the statement of the theorem.  $\square$

In order to expand our analysis of the given class principles, we now explore a phenomenon already unveiled by [13, Theorem 1.11] showing that the existence of weakly shrewd that are not strongly unfoldable yields reflection properties for statements of arbitrary high complexities:

**Lemma 4.6.** *Given natural numbers  $n > m > 0$ , if  $\kappa$  is a  $C^{(m)}$ -weakly shrewd cardinal that is not  $C^{(m)}$ -strongly unfoldable,  $\delta > \kappa$  is a cardinal with the property that the set  $\{\delta\}$  is definable by a  $\Sigma_{m+1}$ -formula with parameters in  $\text{H}(\kappa)$  and  $\alpha < \kappa$  is an ordinal, then the interval  $(\alpha, \delta)$  contains a  $C^{(n)}$ -weakly shrewd cardinal that is not  $C^{(n)}$ -strongly unfoldable.*

*Proof.* Pick  $y \in \text{H}(\kappa)$  with the property that the set  $\{\delta\}$  is definable by a  $\Sigma_{m+1}$ -formula with parameter  $y$ . Since  $\kappa$  is a limit cardinal, we can find a cardinal  $\alpha < \rho < \kappa$  with  $y \in \text{H}(\rho)$ . Moreover, since  $n > m$ , it follows that the interval  $(\rho, \delta)$  does not contain an element of  $C^{(n+1)}$ . Now, assume, towards a contradiction, the interval  $(\rho, \delta)$  does not contain a  $C^{(n)}$ -weakly shrewd cardinal that is not  $C^{(n)}$ -strongly unfoldable. The above observation then allow us to use Lemma 4.1 to conclude that the interval  $(\rho, \delta)$  does not contain  $C^{(n)}$ -weakly shrewd cardinals. This shows that for every cardinal  $\mu$  in the interval  $(\rho, \delta)$ , there exists a set-theoretic formula  $\varphi(v_0, v_1)$ , a cardinal  $\mu < \theta \in C^{(n)}$  and a subset  $z$  of  $\mu$  such that  $\varphi(z, \mu)$  holds in  $\text{H}(\theta)$  and  $\varphi(z \cap \bar{\mu}, \bar{\mu})$  does not hold in  $\text{H}(\bar{\theta})$  for all  $\bar{\theta} \in C^{(n)}$  and all cardinals  $\bar{\mu} < \min(\mu, \bar{\theta})$ . We can then find a cardinal  $\delta < \zeta \in C^{(1)}$  with the property that, in  $\text{H}(\zeta)$ , for every cardinal  $\mu$  in the interval  $(\rho, \delta)$ , there exists a set-theoretic formula  $\varphi(v_0, v_1)$ , a cardinal  $\mu < \theta \in C^{(n)}$  and a subset  $z$  of  $\mu$  such that  $\varphi(z, \mu)$  holds in  $\text{H}(\theta)$  and  $\varphi(z \cap \bar{\mu}, \bar{\mu})$  does not hold in  $\text{H}(\bar{\theta})$  for all  $\bar{\theta} \in C^{(n)}$  and all cardinals  $\bar{\mu} < \min(\mu, \bar{\theta})$ . Let  $\eta$  be the least cardinal with this property. Then the set  $\{\eta\}$  is definable by a  $\Sigma_2$ -formula with parameters  $\delta$  and  $y$ . We then directly know that the set  $\{\eta\}$  is also definable by a  $\Sigma_{m+1}$ -formula with parameter  $y$ .

Now, pick  $\kappa < \vartheta \in C^{(m+1)}$  and use Lemma 3.3 to find  $\bar{\vartheta} \in C^{(m)}$ , a cardinal  $\bar{\kappa} < \min(\kappa, \bar{\vartheta})$ , an elementary submodel  $X$  of  $\text{H}(\bar{\vartheta})$  with  $\bar{\kappa} + 1 \subseteq X$  and an elementary embedding  $j : X \rightarrow \text{H}(\bar{\vartheta})$  with  $j \upharpoonright \bar{\kappa} = \text{id}_{\bar{\kappa}}$ ,  $j(\bar{\kappa}) = \kappa$  and  $y, \rho \in \text{ran}(j)$ . Then  $y \in X$  with  $j(y) = y$ . Moreover, we know that  $\rho < \bar{\kappa}$  and therefore  $j(\rho) = \rho$ . In combination with the facts that all  $\Sigma_{m+1}$ -statements are upwards absolute from  $X$  to  $V$ , this shows that  $\delta, \eta \in X$  with  $j(\delta) = \delta$  and  $j(\eta) = \eta$ . Since  $\bar{\kappa}$  is a cardinal in the interval  $(\rho, \delta)$  in  $\text{H}(\eta)$ , we can now find a set-theoretic formula  $\varphi(v_0, v_1)$ , an ordinal  $\bar{\kappa} < \theta \in X \cap \eta$  and  $z \in \mathcal{P}(\bar{\kappa}) \cap X$  with the property that, in  $\text{H}(\eta)$ , the ordinal  $\theta$  is a cardinal in  $C^{(n)}$  such that  $\varphi(z, \bar{\kappa})$  holds in  $\text{H}(\theta)$  and  $\varphi(z \cap \mu, \mu)$  does not hold in  $\text{H}(\bar{\theta})$  for all  $\bar{\theta} \in C^{(n)}$  and all cardinals  $\mu < \min(\bar{\kappa}, \bar{\theta})$ . The fact that  $\text{H}(\eta) \in X$  with  $j(\text{H}(\eta)) = \text{H}(\eta)$  then allows us to use the elementarity of  $j$  to conclude that, in  $\text{H}(\eta)$ , the ordinal  $j(\theta)$  is a cardinal in  $C^{(n)}$  with the property that  $\varphi(j(z), \kappa)$  holds in  $\text{H}(j(\theta))$  and  $\varphi(j(z) \cap \mu, \mu)$  does not hold in  $\text{H}(\bar{\theta})$  for all  $\bar{\theta} \in C^{(n)}$  and all

cardinals  $\mu < \min(\kappa, \bar{\theta})$ . But, this yields a contradiction, because  $\theta$  is an element of  $C^{(n)}$  in  $H(\eta)$ ,  $\bar{\kappa} < \min(\kappa, \theta)$  and  $j(z) \cap \bar{\kappa} = z$ .  $\square$

As a first application of Lemma 4.6, we show that the principle “Ord is essentially subtle” can be axiomatized over the theory  $\text{ZFC} + \text{“Ord is essentially faint”}$  in a strong way.

*Proof of Proposition 1.12.* For each natural number  $n > 0$ , let  $\phi_n$  denote the sentence in the language of set theory that states that the existence of a proper class of  $C^{(n)}$ -weakly shrewd cardinals implies the existence of a proper class of  $C^{(n)}$ -strongly unfoldable cardinals. Define  $T$  to be the theory consisting of these sentences. Then Theorem 2.4 shows that  $T$  holds in every model of  $\text{ZFC}$  in which Ord is essentially subtle. Moreover, a combination of Theorem 2.4 and Theorem 3.5 shows that Ord is essentially subtle in every model of  $\text{ZFC}$  in which Ord is essentially faint and  $T$  holds. Finally, assume that we work in a model of  $\text{ZFC}$  in which the sentence  $\phi_m$  fails for some natural number  $m > 0$ . Then there exists a proper class of  $C^{(m)}$ -weakly shrewd cardinals and there are only boundedly many  $C^{(m)}$ -strongly unfoldable cardinals. In particular, there exists a proper class of  $C^{(m)}$ -weakly shrewd cardinals that are not  $C^{(m)}$ -strongly unfoldable. In this situation, a combination of Lemmas 4.1 and 4.6 shows that for every natural number  $n > 0$ , there exists a proper class of  $C^{(n)}$ -weakly shrewd cardinals. Hence, we can apply Theorem 3.5 to conclude that Ord is essentially faint in this model.  $\square$

By combing Theorem 1.7 with Proposition 1.12, we can now examine the conditions that allow the proposition’s statement to be strengthened to result in finite axiomatizations.

*Proof of Theorem 1.13.* First, assume that the theory

$$\text{ZFC} + \text{“There is a proper class of subtle cardinals”} + \text{“Ord is essentially subtle”}$$

is inconsistent and let  $\phi$  denote the set-theoretic sentence stating that only boundedly many cardinals are subtle in HOD. Now, if we work in a model  $V$  of  $\text{ZFC}$  in which Ord is essentially subtle, then Ord is essentially subtle in HOD and, since our assumption ensures that HOD has only boundedly many subtle cardinals, we can conclude that  $\phi$  holds in  $V$ . Next, if we work in a model of  $\text{ZFC}$  in which Ord is essentially faint and  $\phi$  holds, then Theorem 1.7 directly shows that Ord is essentially subtle. In combination, this shows that (2) implies (3) in the statement of the theorem.

Next, assume that there is a set-theoretic sentence  $\phi$  satisfying the statements (1a) and (1b) listed in the theorem. Let  $\psi$  be the set-theoretic sentence that is given by the conjunction of  $\phi$  and the sentence stating that there is a proper class of subtle cardinals. Then

$$\text{ZFC} + \psi \vdash \text{“Ord is essentially subtle”}$$

and therefore Proposition 1.9 shows that  $\text{ZFC} \vdash \neg\psi$ . In this situation, our assumptions imply that

$$\text{ZFC} + \text{“Ord is essentially subtle”} \vdash \text{“There are only boundedly many subtle cardinals”}$$

and we can conclude that (3) implies (2) in the statement of the theorem.

Finally, assume that there is a set-theoretic sentence  $\phi$  satisfying the statements (1a) and (1b) listed in the theorem. Let  $T$  be the theory given by Proposition 1.12. We then know that

$$\text{ZFC} + \text{“Ord is essentially faint”} + T \vdash \phi$$

and hence there is a finite subtheory  $F$  of  $T$  with

$$\text{ZFC} + \text{“Ord is essentially faint”} + F \vdash \phi.$$

Let  $\psi$  be the set-theoretic sentence that is given by the disjunction of  $\phi$  with the conjunction of all sentences in  $F$ . Our assumptions then ensure that  $\psi$  satisfies the statement (1a) of the theorem. Moreover, the definition of  $\psi$  implies that

$$\text{ZFC} + \text{“Ord is essentially faint”} + \psi \vdash \phi$$

and hence our assumption ensure that  $\psi$  satisfies statement (1b) of the theorem. Finally, assume that we work in a model of ZFC in which  $\psi$  does not hold. Then there exists a sentence in  $F$  that does not hold. In this situation, the properties of the theory  $T$  allow us to conclude that Ord is essentially faint in this model. This shows that  $\psi$  also satisfies statement (1c) of the theorem. In combination, these arguments show that (1) implies (2) in the statement of the theorem and therefore these assumptions are equivalent.  $\square$

We end this section by using Lemma 4.6 to show that the principle “Ord is essentially faint” does not imply the existence of strongly inaccessible cardinals.

*Proof of Theorem 1.14.* Work in a model of  $\text{ZFC} + V = L$  in which a proper class of subtle cardinals exists and no inaccessible cardinal is a limit of subtle cardinals. Let  $S$  denote the class of all subtle cardinals. Define  $R$  to be the unique class function with domain  $S$  that satisfies  $R(\min(S)) = \omega$  and  $R(\delta) = \sup(S \cap \delta)^+$  for all  $\min(S) < \delta \in S$ . Our assumptions then ensure that  $R$  is a well-defined regressive function. Define  $\mathbb{P}$  to be the class partial order that is given by the Easton support product of all partial orders of the form  $\text{Add}(R(\delta), \delta)$  with  $\delta \in S$  and let  $G$  be  $\mathbb{P}$ -generic over  $V$ . Then standard arguments (as in [10, pp. 235–236]) show that  $V[G]$  is a model of ZFC with the same regular cardinals as  $V$ .

**Claim.** *There are no inaccessible cardinals in  $V[G]$ .*

*Proof of the Claim.* Assume, towards a contradiction, that there is an inaccessible  $\kappa$  in  $V[G]$ . Set  $\delta = \min(S \setminus \kappa) \geq \kappa$ . Since  $\kappa$  is not a limit of subtle cardinals in  $V$ , we know that  $R(\delta) < \kappa$ . Moreover, our setup ensures that

$$(2^{R(\delta)})^{V[G]} \geq \delta \geq \kappa$$

holds, contradicting the inaccessibility of  $\kappa$  in  $V[G]$ .  $\square$

**Claim.** *In  $V[G]$ , there is a proper class of weakly shrewd cardinals.*

*Proof of the Claim.* Fix an ordinal  $\alpha > \omega$  and pick  $\delta \in S$  with  $R(\delta) > \alpha$ . Set  $S_0 = S \cap \delta$  and let  $\mathbb{Q}$  denote the Easton support product of all partial orders of the form  $\text{Add}(R(\gamma), \gamma)$  with  $\gamma \in S_0$ , as constructed in  $V$ . In addition, let  $\mathbb{R}$  denote the partial order  $\text{Add}(R(\delta), \delta)$  as constructed in  $V$ . A standard factor analysis of the class partial order  $\mathbb{P}$  (again, as in [10, pp. 235–236]) now allows us to find an inner model  $M$  of  $V[G]$  such that  $V \subseteq M$ ,  $({}^\delta V)^M \subseteq V$  and there are  $G_0, G_1 \in V[G]$  such that  $G_0 \times G_1$  is  $(\mathbb{Q} \times \mathbb{R})$ -generic over  $M$  with  $V[G] = M[G_0, G_1]$ . Then  $\delta$  is a subtle cardinal in  $M$  and the definition of  $R$  in  $V$  ensures that  $\delta$  is the minimal subtle cardinal above  $R(\delta)$  in  $M$ . In particular, it follows that the set  $\{\delta\}$  is definable by a  $\Sigma_2$ -formula with parameter  $R(\delta)$  in  $M$ .

An application of Lemma 3.4 now shows that, in  $M$ , the interval  $(R(\delta), \delta)$  contains an inaccessible weakly shrewd cardinal  $\kappa$ . In the following, fix a cardinal  $\theta > \delta$  that is an element of  $C^{(1)}$  in  $V[G]$  and an element  $z$  of  $H(\theta)^{V[G]}$ . Pick a  $(\mathbb{Q} \times \mathbb{R})$ -name  $\dot{z}$  in  $M$  with  $z = \dot{z}^{G_0 \times G_1}$  and an ordinal  $\vartheta > \theta$  that is an element of  $C^{(2)}$  in  $M$  and satisfies  $\dot{z} \in H(\vartheta)^M$ . Proposition 3.2 and Lemma 3.3 now show that, in  $M$ , we can find a cardinal  $\bar{\vartheta} \in C^{(1)}$ , a cardinal  $\bar{\kappa} < \min(\kappa, \bar{\vartheta})$ , an elementary submodel  $X$  of  $H(\bar{\vartheta})$  of cardinality  $\bar{\kappa}$  with  $\bar{\kappa} + 1 \subseteq X$  and an elementary embedding  $j : X \rightarrow H(\vartheta)$  with  $j \upharpoonright \bar{\kappa} = \text{id}_{\bar{\kappa}}$ ,  $j(\bar{\kappa}) = \kappa$  and  $R(\delta), \theta, \dot{z} \in \text{ran}(j)$ . We then know that  $R(\delta) < \bar{\kappa}$  and this directly implies



that  $\delta \in X$  with  $j(\delta) = \delta$ . In particular, it follows that  $\bar{\kappa} < \kappa < \delta < \bar{\vartheta}$ . Since the partial order  $\mathbb{Q} \times \mathbb{R}$  has cardinality  $\delta$  in  $M$ , we know that the cardinals  $\vartheta$  and  $\bar{\vartheta}$  are elements of  $C^{(1)}$  in both  $M[G_0]$  and  $V[G]$ . Moreover, since  $X$  has cardinality  $\bar{\kappa} < \delta$  in  $M$ , there is a permutation  $\sigma$  of  $\delta$  in  $M$  that extends the injection  $j \upharpoonright (X \cap \delta) : X \cap \delta \rightarrow \delta$ . Since  $\mathbb{R} = \text{Add}(R(\delta), \delta)^M$ , this shows that  $M$  also contains an automorphism  $\tau$  of  $\mathbb{R}$  that is induced by the action of  $\sigma$  on the supports of the conditions in  $\text{Add}(R(\delta), \delta)$ ,<sup>6</sup> in the sense that  $\text{dom}(\tau(p)) = \sigma[\text{dom}(p)]$  and  $\tau(p)(\sigma(\gamma)) = p(\gamma)$  holds for every condition  $p$  in  $\text{Add}(R(\delta), \delta)$  and all  $\gamma \in \text{dom}(p)$ . Since our setup ensures that  $\bar{\kappa}$  is inaccessible in  $M$  and  $H(\bar{\kappa})^M \subseteq X$  with  $j \upharpoonright H(\bar{\kappa})^M = \text{id}_{H(\bar{\kappa})^M}$ , it follows that  $\text{Add}(R(\delta), 1)^M \cup \{\text{Add}(R(\delta), 1)^M\} \subseteq X$  and  $j \upharpoonright \text{Add}(R(\delta), 1)^M = \text{id}_{\text{Add}(R(\delta), 1)^M}$ . Moreover, we also know that  $\mathbb{R} \in X$  with  $j(\mathbb{R}) = \mathbb{R}$ . Finally, our definitions ensure that  $\tau(p) = j(p)$  holds for every condition  $p$  in  $\mathbb{R}$  that is an element of  $X$ .

Next, since  $\mathcal{P}(\mathbb{Q}) \in H(\bar{\kappa})^M \subseteq X$ , we know that  $G_0$  is  $\mathbb{Q}$ -generic over  $X$ ,  $H(\vartheta)^M[G_0] = H(\vartheta)^{M[G_0]}$  and the model  $X[G_0]$  is an elementary submodel of  $H(\bar{\vartheta})^{M[G_0]}$ . Moreover, the fact that  $j \upharpoonright \mathbb{Q} = \text{id}_{\mathbb{Q}}$  implies that there is a canonical lifting

$$j_0 : X[G_0] \rightarrow H(\vartheta)^{M[G_0]}$$

of  $j$  to  $X[G_0]$ . Now, set  $G'_1 = \tau^{-1}[G_1] \subseteq \mathbb{R}$ . Since  $\tau$  is an element of  $H(\vartheta)^M$ , it follows that  $G'_1$  is  $\mathbb{R}$ -generic over  $M[G_0]$  with

$$H(\vartheta)^{M[G_0]}[G'_1] = H(\vartheta)^{M[G_0]}[G_1] = H(\vartheta)^{V[G]}.$$

Set  $G''_1 = G'_1 \cap X$ . Since the partial order  $\mathbb{Q}$  has cardinality less than  $\bar{\kappa}$  in  $M$  and the partial order  $\mathbb{R} = \text{Add}(R(\delta), \delta)$  satisfies the  $\bar{\kappa}$ -chain condition in  $M$ , it follows that  $\mathbb{R}$  satisfies the  $\bar{\kappa}$ -chain condition in  $M[G_0]$ . Together with the fact that  $\mathbb{R} \cap X = \mathbb{R} \cap X[G_0]$ , this ensures that  $G''_1$  is  $\mathbb{R}$ -generic over  $X[G_0]$  and  $X[G_0, G''_1]$  is an elementary submodel of  $H(\bar{\vartheta})^{V[G]}$ . Moreover, since we have ensured that  $j_0[G''_1] = j[(\tau^{-1}[G_1]) \cap X] \subseteq G_1$ , there is a canonical lifting

$$j_1 : X[G_0, G''_1] \rightarrow H(\vartheta)^{V[G]}$$

of  $j_0$  to  $X[G_0, G''_1]$ . Pick  $\bar{\theta} \in X$  with  $j(\bar{\theta}) = \theta$ . Since  $\bar{\vartheta}$  is an element of  $C^{(1)}$  in  $V[G]$ , elementarity implies that  $\bar{\theta}$  is also an element of this class. Set  $Y = H(\bar{\theta})^{V[G]} \cap X[G_0, G''_1]$  and define  $i = j_1 \upharpoonright Y$ . Since  $H(\bar{\theta})^{V[G]} \in X[G_0, G''_1]$ , elementarity ensures that the set  $Y$  is an elementary submodel of  $H(\bar{\theta})^{V[G]}$ . Moreover, our setup ensures that  $i : Y \rightarrow H(\bar{\theta})^{V[G]}$  is an elementary embedding with  $i \upharpoonright \bar{\kappa} = \text{id}_{\bar{\kappa}}$ ,  $i(\bar{\kappa}) = \kappa$ ,  $i(\delta) = \delta$  and  $z \in \text{ran}(i)$ . These computations show that, in  $V[G]$ , the cardinal  $\kappa > \alpha$  is weakly shrewd.  $\square$

In combination, the above claims show that  $V[G]$  is a model of ZFC that contains a proper class of weakly shrewd cardinals and in which no strongly inaccessible cardinals exist.

Now, let  $\phi$  denote the sentence in the language of set-theory stating that there are no strongly inaccessible cardinals and there is a proper class of weakly shrewd cardinals. Then  $\phi$  implies that there is a proper class of weakly shrewd cardinals that are not strongly unfoldable and, by Lemma 4.6, this conclusion ensures that for every natural number  $n > 0$ , there is a proper class of  $C^{(n)}$ -weakly shrewd cardinals. Therefore, Theorem 3.5 shows that

$$\text{ZFC} + \phi \vdash \text{“Ord is essentially faint”} + \text{“There are no strongly inaccessible cardinals”}.$$

In addition, the above computations show that the consistency of the theory

$$\text{ZFC} + \text{“There is a proper class of subtle cardinals”}$$

<sup>6</sup>Here, we view conditions in partial orders of the form  $\text{Add}(\mu, \nu)$  as functions  $p$  of cardinality less than  $\mu$  with  $\text{dom}(p) \subseteq \nu$  and  $p(\gamma) \in \text{Add}(\mu, 1)$  for every  $\gamma \in \text{dom}(p)$ .

implies the consistency of  $\text{ZFC} + \phi$ . Hence, we know that (1) implies (3) in the statement of the theorem.

Next, if we work in a model of  $\text{ZFC}$  in which  $\text{Ord}$  is essentially faint and there are no strongly inaccessible cardinals, then Theorem 2.4 shows that  $\text{Ord}$  is not essentially subtle and Theorem 1.7 allows us to conclude that a proper class of cardinals is subtle in  $\text{HOD}$ . In particular, we know that (2) implies (1) in the statement of the theorem.

This completes the proof of the theorem, because the remaining implication from (3) to (2) in its statement is trivial.  $\square$

## 5. ABSTRACT LOGICS

In the remainder of this paper, we will connect the theory developed above with the study of structural properties of abstract logics. We start by recalling the relevant definitions. Our setup is based on the definitions given in [6, Section 2.5] and our presentation closely follows those provided in [4], [5] and [9].

**Definition 5.1.** (1) A language is a tuple

$$\tau = \langle \mathfrak{C}_\tau, \mathfrak{F}_\tau, \mathfrak{R}_\tau, \mathfrak{a}_\tau \rangle,$$

where  $\mathfrak{C}_\tau$ ,  $\mathfrak{F}_\tau$  and  $\mathfrak{R}_\tau$  are pairwise disjoint sets and  $\mathfrak{a}_\tau : \mathfrak{F}_\tau \cup \mathfrak{R}_\tau \rightarrow \omega \setminus \{0\}$  is a function. We then call  $\mathfrak{C}_\tau$  the set of constant symbols of  $\tau$ ,  $\mathfrak{F}_\tau$  the set of function symbols of  $\tau$ ,  $\mathfrak{R}_\tau$  the set of relation symbols of  $\tau$ ,  $\mathfrak{a}_\tau$  the arity function of  $\tau$  and  $\mathfrak{S}_\tau = \mathfrak{C}_\tau \cup \mathfrak{F}_\tau \cup \mathfrak{R}_\tau$  the set of logical symbols of  $\tau$ .

(2) Given a language  $\tau$ , a  $\tau$ -structure is a tuple

$$M = \langle \text{dom}(M), \langle c^M \mid c \in \mathfrak{C}_\tau \rangle, \langle f^M \mid f \in \mathfrak{F}_\tau \rangle, \langle r^M \mid r \in \mathfrak{R}_\tau \rangle \rangle,$$

where  $\text{dom}(M)$  is a non-empty set, each  $c^M$  is an element of  $\text{dom}(M)$ , each  $f^M$  is an  $\mathfrak{a}_\tau(f)$ -ary function on  $\text{dom}(M)$  and each  $r^M$  is an  $\mathfrak{a}_\tau(r)$ -ary relation on  $\text{dom}(M)$ .

(3) Given a language  $\tau$ , we let  $\text{Str}_\tau$  denote the class of all  $\tau$ -structures.

(4) Given a language  $\tau$  and  $M, N \in \text{Str}_\tau$ , an isomorphism between  $M$  and  $N$  is a bijection  $\pi : \text{dom}(M) \rightarrow \text{dom}(N)$  satisfying:

(a) If  $c \in \mathfrak{C}_\tau$ , then  $\pi(c^M) = c^N$ .

(b) If  $f \in \mathfrak{F}_\tau$  and  $x_0, \dots, x_{\mathfrak{a}_\tau(f)} \in \text{dom}(M)$ , then

$$\pi(f^M(x_0, \dots, x_{\mathfrak{a}_\tau(f)})) = f^N(\pi(x_0), \dots, \pi(x_{\mathfrak{a}_\tau(f)})).$$

(c) If  $r \in \mathfrak{R}_\tau$  and  $x_0, \dots, x_{\mathfrak{a}_\tau(r)} \in \text{dom}(M)$ , then

$$r^M(x_0, \dots, x_{\mathfrak{a}_\tau(r)}) \iff r^N(\pi(x_0), \dots, \pi(x_{\mathfrak{a}_\tau(r)})).$$

(5) Given a language  $\tau$  and  $M, N \in \text{Str}_\tau$ , the structure  $M$  is a substructure of the structure  $N$  if  $\text{dom}(M) \subseteq \text{dom}(N)$ ,  $c^M = c^N$  for all  $c \in \mathfrak{C}_\tau$ ,  $f^M = f^N \upharpoonright \text{dom}(M)^{\mathfrak{a}_\tau(f)}$  for all  $f \in \mathfrak{F}$  and  $r^M = r^N \cap \text{dom}(M)^{\mathfrak{a}_\tau(r)}$  for all  $r \in \mathfrak{R}$ .

(6) A language  $\sigma$  is a sublanguage of a language  $\tau$  if  $\mathfrak{C}_\sigma \subseteq \mathfrak{C}_\tau$ ,  $\mathfrak{F}_\sigma \subseteq \mathfrak{F}_\tau$ ,  $\mathfrak{R}_\sigma \subseteq \mathfrak{R}_\tau$  and  $\mathfrak{a}_\sigma = \mathfrak{a}_\tau \upharpoonright (\mathfrak{F}_\sigma \cup \mathfrak{R}_\sigma)$ .

(7) Given a sublanguage  $\sigma$  of a language  $\tau$  and  $M \in \text{Str}_\tau$ , the  $\sigma$ -reduct  $M \upharpoonright \sigma$  of  $M$  is the unique element  $N$  of  $\text{Str}_\sigma$  with  $\text{dom}(N) = \text{dom}(M)$ ,  $c^N = c^M$  for all  $c \in \mathfrak{C}_\sigma$ ,  $f^N = f^M$  for all  $f \in \mathfrak{F}_\sigma$  and  $r^N = r^M$  for all  $r \in \mathfrak{R}_\sigma$ .

(8) A renaming of a language  $\sigma$  into a language  $\tau$  is a bijection

$$u : \mathfrak{S}_\sigma \longrightarrow \mathfrak{S}_\tau$$

satisfying  $u[\mathfrak{C}_\sigma] = \mathfrak{C}_\tau$ ,  $u[\mathfrak{F}_\sigma] = \mathfrak{F}_\tau$ ,  $u[\mathfrak{R}_\sigma] = \mathfrak{R}_\tau$  and  $a_\sigma(s) = a_\tau(u(s))$  for all  $s \in \mathfrak{F}_\sigma \cup \mathfrak{R}_\sigma$ .

(9) Given a renaming  $u$  of a language  $\sigma$  into a language  $\tau$ , we let  $u^* : \text{Str}_\sigma \longrightarrow \text{Str}_\tau$  denote the bijective class function with the property that for all  $M \in \text{Str}_\sigma$ , we have  $\text{dom}(u^*(M)) = \text{dom}(M)$  and  $u(s)^{u^*(M)} = s^M$  for all  $s \in \mathfrak{S}_\sigma$ .

Note that, if  $u$  is a renaming of a language  $\sigma$  into a language  $\tau$ , then the inverse  $u^{-1}$  of  $u$  is a renaming of  $\tau$  into  $\sigma$  with the property that  $(u^{-1})^* = (u^*)^{-1}$  holds.

We are now ready to introduce the notion of an abstract logic:

**Definition 5.2.** An abstract logic consists of a class function  $\mathcal{L}$  and a binary class relation  $\models_{\mathcal{L}}$  satisfying the following statements:

- (1) The domain of  $\mathcal{L}$  is the class of all languages.
- (2) If  $M \models_{\mathcal{L}} \phi$  holds, then there exists a language  $\tau$  such that  $M \in \text{Str}_\tau$  and  $\phi \in \mathcal{L}(\tau)$ .
- (3) (Monotonicity) If  $\sigma$  is a sublanguage of  $\tau$ , then  $\mathcal{L}(\sigma) \subseteq \mathcal{L}(\tau)$ .
- (4) (Expansion) If  $\sigma$  is a sublanguage of  $\tau$ ,  $M \in \text{Str}_\tau$  and  $\phi \in \mathcal{L}(\sigma)$ , then

$$M \models_{\mathcal{L}} \phi \iff M \upharpoonright \sigma \models_{\mathcal{L}} \phi.$$

- (5) (Isomorphism) Given a language  $\tau$  and isomorphic  $\tau$ -structures  $M$  and  $N$ , we have

$$M \models_{\mathcal{L}} \phi \iff N \models_{\mathcal{L}} \phi$$

for all  $\phi \in \mathcal{L}(\tau)$ .

- (6) (Renaming) Every renaming  $u$  of a language  $\sigma$  into a language  $\tau$  induces a unique bijection  $u_* : \mathcal{L}(\sigma) \longrightarrow \mathcal{L}(\tau)$  with the property that

$$M \models_{\mathcal{L}} \phi \iff u^*(M) \models_{\mathcal{L}} u_*(\phi)$$

holds for every  $M \in \text{Str}_\sigma$  and every  $\phi \in \mathcal{L}(\sigma)$ .

- (7) (Occurrence number) There exists an infinite cardinal  $o$  with the property that for every language  $\tau$  and every  $\phi \in \mathcal{L}(\tau)$ , there exists a sublanguage  $\sigma$  of  $\tau$  with  $|\mathfrak{S}_\sigma| < o$  and  $\phi \in \mathcal{L}(\sigma)$ . The least such cardinal is called the occurrence number of the logic.

In the following, we just write  $\mathcal{L}$  to denote an abstract logic. We then call the corresponding relation  $\models_{\mathcal{L}}$  the *satisfaction relation* of  $\mathcal{L}$  and, given a language  $\tau$ , we say that  $\mathcal{L}(\tau)$  is the set of  $\tau$ -sentences. In addition, we write  $o_{\mathcal{L}}$  to denote the occurrence number of an abstract logic  $\mathcal{L}$ . Note that the uniqueness of  $u_*$  in item (6) of Definition 5.2 implies that  $(u^{-1})_* = (u_*)^{-1}$  holds for every renaming  $u$ .

The above definitions now allow us to generalize fundamental concepts from first-order logic to all abstract logics.

**Definition 5.3.** Let  $\mathcal{L}$  be an abstract logic.

- (1) An  $\mathcal{L}$ -theory is a set  $T$  with the property that  $T \subseteq \mathcal{L}(\tau)$  for some language  $\tau$ .
- (2) An  $\mathcal{L}$ -theory  $T$  is satisfiable if there is a language  $\tau$  with  $T \subseteq \mathcal{L}(\tau)$  and  $M \in \text{Str}_\tau$  with  $M \models_{\mathcal{L}} \phi$  for all  $\phi \in T$ .
- (3) Given an infinite cardinal  $\kappa$ , an  $\mathcal{L}$ -theory  $T$  is  $<\kappa$ -satisfiable if every subset of  $T$  of cardinality less than  $\kappa$  is satisfiable.
- (4) An infinite cardinal  $\kappa$  is a strong compactness cardinal for  $\mathcal{L}$  if every  $<\kappa$ -satisfiable  $\mathcal{L}$ -theory is satisfiable.

- (5) An infinite cardinal  $\kappa$  is a weak compactness cardinal for  $\mathcal{L}$  if every  $<\kappa$ -satisfiable  $\mathcal{L}$ -theory of cardinality  $\kappa$  is satisfiable.
- (6) Given a language  $\tau$  and  $N \in \text{Str}_\tau$ , an  $\mathcal{L}$ -elementary substructure of  $N$  is a substructure  $M$  of  $N$  with the property that

$$M \models \phi \iff N \models \phi$$

holds for all  $\phi \in \mathcal{L}(\tau)$ .

- (7) An uncountable cardinal  $\kappa$  is a Löwenheim–Skolem–Tarski number for  $\mathcal{L}$  if for every language  $\tau$  with  $|\mathfrak{S}_\tau| < \kappa$  and every  $\tau$ -structure  $N$ , there exists an  $\mathcal{L}$ -elementary substructure  $M$  of  $N$  with  $|\text{dom}(M)| < \kappa$ .
- (8) An uncountable cardinal  $\kappa$  is a weak Löwenheim–Skolem–Tarski number for  $\mathcal{L}$  if for every language  $\tau$  with the property that  $|\mathfrak{S}_\tau| < \kappa$  and every  $\tau$ -structure  $N$  with  $|\text{dom}(N)| = \kappa$ , there exists an  $\mathcal{L}$ -elementary substructure  $M$  of  $N$  with  $|\text{dom}(M)| < \kappa$ .
- (9) An uncountable cardinal  $\kappa$  is a strict Löwenheim–Skolem–Tarski number for  $\mathcal{L}$  if for every language  $\tau$  with the property that  $|\mathfrak{S}_\tau| < \kappa$ , every  $\tau$ -structure  $N$  with  $|\text{dom}(N)| = \kappa$  and every  $\phi \in \mathcal{L}(\tau)$  with  $N \models_{\mathcal{L}} \phi$ , there exists a substructure  $M$  of  $N$  with  $|\text{dom}(M)| < \kappa$  and  $M \models_{\mathcal{L}} \phi$ .

In [5, Section 4], it is verified that Makowsky’s Theorem 1.1 holds true with respect to the above definitions. In another direction, an unpublished result of Stavi shows that the validity of Vopěnka’s Principle is also equivalent to the fact that analogs of the Löwenheim–Skolem Downward Theorem holds for all abstract logics (see [15, Theorem 6]). For sake of completeness and to motivate the notions introduced in items (8) and (9) of Definition 5.3, we present a proof of Stavi’s result.

**Theorem 5.4** (Stavi). *The following schemes are equivalent over ZFC:*

- (1) Vopěnka’s Principle.
- (2) Every abstract logic has a Löwenheim–Skolem–Tarski number.
- (3) For every abstract logic  $\mathcal{L}$ , there exists an uncountable cardinal  $\kappa$  with the property that for every language  $\tau$  with  $|\mathfrak{S}_\tau| < \kappa$ , every  $\tau$ -structure  $N$  and every  $\phi \in \mathcal{L}(\tau)$  with  $N \models_{\mathcal{L}} \phi$ , there exists a substructure  $M$  of  $N$  with  $|\text{dom}(M)| < \kappa$  and  $M \models_{\mathcal{L}} \phi$ .

*Proof.* First, assume that (1) holds and fix an abstract logic  $\mathcal{L}$ . Then there is a natural number  $n > 0$ , a  $\Sigma_n$ -formula  $\varphi(v_0, v_1, v_2)$ , a  $\Sigma_n$ -formula  $\psi(v_0, v_1, v_2)$  and a set  $z$  with the property that the function  $\mathcal{L}$  is defined by the formula  $\varphi(v_0, v_1, v_2)$  and the parameter  $z$  and the relation  $\models_{\mathcal{L}}$  is defined by the formula  $\psi(v_0, v_1, v_2)$  and the parameter  $z$ . By [1, Corollary 4.15], we can now find a  $C^{(n)}$ -extendible cardinal  $\kappa$  (i.e., a cardinal  $\kappa$  with the property that for every  $\eta > \kappa$ , there exists an ordinal  $\zeta$  and an elementary embedding  $j : V_\eta \rightarrow V_\zeta$  with  $\text{crit}(j) = \kappa$ ,  $j(\kappa) > \eta$  and  $j(\kappa) \in C^{(n)}$ ) with  $z \in H(\kappa)$ . Now, fix a language  $\sigma$  with  $|\mathfrak{S}_\sigma| < \kappa$  and a  $\sigma$ -structure  $K$ . Then there exists a renaming  $u$  of  $\sigma$  into a language  $\tau \in V_\kappa$ . Since [1, Proposition 3.4] shows that  $\kappa \in C^{(n+2)}$ , we know that  $\mathcal{L}(\tau)$  is also an element of  $V_\kappa$ . Set  $M = u^*(K)$ . Pick  $\kappa < \eta \in C^{(n+2)}$  with  $M \in V_\eta$ . Then, there is an ordinal  $\zeta$  and an elementary embedding  $j : V_\eta \rightarrow V_\zeta$  with  $\text{crit}(j) = \kappa$ ,  $j(\kappa) > \eta$  and  $j(\kappa) \in C^{(n)}$ . Since  $\tau \in V_\kappa$  and  $j(\tau) = \tau$ , we now know that  $j(M)$  is a  $\tau$ -structure with  $s^{j(M)} = j(s^M)$  for every  $s \in \mathfrak{S}_\tau$ . This shows that  $j(M)$  has a substructure  $N$  with  $\text{dom}(N) = j[\text{dom}(M)]$ . We then know that the map  $j \upharpoonright \text{dom}(M)$  is an isomorphism between  $M$  and  $N$  and this isomorphism is an element of  $V_\zeta$ . In this situation, the correctness properties of  $V_\eta$ , the elementarity of  $j$  and item (5) in Definition 5.2 ensure that

$$V_\zeta \models \psi(M, \phi, z) \iff \psi(N, \phi, z) \tag{A}$$

holds for all  $\phi \in \mathcal{L}(\tau)$ . Moreover, since  $\kappa$  and  $\eta$  are both elements of  $C^{(n)}$ , we know that the formula  $\psi(v_0, v_1, v_2)$  is absolute between  $V_\kappa$  and  $V_\eta$ . Elementarity then ensures that this formula is also absolute between  $V_{j(\kappa)}$  and  $V_\zeta$ . Moreover, since  $j(\kappa)$  is also an element of  $C^{(n)}$ , it follows that  $\psi(v_0, v_1, v_2)$  is absolute between  $V_\eta$  and  $V_\zeta$ . For all  $\phi \in \mathcal{L}(\tau)$ , we then have

$$V_\zeta \models \psi(j(M), \phi, z) \iff V_\eta \models \psi(M, \phi, z) \iff V_\zeta \models \psi(M, \phi, z) \iff V_\zeta \models \psi(N, \phi, z),$$

where the first equivalence is given by elementarity of  $j$ , the second equivalence follows from the above absoluteness considerations and the third equivalence is given by (A). The elementarity of  $j$  now yields a substructure  $S$  of  $M$  with  $|\text{dom}(S)| < \kappa$  and the property that

$$V_\eta \models \psi(M, \phi, z) \iff \psi(S, \phi, z)$$

holds for all  $\phi$  in  $\mathcal{L}(\tau)$ . Since  $\eta$  is an element of  $C^{(n)}$ , this shows that  $S$  is an  $\mathcal{L}$ -elementary submodel of  $M$ . Set  $R = (u^*)^{-1}(M) \in \text{Str}_\sigma$ . Then  $R$  is a submodel of  $K$  with  $|\text{dom}(R)| < \kappa$ . Moreover, we can apply item (6) in Definition 5.2 to conclude that  $R$  is an  $\mathcal{L}$ -elementary submodel of  $K$ . We can conclude that  $\kappa$  is a Löwenheim–Skolem–Tarski number for  $\mathcal{L}$ . These computations show that (1) implies (2).

Now, assume that (3) holds and fix a proper class  $\Gamma$  of graphs. Then there is an abstract logic  $\mathcal{L}_\Gamma$  such that the following statements hold:

- For every language  $\tau$ , the set  $\mathcal{L}_\Gamma(\tau)$  consists of all first-order sentences of  $\tau$  together with a distinguished sentence  $\phi_r$  for every binary relation symbol  $r$  in  $\tau$ .
- Given is a language  $\tau$ ,  $M \in \text{Str}_\tau$  and a first-order sentence  $\phi \in \mathcal{L}_\Gamma(\tau)$ , then  $M \models_{\mathcal{L}_\Gamma} \phi$  holds if and only if  $\phi$  holds in  $M$  (in the sense of first-order structures).
- Given is a language  $\tau$ ,  $M \in \text{Str}_\tau$  and a binary relation symbol  $r$  in  $\tau$ , then  $M \models_{\mathcal{L}_\Gamma} \phi_r$  holds if and only if  $\langle \bar{M}, r^M \rangle$  is a graph that is isomorphic to a graph in  $\Gamma$ .

Our assumption now yields an uncountable cardinal  $\kappa$  with the property that for every language  $\tau$  with  $|\mathfrak{S}_\tau| < \kappa$ , every  $\tau$ -structure  $N$  and every  $\phi \in \mathcal{L}_\Gamma(\tau)$  with  $N \models_{\mathcal{L}_\Gamma} \phi$ , there exists a substructure  $M$  of  $N$  with  $|\text{dom}(M)| < \kappa$  and  $M \models_{\mathcal{L}_\Gamma} \phi$ . Since  $\Gamma$  is a proper class, we can fix a graph  $G$  in  $\Gamma$  that has cardinality greater than  $\kappa$ . Let  $\sigma$  denote a language with a single binary relation symbol  $e$  and let  $\tau$  be a language extending  $\sigma$  that adds function symbols for first-order Skolem functions to  $\sigma$ . View  $G$  as a  $\sigma$ -structure and let  $G'$  be a  $\tau$ -structure whose  $\sigma$ -reduct is  $G$  and that interprets the new function symbols as the corresponding first-order Skolem functions. Then item (4) in Definition 5.2 ensures that  $G' \models_{\mathcal{L}_\Gamma} \phi_e$  holds and, since  $|\mathfrak{S}_\tau| < \kappa$ , we can use our assumption to find a substructure  $H'$  of  $G'$  with  $|H'| < \kappa$  and  $H' \models_{\mathcal{L}_\Gamma} \phi_e$ . Let  $H$  denote the  $\sigma$ -reduct of  $H'$ . Our setup now ensures that  $H$  is a first-order elementary submodel of  $G$  and  $H \models_{\mathcal{L}_\Gamma} \phi_e$  holds. This shows that  $\langle \text{dom}(H), e^H \rangle$  is a graph that is isomorphic to an element  $K$  of  $\Gamma$ . Since  $\text{dom}(H)$  has cardinality less than  $\kappa$ , we know that  $K \neq G$  and  $K$  embeds into  $G$ . These computations show that (1) holds.

This completes the proof of the theorem, because (2) obviously implies (3).  $\square$

Motivated by Theorem 5.4, we will also use the developed theory to prove the following analogs of Theorems 1.4 and 1.6 for strict Löwenheim–Skolem–Tarski numbers in the next section:

**Theorem 5.5.** (1) *The following schemes of sentences are equivalent over ZFC:*

- (a) *Ord is essentially faint.*
  - (b) *Every abstract logic has a strict Löwenheim–Skolem–Tarski number.*
- (2) *The following schemes of sentences are equivalent over ZFC:*
- (a) *Ord is essentially subtle.*

- (b) *Every abstract logic has a stationary class of strict Löwenheim–Skolem–Tarski numbers.*
- (c) *Every abstract logic has a stationary class of weak Löwenheim–Skolem–Tarski numbers.*

In the last part of the next section, we will consider the question of whether the first equivalence can also be extended to characterize the existence of weak Löwenheim–Skolem–Tarski numbers for all abstract logics.

## 6. STRUCTURAL PROPERTIES OF ABSTRACT LOGICS

This section contains the proofs of Theorems 1.4, 1.6 and 5.5. The following lemma shows how strong structural properties of abstract logics can be derived from the existence of  $C^{(n)}$ -weakly shrewd cardinals.

**Lemma 6.1.** *Let  $n > 0$  be a natural number, let  $\varphi(v_0, v_1, v_2)$  and  $\psi(v_0, v_1, v_2)$  be  $\Sigma_n$ -formulas and let  $z$  be a set with the property that there is an abstract logic  $\mathcal{L}$  such that the function  $\mathcal{L}$  is defined by the formula  $\varphi(v_0, v_1, v_2)$  and the parameter  $z$  and the relation  $\models_{\mathcal{L}}$  is defined by the formula  $\psi(v_0, v_1, v_2)$  and the parameter  $z$ . Then the following statements hold:*

- (1) *There exists a cardinal  $\mu \in C^{(1)}$  such that the following statements hold:*
  - (a)  $\mu > o_{\mathcal{L}}$  and  $z \in \mathbf{H}(\mu)$ .
  - (b) *If  $\sigma$  is a language that is an element of  $\mathbf{H}(o_{\mathcal{L}})$ , then  $\mathcal{L}(\sigma) \in \mathbf{H}(\mu)$ .*
  - (c) *If  $\sigma$  is a language that is an element of  $\mathbf{H}(o_{\mathcal{L}})$  and  $\pi$  is a non-trivial permutation of  $\mathcal{L}(\sigma)$ , then there exists a  $\sigma$ -structure  $M_{\sigma, \pi} \in \mathbf{H}(\mu)$  and  $\phi_{\sigma, \pi} \in \mathcal{L}(\sigma)$  with*

$$M_{\sigma, \pi} \models_{\mathcal{L}} \phi_{\sigma, \pi} \iff \neg(M_{\sigma, \pi} \models_{\mathcal{L}} \pi(\phi_{\sigma, \pi})). \quad (\text{B})$$

- (2) *If  $\mu$  is a cardinal with the properties listed in (1),  $\sigma$  is a language in  $\mathbf{H}(o_{\mathcal{L}})$  and  $u$  is a renaming of  $\sigma$  into a language  $\tau$ , then  $u_*$  is the unique bijection  $\varpi$  between  $\mathcal{L}(\sigma)$  and  $\mathcal{L}(\tau)$  with the property that*

$$M \models_{\mathcal{L}} \phi \iff u^*(M) \models_{\mathcal{L}} \varpi(\phi) \quad (\text{C})$$

*holds for every  $\sigma$ -structure  $M$  in  $\mathbf{H}(\mu)$  and every  $\phi \in \mathcal{L}(\sigma)$ .*

- (3) *If  $\mu$  is a cardinal with the properties listed in (1),  $\kappa > \mu$  is a  $C^{(n)}$ -weakly shrewd cardinal and  $\tau$  is a language with  $|\mathfrak{S}_{\tau}| \leq \kappa$  and  $\mathfrak{S}_{\tau} \subseteq \mathbf{H}(\kappa)$ , then for every cardinal  $\kappa < \theta \in C^{(n+2)}$  and every  $y \in \mathbf{H}(\theta)$ , there exists*
  - a cardinal  $\bar{\theta} \in C^{(n)}$ ,
  - a cardinal  $\mu < \bar{\kappa} < \min(\kappa, \bar{\theta})$ ,
  - a sublanguage  $\bar{\tau}$  of  $\tau$  with  $|\mathfrak{S}_{\bar{\tau}}| \leq \bar{\kappa}$  and  $\mathfrak{S}_{\bar{\tau}} = \mathfrak{S}_{\tau} \cap \mathbf{H}(\bar{\kappa})$ ,
  - an elementary submodel  $X$  of  $\mathbf{H}(\bar{\theta})$  with  $\bar{\kappa} \cup \{\bar{\kappa}, \bar{\tau}, \mathcal{L}(\bar{\tau}), z\} \subseteq X$  and
  - an elementary embedding  $j : X \rightarrow \mathbf{H}(\theta)$  with  $j \upharpoonright \bar{\kappa} = \text{id}_{\bar{\kappa}}$ ,  $j(\bar{\kappa}) = \kappa$ ,  $j(z) = z$ ,  $j(\bar{\tau}) = \tau$ ,  $j(\mathcal{L}(\bar{\tau})) = \mathcal{L}(\tau)$ ,  $j \upharpoonright (\mathcal{L}(\bar{\tau}) \cap X) = \text{id}_{\mathcal{L}(\bar{\tau}) \cap X}$  and  $y \in \text{ran}(j)$ .
- (4) *If  $\mu$  is a cardinal with the properties listed in (1), then every  $C^{(n)}$ -weakly shrewd cardinal greater than  $\mu$  is a weak compactness cardinal for  $\mathcal{L}$ .*
- (5) *If  $\mu$  is a cardinal with the properties listed in (1), then every  $C^{(n)}$ -weakly shrewd cardinal greater than  $\mu$  is a strict Löwenheim–Skolem–Tarski number for  $\mathcal{L}$ .*
- (6) *If  $\mu$  is a cardinal with the properties listed in (1), then every  $C^{(n)}$ -weakly shrewd cardinal  $\kappa > \mu$  with the property that  $\nu^{< o_{\mathcal{L}}} \leq \kappa$  holds for all  $\nu < \kappa$  is a weak Löwenheim–Skolem–Tarski number for  $\mathcal{L}$ .*

*Proof.* (1) Note that, given a language  $\sigma$ , an application of (6) in Definition 5.2 to the trivial renaming of  $\sigma$  into itself ensures that the identity function on  $\mathcal{L}(\sigma)$  is the unique permutation  $\pi$  on  $\mathcal{L}(\sigma)$  with the property that

$$M \models_{\mathcal{L}} \phi \iff M \models_{\mathcal{L}} \pi(\phi)$$

holds for every  $\sigma$ -structure  $M$  and every  $\phi \in \mathcal{L}(\sigma)$ . Hence, for every non-trivial permutation  $\pi$  of  $\mathcal{L}(\sigma)$ , there exists a  $\sigma$ -structure  $M_{\sigma,\pi}$  and  $\phi_{\sigma,\pi} \in \mathcal{L}(\sigma)$  with (B). Using this observation, it is now easy to find a cardinal  $\mu$  with the desired properties.

(2) Let  $\mu$  be a cardinal as in (1), let  $\sigma$  be a language in  $\mathsf{H}(o_{\mathcal{L}})$ , let  $u$  be a renaming of  $\sigma$  into a language  $\tau$  and let  $\varpi$  be a bijection  $\pi$  between  $\mathcal{L}(\sigma)$  and  $\mathcal{L}(\tau)$  such that (C) holds. Assume, towards a contradiction that  $\varpi \neq u_*$ . Then  $\pi = (u_*)^{-1} \circ \varpi$  is a non-trivial permutation of  $\mathcal{L}(\sigma)$  and, since  $M_{\sigma,\pi}$  is an element of  $\mathsf{H}(\mu)$ , our assumptions on  $\varpi$  ensure that

$$M_{\sigma,\pi} \models_{\mathcal{L}} \phi_{\sigma,\pi} \iff u^*(M_{\sigma,\pi}) \models_{\mathcal{L}} \varpi(\phi_{\sigma,\pi}) \iff M_{\sigma,\pi} \models_{\mathcal{L}} \pi(\phi_{\sigma,\pi}),$$

contradicting the definitions of  $M_{\sigma,\pi}$  and  $\phi_{\sigma,\pi}$ . This shows that  $\varpi$  is equal to  $u_*$ .

(3) Fix a cardinal  $\mu$  as in (1), a  $C^{(n)}$ -weakly shrewd cardinal  $\kappa > \mu$ , a cardinal  $\kappa < \theta \in C^{(n+2)}$ , an element  $y$  of  $\mathsf{H}(\theta)$  and a language  $\tau$  with  $|\mathfrak{S}_{\tau}| \leq \kappa$  and  $\mathfrak{S}_{\tau} \subseteq \mathsf{H}(\kappa)$ . Using Lemma 3.3, we can find a cardinal  $\theta \in C^{(n)}$ , a cardinal  $\bar{\kappa} < \min(\kappa, \theta)$ , an elementary submodel  $X$  of  $\mathsf{H}(\theta)$  with  $\bar{\kappa} + 1 \subseteq X$  and an elementary embedding  $j : X \rightarrow \mathsf{H}(\theta)$  with  $j \upharpoonright \bar{\kappa} = \text{id}_{\bar{\kappa}}$ ,  $j(\bar{\kappa}) = \kappa$  and  $\mu, \tau, y \in \text{ran}(j)$ . Then  $\mu < \bar{\kappa}$  and the fact that  $\mu \in C^{(1)}$  implies that  $\mathsf{H}(\mu) \cup \{\mathsf{H}(\mu)\} \subseteq X$  with  $j \upharpoonright \mathsf{H}(\mu) = \text{id}_{\mathsf{H}(\mu)}$ . In particular, we know that  $z \in X$  with  $j(z) = z$ . Moreover, our setup ensures that there is a language  $\bar{\tau}$  in  $X$  with  $j(\bar{\tau}) = \tau$ ,  $\mathfrak{S}_{\bar{\tau}} \subseteq \mathsf{H}(\bar{\kappa})$  and  $|\mathfrak{S}_{\bar{\tau}}| \leq \bar{\kappa}$ . It then follows that  $j \upharpoonright \mathfrak{S}_{\bar{\tau}} = \text{id}_{\mathfrak{S}_{\bar{\tau}}}$  and  $\bar{\tau}$  is a sublanguage of  $\tau$  with  $\mathfrak{S}_{\bar{\tau}} = \mathfrak{S}_{\tau} \cap \mathsf{H}(\bar{\kappa})$ . Moreover, since  $\mathcal{L}(\tau)$  is the unique set  $\ell$  such that  $\varphi(\tau, \ell, z)$  holds and  $\mathcal{L}(\bar{\tau})$  is the unique set  $\bar{\ell}$  such that  $\varphi(\bar{\tau}, \bar{\ell}, z)$  holds, the correctness properties of  $\mathsf{H}(\theta)$  and  $\mathsf{H}(\bar{\theta})$  imply that  $\mathcal{L}(\bar{\tau}) \in X$  with  $j(\mathcal{L}(\bar{\tau})) = \mathcal{L}(\tau)$ .

Now, assume, towards a contradiction, that there is  $\phi_* \in \mathcal{L}(\bar{\tau}) \cap X$  with  $j(\phi_*) \neq \phi_*$ . The correctness properties of  $\mathsf{H}(\bar{\theta})$  and (7) in Definition 5.2 then allow us to find a sublanguage  $\tau_0$  of  $\bar{\tau}$  in  $X$  with  $\phi_* \in \mathcal{L}(\tau_0)$  and  $|\mathfrak{S}_{\tau_0}| < o_{\mathcal{L}}$ . We then know that  $\tau_0$  is an element of  $\mathsf{H}(\bar{\kappa}) \cap X$  with  $j(\tau_0) = \tau_0$  and  $\mathcal{L}(\tau_0)$  is an element of  $X$  with  $j(\mathcal{L}(\tau_0)) = \mathcal{L}(\tau_0)$ . This implies that  $j(\phi_*)$  is an element of  $\mathcal{L}(\tau_0)$ . In this situation, we can now pick a language  $\tau_1$  that is an element of  $\mathsf{H}(o_{\mathcal{L}})$  and a renaming  $u$  of  $\tau_1$  into  $\tau_0$  that is an element of  $\mathsf{H}(\bar{\kappa}) \cap X$ . It follows that  $j(u) = u$ . Let  $u^*$  denote the canonical bijection between  $\text{Str}_{\tau_1}$  and  $\text{Str}_{\tau_0}$  induced by  $u$ . Since  $\tau_0$  and  $u$  are both elements of  $\mathsf{H}(\bar{\kappa}) \cap X$ , it follows that  $u^*(M) \in \mathsf{H}(\bar{\kappa}) \cap X$  holds for every  $\tau_1$ -structure  $M$  in  $\mathsf{H}(\mu)$ . In particular, we know that  $j(u^*(M)) = u^*(M)$  holds for every  $\tau_1$ -structure  $M$  in  $\mathsf{H}(\mu)$ . Let  $u_* : \mathcal{L}(\tau_1) \rightarrow \mathcal{L}(\tau_0)$  denote the unique bijection given by (6) in Definition 5.2.

**Claim.** *The map  $u_*$  is an element of  $X$ .*

*Proof of the Claim.* First, since  $\theta$  is an element of  $C^{(n+2)}$ , it follows that  $u_*$  is an element of  $\mathsf{H}(\theta)$ . The elementarity of  $j$  and the correctness properties of  $\mathsf{H}(\bar{\theta})$  then imply that there is a bijection  $\varpi : \mathcal{L}(\tau_1) \rightarrow \mathcal{L}(\tau_0)$  in  $X$  with the property that

$$M \models_{\mathcal{L}} \phi \iff u^*(M) \models_{\mathcal{L}} \varpi(\phi)$$

holds for all  $M \in \text{Str}_{\tau_1} \cap X$  and all  $\phi \in \mathcal{L}(\tau_1)$ . Since  $\mathsf{H}(\mu) \subseteq X$ , we can now apply (2) to conclude that  $\varpi = u_*$ .  $\square$

Since our setup ensures that  $\mathcal{L}(\tau_1) \subseteq X$ , we now know that  $\mathcal{L}(\tau_0) = u_*[\mathcal{L}(\tau_1)] \subseteq X$  and the fact that  $j(\mathcal{L}(\tau_0)) = \mathcal{L}(\tau_0)$  implies that

$$j \upharpoonright \mathcal{L}(\tau_0) : \mathcal{L}(\tau_0) \rightarrow \mathcal{L}(\tau_0)$$

is a bijection. If we now define

$$\pi = (u_*)^{-1} \circ (j \upharpoonright \mathcal{L}(\tau_0)) \circ u_* : \mathcal{L}(\tau_1) \longrightarrow \mathcal{L}(\tau_1),$$

then we have

$$\pi((u_*)^{-1}(\phi_*)) = (u_*)^{-1}(j(\phi_*)) \neq (u_*)^{-1}(\phi_*)$$

and this shows that  $\pi$  is a non-trivial permutation of  $\mathcal{L}(\tau_1)$ . Since  $M_{\tau_1, \pi} \in \mathbf{H}(\mu) \subseteq \mathbf{H}(\bar{\kappa}) \cap X$ , we know that  $j(M_{\tau_1, \pi}) = M_{\tau_1, \pi}$ . Moreover, earlier computations show that  $u^*(M_{\tau_1, \pi})$  is an element of  $X$  with  $j(u^*(M_{\tau_1, \pi})) = u^*(M_{\tau_1, \pi})$ . Since  $u_*(\phi_{\tau_1, \pi}) \in \mathcal{L}(\tau_0) \subseteq X$ , we can now conclude that

$$\begin{aligned} M_{\tau_1, \pi} \models_{\mathcal{L}} \phi_{\tau_1, \pi} &\iff u^*(M_{\tau_1, \pi}) \models_{\mathcal{L}} u_*(\phi_{\tau_1, \pi}) \\ &\iff X \models \psi(u^*(M_{\tau_1, \pi}), u_*(\phi_{\tau_1, \pi}), z) \\ &\iff \mathbf{H}(\theta) \models \psi(u^*(M_{\tau_1, \pi}), (j \circ u_*)(\phi_{\tau_1, \pi}), z) \\ &\iff u^*(M_{\tau_1, \pi}) \models_{\mathcal{L}} (u_* \circ \pi)(\phi_{\tau_1, \pi}) \\ &\iff M_{\tau_1, \pi} \models_{\mathcal{L}} \pi(\phi_{\tau_1, \pi}), \end{aligned}$$

where the first and fifth equivalence is given by (6) in Definition 5.2, the second equivalence is given by the correctness properties of  $\mathbf{H}(\bar{\theta})$ , the third equivalence is given by the elementarity of  $j$  and the fourth equivalence is given by the correctness properties of  $\mathbf{H}(\theta)$ . The above equivalences now contradict the defining properties of  $M_{\tau_1, \pi}$  and  $\phi_{\tau_1, \pi}$ . This shows that  $j \upharpoonright (\mathcal{L}(\bar{\tau}) \cap X) = \text{id}_{\mathcal{L}(\bar{\tau}) \cap X}$ .

(4) Let  $\mu$  be a cardinal as in (1), let  $\kappa > \mu$  be a  $C^{(n)}$ -weakly shrewd cardinal and let  $U$  be a  $<\kappa$ -satisfiable theory of cardinality  $\kappa$ . Since  $o_{\mathcal{L}} < \kappa$ , we can find a language  $\sigma$  with  $|\mathfrak{S}_{\sigma}| \leq \kappa$  and  $U \subseteq \mathcal{L}(\sigma)$ . Then there is a renaming  $u$  of  $\sigma$  into a language  $\tau$  with  $\mathfrak{S}_{\tau} \subseteq \mathbf{H}(\kappa)$ . Set  $T = u_*[U]$ . Then (6) in Definition 5.2 ensures that  $T$  is also  $<\kappa$ -satisfiable. Now, pick a cardinal  $\kappa < \theta \in C^{(n+2)}$  with  $T \in \mathbf{H}(\theta)$  and use (3) to find a cardinal  $\bar{\theta} \in C^{(n)}$ , a cardinal  $\mu < \bar{\kappa} < \min(\kappa, \bar{\theta})$ , a sublanguage  $\bar{\tau}$  of  $\tau$  with  $|\mathfrak{S}_{\bar{\tau}}| \leq \bar{\kappa}$  and  $\mathfrak{S}_{\bar{\tau}} = \mathfrak{S}_{\tau} \cap \mathbf{H}(\bar{\kappa})$ , an elementary submodel  $X$  of  $\mathbf{H}(\bar{\theta})$  with  $\bar{\kappa} \cup \{\bar{\kappa}, \bar{\tau}, \mathcal{L}(\bar{\tau}), z\} \subseteq X$  and an elementary embedding  $j : X \longrightarrow \mathbf{H}(\theta)$  with  $j \upharpoonright \bar{\kappa} = \text{id}_{\bar{\kappa}}$ ,  $j(\bar{\kappa}) = \kappa$ ,  $j(z) = z$ ,  $j(\bar{\tau}) = \tau$ ,  $j(\mathcal{L}(\bar{\tau})) = \mathcal{L}(\tau)$ ,  $j \upharpoonright (\mathcal{L}(\bar{\tau}) \cap X) = \text{id}_{\mathcal{L}(\bar{\tau}) \cap X}$  and  $T \in \text{ran}(j)$ . Pick  $\bar{T} \in X$  with  $j(\bar{T}) = T$ . Then  $\bar{T}$  is a subset of  $\mathcal{L}(\bar{\tau})$  of cardinality  $\bar{\kappa}$  and the fact that  $\bar{\kappa} \subseteq X$  ensures that  $\bar{T} \subseteq X$ . Moreover, since  $j \upharpoonright (\mathcal{L}(\bar{\tau}) \cap X) = \text{id}_{\mathcal{L}(\bar{\tau}) \cap X}$ , it follows that  $\bar{T} = j[\bar{T}] \subseteq T$  and hence  $\bar{T}$  is satisfiable. This shows that there is a  $\bar{\tau}$ -structure  $M$  with the property that  $\psi(M, \phi, z)$  holds for all  $\phi$  in  $\bar{T}$ . The correctness properties of  $\mathbf{H}(\bar{\theta})$  now ensure that this statement holds in  $X$  and hence we know that, in  $\mathbf{H}(\theta)$ , there is a  $\tau$ -structure  $N$  with the property that  $\psi(N, \phi, z)$  holds for all  $\phi$  in  $T$ . Using the correctness properties of  $\mathbf{H}(\theta)$ , we can now conclude that  $T$  is satisfiable and, by (6) in Definition 5.2, we also know that  $U$  is satisfiable.

(5) Let  $\mu$  be a cardinal as in (1), let  $\kappa > \mu$  be a  $C^{(n)}$ -weakly shrewd cardinal, let  $\sigma$  be a language with the property that  $|\mathfrak{S}_{\sigma}| < \kappa$ , let  $I$  be a  $\sigma$ -structure with  $|\text{dom}(I)| = \kappa$  and let  $\phi$  be an element of  $\mathcal{L}(\sigma)$  with  $I \models_{\mathcal{L}} \phi$ . Let  $u$  be a renaming of  $\sigma$  into a language  $\tau \in \mathbf{H}(\kappa)$ . Set  $K = u^*(I)$  and  $\phi_1 = u_*(\phi)$ . Then  $K \models_{\mathcal{L}} \phi_1$  and there is an isomorphism  $\pi$  between  $K$  and a  $\tau$ -structure  $N$  with  $\text{dom}(N) = \kappa$  and  $N \models_{\mathcal{L}} \phi_1$ . Fix a cardinal  $\kappa < \theta \in C^{(n+2)}$ . Then  $K$  is an element of  $\mathbf{H}(\theta)$  and we can use (3) to find a cardinal  $\bar{\theta} \in C^{(n)}$ , a cardinal  $\mu < \bar{\kappa} < \min(\kappa, \bar{\theta})$ , a sublanguage  $\bar{\tau}$  of  $\tau$  with  $|\mathfrak{S}_{\bar{\tau}}| \leq \bar{\kappa}$  and  $\mathfrak{S}_{\bar{\tau}} = \mathfrak{S}_{\tau} \cap \mathbf{H}(\bar{\kappa})$ , an elementary submodel  $X$  of  $\mathbf{H}(\bar{\theta})$  with  $\bar{\kappa} \cup \{\bar{\kappa}, \bar{\tau}, \mathcal{L}(\bar{\tau}), z\} \subseteq X$  and an elementary embedding  $j : X \longrightarrow \mathbf{H}(\theta)$  with  $j \upharpoonright \bar{\kappa} = \text{id}_{\bar{\kappa}}$ ,  $j(\bar{\kappa}) = \kappa$ ,  $j(z) = z$ ,  $j(\bar{\tau}) = \tau$ ,  $j(\mathcal{L}(\bar{\tau})) = \mathcal{L}(\tau)$ ,  $j \upharpoonright (\mathcal{L}(\bar{\tau}) \cap X) = \text{id}_{\mathcal{L}(\bar{\tau}) \cap X}$  and  $N, \phi_1 \in \text{ran}(j)$ . The fact that  $\tau$  is an element of  $\mathbf{H}(\kappa)$  then implies that  $\tau = \bar{\tau} \in \mathbf{H}(\bar{\kappa})$ ,  $\mathfrak{S}_{\tau} \subseteq X$  and  $j \upharpoonright \mathfrak{S}_{\tau} = \text{id}_{\mathfrak{S}_{\tau}}$ . Pick  $M$  in  $X$  with  $j(M) = N$ . Then  $M$  is a  $\tau$ -structure with  $\text{dom}(M) = \bar{\kappa}$  and it follows that  $M$  is a substructure of  $N$ . Next, pick  $\phi_0$  in  $X$  with  $j(\phi_0) = \phi_1$ . Then  $\phi_0 \in \mathcal{L}(\tau)$  and the fact that  $j \upharpoonright (\mathcal{L}(\tau) \cap X) = \text{id}_{\mathcal{L}(\tau) \cap X}$



implies that  $\phi_0 = \phi_1 \in X$ . Moreover, using the elementarity of  $j$  and the correctness properties of  $\mathbf{H}(\theta)$  and  $\mathbf{H}(\bar{\theta})$ , we know that  $M \models_{\mathcal{L}} \phi_1$  holds. Then  $\pi^{-1} \upharpoonright \bar{\kappa}$  is an isomorphism between  $M$  and a substructure  $J$  of  $K$ , and we can apply (5) in Definition 5.2 to show that  $J \models_{\mathcal{L}} \phi_1$ . Finally, we can use (5) in Definition 5.2 to conclude that  $H = (u^*)^{-1}(J)$  is a substructure of  $I$  with  $|\text{dom}(H)| < \kappa$  and  $H \models_{\mathcal{L}} \phi$ .

(6) Given a cardinal  $\mu$  as in (1), a  $C^{(n)}$ -weakly shrewd cardinal  $\kappa > \mu$  satisfying  $\nu^{< o_{\mathcal{L}}} \leq \kappa$  for all  $\nu < \kappa$ , a language  $\sigma$  with  $|\mathfrak{S}_{\sigma}| < \kappa$  and a  $\sigma$ -structure  $I$  with  $|\text{dom}(I)| = \kappa$ , we proceed as in (5) to first find a renaming  $u$  of  $\sigma$  into a language  $\tau \in \mathbf{H}(\kappa)$  and an isomorphism  $\pi$  between  $u^*(I)$  and a  $\tau$ -structure  $N$  with  $\text{dom}(N) = \kappa$ . As above, fix a cardinal  $\kappa < \theta \in C^{(n+2)}$  and use (3) to find a cardinal  $\bar{\theta} \in C^{(n)}$ , a cardinal  $\mu < \bar{\kappa} < \min(\kappa, \bar{\theta})$  with  $\tau \in \mathbf{H}(\bar{\kappa})$ , an elementary submodel  $X$  of  $\mathbf{H}(\bar{\theta})$  with  $\bar{\kappa} \cup \{\bar{\kappa}, \tau, \mathcal{L}(\tau), z\} \subseteq X$  and an elementary embedding  $j : X \rightarrow \mathbf{H}(\theta)$  with  $j \upharpoonright \bar{\kappa} = \text{id}_{\bar{\kappa}}$ ,  $j(\bar{\kappa}) = \kappa$ ,  $j(z) = z$ ,  $j(\tau) = \tau$ ,  $j(\mathcal{L}(\tau)) = \mathcal{L}(\tau)$ ,  $j \upharpoonright (\mathcal{L}(\tau) \cap X) = \text{id}_{\mathcal{L}(\tau) \cap X}$  and  $N \in \text{ran}(j)$ . This setup then ensures that  $\nu^{< o_{\mathcal{L}}} \leq \bar{\kappa}$  holds for all  $\nu < \bar{\kappa}$ .

**Claim.**  $\mathcal{L}(\tau) \subseteq X$ .

*Proof of the Claim.* Fix  $\phi \in \mathcal{L}(\tau)$ . By (7) in Definition 5.2, there is a sublanguge  $\tau_0$  of  $\tau$  with  $|\mathfrak{S}_{\tau_0}| < o_{\mathcal{L}}$  and  $\phi \in \mathcal{L}(\tau_0)$ . Since  $|\mathfrak{S}_{\tau}|^{< o_{\mathcal{L}}} \leq \bar{\kappa} \subseteq X$ , it follows that  $\tau_0$  is an element of  $\mathbf{H}(\bar{\kappa}) \cap X$  and this implies that  $j(\tau_0) = \tau_0$  and  $j(\mathcal{L}(\tau_0)) = \mathcal{L}(\tau_0)$ . Now, we can find a language  $\tau_1 \in \mathbf{H}(o_{\mathcal{L}})$  and a renaming  $v$  of  $\tau_1$  into  $\tau_0$  that is an element of  $X$ . Our setup then ensures that  $\mathcal{L}(\tau_1) \in \mathbf{H}(\mu) \subseteq X$  and  $j(\mathcal{L}(\tau_1)) = \mathcal{L}(\tau_1)$ . Since (2) shows that  $v_*$  is the unique bijection between  $\mathcal{L}(\tau_1)$  and  $\mathcal{L}(\tau_0)$  with the property that (C) holds for all  $\tau_1$ -structures  $M$  in  $\mathbf{H}(\mu)$  and every  $\phi \in \mathcal{L}(\tau_1)$ , the fact that all parameters of this existence statements are elements of  $X$  that are fixed by  $j$  ensures that  $v_*$  is an element of  $X$ . In particular, since  $\mathcal{L}(\tau_1) \subseteq \mathbf{H}(\mu) \subseteq X$ , we know that  $\mathcal{L}(\tau_0) \subseteq X$  and hence  $\phi$  is an element of  $X$ .  $\square$

Now, pick  $M \in X$  with  $j(M) = N$ . Then  $M$  is a  $\tau$ -structure with  $\text{dom}(M) = \bar{\kappa}$ . Moreover, since  $j \upharpoonright \mathfrak{S}_{\tau} = \text{id}_{\mathfrak{S}_{\tau}}$ , we know that  $M$  is a substructure of  $N$ . Fix  $\phi \in \mathcal{L}(\tau)$ . Then  $\phi \in X$  with  $j(\phi) = \phi$  and we can conclude that

$$M \models_{\mathcal{L}} \phi \iff X \models \psi(M, \phi, z) \iff \mathbf{H}(\theta) \models \psi(N, \phi, z) \iff N \models_{\mathcal{L}} \phi.$$

This shows that  $M$  is an  $\mathcal{L}$ -elementary submodel of  $N$  and, by arguing as in the proof of (5), this shows that there is an  $\mathcal{L}$ -elementary submodel  $H$  of  $I$  with  $|\text{dom}(H)| < \kappa$ .  $\square$

By combining the above lemma with Theorems 2.4 and 3.5, we obtain the forward implications in Theorems 1.4, 1.6 and 5.5:

**Corollary 6.2.** (1) *If Ord is essentially faint and  $\mathcal{L}$  is an abstract logic, then there is a proper class of cardinals that are both weak compactness cardinals for  $\mathcal{L}$  and strict Löwenheim–Skolem–Tarski numbers for  $\mathcal{L}$ .*

(2) *If Ord is essentially subtle and  $\mathcal{L}$  is an abstract logic, then the class of cardinals that are both weak compactness cardinals for  $\mathcal{L}$  and weak Löwenheim–Skolem–Tarski numbers for  $\mathcal{L}$  is stationary.*

*Proof.* Given a natural number  $n > 0$ , assume that  $\mathcal{L}$  is an abstract logic with the property that the function  $\mathcal{L}$  and the relation  $\models_{\mathcal{L}}$  are definable by  $\Sigma_n$ -formulas with parameters. Then (4) and (5) of Lemma 6.1 show that all sufficiently large  $C^{(n)}$ -weakly shrewd cardinals are both weak compactness cardinals for  $\mathcal{L}$  and strict Löwenheim–Skolem–Tarski numbers for  $\mathcal{L}$ . Theorem 3.5 then shows that the assumption that Ord is essentially faint implies that there is a proper class of cardinals that are both weak compactness cardinals for  $\mathcal{L}$  and strict Löwenheim–Skolem–Tarski numbers for  $\mathcal{L}$ .

Now, let  $\mathcal{L}$  be an abstract logic and let  $C$  be a closed unbounded class of ordinals. Then there is a natural number  $n > 0$  such that both  $\mathcal{L}$  and  $C$  are definable by  $\Sigma_n$ -formulas with parameters. Since every sufficiently large cardinal in  $C^{(n)}$  is an element of  $C$ , we can combine the above computations with Lemma 4.1 to conclude that every sufficiently large  $C^{(n)}$ -strongly unfoldable cardinal is an element of  $C$  that is both a weak compactness cardinal for  $\mathcal{L}$  and a weak Löwenheim–Skolem–Tarski number for  $\mathcal{L}$ . In particular, Theorem 2.4 shows that the assumption that Ord is essentially subtle implies that every closed unbounded class of ordinals contains a cardinal that is both a weak compactness cardinal for  $\mathcal{L}$  and a weak Löwenheim–Skolem–Tarski number for  $\mathcal{L}$ .  $\square$

By combining the above corollary with arguments contained in [5], we can also use the obtained results to separate the principle “Ord is subtle” from the principle “Ord is essentially subtle”:

**Corollary 6.3.** *If the theory*

$$\text{ZFC} + \text{“Ord is subtle”}$$

*is consistent, then this theory does not prove that Ord is essentially subtle.*

*Proof.* The proof of [5, Theorem 5.6] shows that, if the above theory is consistent, then it is consistent with the assumption that there is no weak compactness cardinal for second-order logic. An application of Corollary 6.2 then shows that this theory does not prove that Ord is essentially subtle.  $\square$

We now work towards proving the backward implications in Theorems 1.4, 1.6 and 5.5. Given a natural number  $n > 0$  and a cardinal  $\kappa$  that is not  $C^{(n)}$ -weakly shrewd, we let  $\theta_\kappa^n$  denote the minimal element  $\theta$  of  $C^{(n)}$  greater than  $\kappa$  with the property that there exists  $z \in \text{H}(\theta)$  such that for all  $\bar{\theta} \in C^{(n)}$ , all cardinals  $\bar{\kappa} < \min(\kappa, \bar{\theta})$  and all elementary submodels  $X$  of  $\text{H}(\bar{\theta})$  with  $\bar{\kappa} + 1 \subseteq X$ , there is no elementary embedding  $j : X \rightarrow \text{H}(\theta)$  with  $j \upharpoonright \bar{\kappa} = \text{id}_{\bar{\kappa}}$ ,  $j(\bar{\kappa}) = \kappa$  and  $z \in \text{ran}(j)$ . In addition, given a natural number  $n > 0$  and a cardinal  $\kappa$  that is not  $C^{(n)}$ -weakly shrewd, we let  $Z_\kappa^n$  denote the non-empty set of all elements  $z$  of  $\text{H}(\theta_\kappa^n)$  with the property that for all  $\bar{\theta} \in C^{(n)}$ , all cardinals  $\bar{\kappa} < \min(\kappa, \bar{\theta})$  and all elementary submodels  $X$  of  $\text{H}(\bar{\theta})$  with  $\bar{\kappa} + 1 \subseteq X$ , there is no elementary embedding  $j : X \rightarrow \text{H}(\theta_\kappa^n)$  with  $j \upharpoonright \bar{\kappa} = \text{id}_{\bar{\kappa}}$ ,  $j(\bar{\kappa}) = \kappa$  and  $z \in \text{ran}(j)$ . In the following, let  $\sigma$  denote a language consisting of a binary relation symbol  $E$  and four constant symbols  $d$ ,  $e$ ,  $f$  and  $g$ . Given a natural number  $m$ , a natural number  $n > 0$  and a cardinal  $\rho \in C^{(1)}$  of uncountable cofinality, we define  $\mathcal{C}_\rho^{m,n}$  to be the class of all  $\sigma$ -structures  $M$  with the property that there exists a (necessarily unique) cardinal  $\rho < \kappa_M \in \text{Lim}(C^{(m)})$  such that the following statements hold:

- There exists a cardinal  $\vartheta > \kappa_M$  with the property that  $\vartheta > \theta_\mu^n$  holds for every cardinal  $\mu$  in  $C^{(m)} \cap (\rho, \kappa_M]$  that is not  $C^{(n)}$ -weakly shrewd and  $\text{dom}(M)$  is an elementary submodel of  $\text{H}(\vartheta)$  of cardinality  $\kappa_M$  with  $\kappa_M + 1 \subseteq \text{dom}(M)$ .
- $E^M = \in \upharpoonright (\text{dom}(M) \times \text{dom}(M))$ ,  $d^M = \kappa_M$  and  $e^M = \rho$ .
- $f^M$  and  $g^M$  are functions whose domains are the set of all cardinals in  $C^{(m)} \cap (\rho, \kappa_M]$  that are not  $C^{(n)}$ -weakly shrewd.
- If  $\mu \in \text{dom}(f^M)$ , then  $f^M(\mu)$  is an elementary submodel of  $\text{H}(\theta_\mu^n)$  of cardinality  $\mu$  with  $\mu + 1 \subseteq f^M(\mu)$  and  $g^M(\mu) \in Z_\mu^n \cap f^M(\mu)$ .

The following lemma shows how the above class of structures can be used to derive the existence of  $C^{(n)}$ -weakly shrewd cardinals from strong structural properties of abstract logics:

**Lemma 6.4.** *Given a natural number  $m$ , a natural number  $n > 0$  and a cardinal  $\rho \in C^{(1)}$  of uncountable cofinality, assume that  $\mathcal{L}$  is an abstract logic with the property that the following statements hold whenever  $\tau$  is a language extending  $\sigma$ :*

- (1) The set  $\mathcal{L}(\tau)$  contains all first-order sentences of  $\tau$ , a distinguished sentence  $\phi_*$  and distinguished sentences  $\phi_{<\xi}$  and  $\phi_{\geq\xi}$  for every ordinal  $\xi \leq \rho$ .
- (2) Given  $M \in \text{Str}_\tau$  and a first-order sentence  $\phi \in \mathcal{L}(\tau)$ , then  $M \models_{\mathcal{L}} \phi$  holds if and only if  $\phi$  holds in  $M$  (in the sense of first-order structures).
- (3) Given  $M \in \text{Str}_\tau$ , then  $M \models_{\mathcal{L}} \phi_*$  holds if and only if  $M \upharpoonright \sigma$  is isomorphic to a structure in  $\mathcal{C}_\rho^{m,n}$ .
- (4) Given  $\xi \leq \rho$  and  $M \in \text{Str}_\tau$ , then  $M \models_{\mathcal{L}} \phi_{<\xi}$  holds if and only if  $\langle \text{dom}(M), E^M \rangle$  is a well-order of order-type less than  $\xi$ .
- (5) Given  $\xi \leq \rho$  and  $M \in \text{Str}_\tau$ , then  $M \models_{\mathcal{L}} \phi_{\geq\xi}$  holds if and only if  $\langle \text{dom}(M), E^M \rangle$  is a well-order of order-type at least  $\xi$ .

If  $\kappa \in \text{Lim}(C^{(m)})$  is either a weak compactness cardinal for  $\mathcal{L}$  or a strict Löwenheim–Skolem–Tarski number for  $\mathcal{L}$ , then  $\kappa > \rho$  and the set  $C^{(m)} \cap (\rho, \kappa]$  contains a  $C^{(n)}$ -weakly shrewd cardinal  $\mu$  with the property that  $\delta^\eta < \mu$  holds for all  $\delta < \mu$  and  $\eta \leq \rho$  with  $\delta^\eta < \kappa$ .

*Proof.* First, assume towards a contradiction, that  $\kappa \leq \rho$ . Define

$$T = \{\phi_{<\kappa}\} \cup \{\phi_{\geq\xi} \mid \xi < \kappa\} \subseteq \mathcal{L}(\sigma).$$

Then  $T$  is an unsatisfiable  $\mathcal{L}$ -theory that is  $<\kappa$ -satisfiable. This shows that  $\kappa$  is not a weak compactness cardinal for  $\mathcal{L}$ . If we now pick a  $\sigma$ -structure  $N$  with  $\text{dom}(N) = \kappa$  and  $E^N = \in \upharpoonright (\kappa \times \kappa)$ , then  $N \models_{\mathcal{L}} \phi_{\geq\kappa}$  holds and  $M \models_{\mathcal{L}} \phi_{\geq\kappa}$  fails for every substructure  $M$  of  $N$  with  $|\text{dom}(M)| < \kappa$ . This shows that  $\kappa$  is also not a strict Löwenheim–Skolem–Tarski number for  $\mathcal{L}$ , contradicting our initial assumptions. Therefore, we know that  $\kappa > \rho$  and we can pick  $M \in \mathcal{C}_\rho^{m,n}$  with  $\kappa_M = \kappa$ .

**Claim.** *There exist  $\sigma$ -structures  $H$  and  $K$  in  $\mathcal{C}_\rho^{m,n}$  with  $\kappa \in \{\kappa_H, \kappa_K\}$  and the property that there is an elementary embedding  $j : \text{dom}(H) \rightarrow \text{dom}(K)$  satisfying  $j(\kappa_H) = \kappa_K$ ,  $j(\rho) = \rho$ ,  $j \upharpoonright (\kappa_H + 1) \neq \text{id}_{\kappa_H + 1}$ ,  $j(f^H) = f^K$  and  $j(g^H) = g^K$ .*

*Proof of the Claim.* First, assume that  $\kappa$  is a weak compactness cardinal for  $\mathcal{L}$ . Let  $\tau$  denote the language that extends  $\sigma$  by a constant symbol  $c$  and a constant symbol  $c_x$  for every  $x \in \text{dom}(M)$ . Next, let  $T$  denote the  $\mathcal{L}(\tau)$ -theory consisting of the union of the first-order elementary diagram of  $M$  using the constant symbols  $c_x$  and the set

$$\{\phi_*\} \cup \{c E d\} \cup \{c \neq c_\alpha \mid \alpha < \kappa\}.$$

This theory has cardinality  $\kappa$  and  $\tau$ -expansions of the structure  $M$  witness that it is  $<\kappa$ -satisfiable. Hence, our assumption implies that  $T$  is satisfiable. Our setup and (5) in Definition 5.2 now ensure that  $T$  has a model whose  $\sigma$ -reduct  $N$  is an element of  $\mathcal{C}_\rho^{m,n}$ . We can now find an elementary embedding  $j : \text{dom}(M) \rightarrow \text{dom}(N)$  with  $j(\kappa) = \kappa_N \geq \kappa$ ,  $j(\rho) = \rho$ ,  $c^N \in \kappa_N \setminus j[\kappa]$ ,  $j(f^M) = f^N$  and  $j(g^M) = g^N$ . In particular, we know that  $j \upharpoonright (\kappa + 1) \neq \text{id}_{\kappa + 1}$ .

Now, assume that  $\kappa$  is a strict Löwenheim–Skolem–Tarski number for  $\mathcal{L}$ . Let  $\tau$  be a language extending  $\sigma$  that adds function symbols for first-order Skolem functions to  $\sigma$  and let  $N$  be  $\tau$ -structure such that  $N \upharpoonright \sigma = M$  and  $N$  interprets all new function symbols as Skolem functions. Then  $N \models_{\mathcal{L}} \phi_*$  and  $|\text{dom}(N)| = \kappa$ . Therefore, our assumptions yield a substructure  $K$  of  $N$  with  $K \models_{\mathcal{L}} \phi_*$  and  $|\text{dom}(K)| < \kappa$ . We can now find a  $\sigma$ -structure  $H$  in  $\mathcal{C}$  that is isomorphic to  $K \upharpoonright \sigma$ . Our setup now ensures that there is an elementary embedding  $j : \text{dom}(H) \rightarrow \text{dom}(M)$  with  $j(\kappa_H) = \kappa$ ,  $j(\rho) = \rho$ ,  $j(f^H) = f^M$  and  $j(g^H) = g^M$ . Moreover, since  $\kappa_H = |\text{dom}(H)| < \kappa$ , we know that  $j \upharpoonright (\kappa_H + 1) \neq \text{id}_{\kappa_H + 1}$ .  $\square$

Fix  $\sigma$ -structures  $H$  and  $K$  in  $\mathcal{C}_\rho^{m,n}$  and an elementary embedding  $j : \text{dom}(H) \rightarrow \text{dom}(K)$  as in the above claim. Since  $j \upharpoonright (\kappa_H + 1) \neq \text{id}_{\kappa_H + 1}$ , we can find an ordinal  $\mu \leq \kappa_H$  with  $j \upharpoonright \mu = \text{id}_\mu$  and

$j(\mu) > \mu$ . The fact that  $\text{dom}(H)$  is an elementary submodel of  $H(\vartheta)$  for some cardinal  $\vartheta > \kappa$  then implies that  $\mu$  is a regular cardinal.

**Claim.**  $\mu > \rho$ .

*Proof of the Claim.* Since  $\rho$  is an element of  $C^{(1)} \cap \kappa_H$ , we know that  $V_\rho \cup \{V_\rho\} \subseteq \text{dom}(H)$ . Moreover, since  $j(\rho) = \rho$ , it follows that  $j(V_\rho) = V_\rho$  and  $j \upharpoonright V_\rho : V_\rho \rightarrow V_\rho$ . In this situation, the fact that  $\rho$  has uncountable cofinality now allows us to use the *Kunen Inconsistency* to conclude that  $j \upharpoonright V_\rho = \text{id}_{V_\rho}$ .  $\square$

**Claim.**  $\mu \in C^{(m)}$ .

*Proof of the Claim.* Assume, towards a contradiction, that  $\mu \notin C^{(m)}$ . Set  $\xi = \min(C^{(m)} \setminus \mu) < \kappa_H$ . Then there is an ordinal  $\eta < \mu$  with the property that the ordinal  $\xi$  is definable in  $\text{dom}(H)$  by a formula using the parameters  $\kappa_H$  and  $\eta$  and the same formula with the parameters  $\kappa_K$  and  $\eta$  defines  $\xi$  in  $\text{dom}(K)$ . It follows that  $j(\xi) = \xi > \mu$ . Since  $C^{(0)}$  is the class of all ordinals, we now know  $m > 0$  and this implies that  $\kappa_H \in C^{(1)}$  and  $V_{\kappa_H} \subseteq \text{dom}(H)$ . The fact that  $j(V_{\xi+2}) = V_{\xi+2}$  then implies that  $j \upharpoonright V_{\xi+2} : V_{\xi+2} \rightarrow V_{\xi+2}$  is a non-trivial elementary embedding, contradicting the *Kunen Inconsistency*.  $\square$

**Claim.** If  $\delta < \mu$  and  $\eta \leq \rho$  with  $\delta^\eta < \kappa$ , then  $\delta^\eta < \mu$ .

*Proof of the Claim.* Assume, towards a contradiction, that there are cardinals  $\delta < \mu$  and  $\eta \leq \rho$  with  $\mu \leq \delta^\eta < \kappa$ . Then  $\text{dom}(H)$  contains an injection  $\iota : \mu \rightarrow {}^\eta\delta$ . Moreover, since  $\delta, \eta < \mu$ ,  $j(\kappa_H) = \kappa_K$  and  $\kappa \in \{\kappa_H, \kappa_K\}$ , the elementarity of  $j$  ensures that  $\delta^\eta < \kappa_H$  and hence we know that  ${}^\eta\delta \subseteq \text{dom}(H)$ . Set  $s = j(\iota)(\mu) \in {}^\eta\delta$ . Since  $j(s) = s$ , we can now find an  $\alpha < \mu$  with  $\iota(\alpha) = s$ . But then  $j(\iota)(\alpha) = j(\iota(\alpha)) = j(s) = j(\iota)(\mu)$ , contradicting the injectivity of  $\iota$ .  $\square$

Since  $\kappa_H \leq j(\kappa_H) = \kappa_K$  and  $\kappa \in \{\kappa_H, \kappa_K\}$ , we know that  $\mu \leq \kappa_H \leq \kappa$  and the above computations show that the cardinal  $\mu$  is an element of the set  $C^{(m)} \cap (\rho, \kappa]$ . Assume, towards a contradiction, that  $\mu$  is not a  $C^{(n)}$ -weakly shrewd cardinal. Then  $\mu \in \text{dom}(f^H)$ . Set  $\nu = j(\mu) > \mu$ . Then  $\nu$  is a regular cardinal in the interval  $(\rho, \kappa_K]$  and elementarity implies that  $\nu \in \text{dom}(f^K)$ . Set  $X = f^H(\mu) \subseteq \text{dom}(H)$  and  $z = g^N(\nu) = j(g^M(\mu)) \in Z_\nu^n$ . Then  $\theta_\mu^n$  is an element of  $C^{(n)}$ ,  $\mu < \min(\nu, \theta_\mu^n)$ ,  $X$  is an elementary submodel of  $H(\theta_\mu^n)$  with  $\mu+1 \subseteq X$  and  $i = j \upharpoonright X : X \rightarrow H(\theta_\mu^n)$  is an elementary embedding with  $i \upharpoonright \mu = \text{id}_\mu$ ,  $i(\mu) = \nu$  and  $z \in \text{ran}(i)$ . But, this contradicts the fact that  $z$  is an element of  $Z_\nu^n$ . This yields the conclusion of the lemma.  $\square$

We can now also combine the above lemma with Theorems 2.4 and 3.5 to derive the backward implications of Theorems 1.4, 1.6 and 5.5. This completes the proofs of these results.

**Corollary 6.5.** (1) *If every abstract logic has either a weak compactness cardinal or a strict Löwenheim–Skolem–Tarski number, then Ord is essentially faint.*  
 (2) *If every abstract logic has, for every natural number  $n$ , either a weak compactness cardinal or a strict Löwenheim–Skolem–Tarski number that is an element of  $C^{(n)}$ , then Ord is essentially subtle.*

*Proof.* (1) If  $n > 0$  is a natural number and  $\rho$  is a cardinal in  $C^{(1)}$  of uncountable cofinality, then there exists an abstract logic  $\mathcal{L}$  that satisfies the statements (1)–(5) listed in Lemma 6.4 for  $m = 0$ ,  $n$  and  $\rho$ . The fact that  $C^{(m)}$  is the class of all ordinals then allows us to apply the lemma to find a  $C^{(n)}$ -weakly shrewd cardinal greater than  $\rho$ . By Theorem 3.5, these computations show that Ord is essentially faint.

(2) Given a natural number  $n > 0$  and a cardinal  $\rho \in C^{(1)}$  of uncountable cofinality, there exists an abstract logic  $\mathcal{L}$  that satisfies the statements (1)–(5) listed in Lemma 6.4 for  $m = n + 1$ ,  $n$  and  $\rho$ . Since  $C^{(n+2)} \subseteq \text{Lim}(C^{(n+1)})$ , our assumptions allows us to use the lemma to find a  $C^{(n)}$ -weakly shrewd cardinal greater than  $\rho$  that is an element of  $C^{(n+1)}$ . We can then combine Theorem 2.4 with Lemma 4.1 to conclude that  $\text{Ord}$  is essentially subtle.  $\square$

In the remainder of this section, we consider the questions whether it is possible to extend the conclusion of Theorem 5.5.(1) to obtain a characterization of the existence of weak Löwenheim–Skolem–Tarski numbers for all abstract logics through the assumption that  $\text{Ord}$  is essentially faint.

**Theorem 6.6.** *The following schemes are equivalent over ZFC:*

- (1) *Every abstract logic has a weak Löwenheim–Skolem–Tarski number.*
- (2) *For every natural number  $n$  and every cardinal  $\eta$ , there is a  $C^{(n)}$ -weakly shrewd cardinal  $\kappa$  with  $\delta^\eta < \kappa$  for all  $\delta < \kappa$ .*

*Proof.* First, assume that (1) holds,  $n > 0$  is a natural number and  $\mu$  is a cardinal. Pick a cardinal  $\rho$  in  $C^{(1)}$  above  $\mu$ . Then there is an abstract logic  $\mathcal{L}$  that satisfies statements (1)–(5) listed in Lemma 6.4 for  $m = 0$ ,  $n$  and  $\rho$  together with the following statements:

- (6) If  $\tau$  is a language, then  $\mathcal{L}(\tau)$  contains the set  $\mathcal{L}_{\rho,\omega}(\tau)$  of all sentences of the infinitary logic  $\mathcal{L}_{\rho,\omega}$  that extends first-order logic by allowing conjunctions and disjunctions of less than  $\rho$ -many formulas.
- (7) Given a language  $\tau$ ,  $M \in \text{Str}_\tau$  and  $\phi \in \mathcal{L}_{\rho,\omega}(\tau)$ , then  $M \models_{\mathcal{L}} \phi$  if and only if  $M \models_{\mathcal{L}_{\rho,\omega}} \phi$ .

Our assumption now ensure that there is a weak Löwenheim–Skolem–Tarski number  $\kappa$  for  $\mathcal{L}$ .

**Claim.** *If  $\delta < \kappa$ , then  $\delta^\eta < \kappa$ .*

*Proof of the Claim.* Assume, towards a contradiction, that there exists a cardinal  $\eta \leq \delta < \kappa$  with  $\delta^\eta \geq \kappa$ . Let  $\tau$  denote a language that is obtained from  $\sigma$  by first adding a constant symbol  $c_\gamma$  for each  $\gamma < \delta$  and then closing the language under Skolem functions for first order formulas. Pick a sufficiently large cardinal  $\theta > \kappa$  and a  $\tau$ -structure  $N$  such that  $\text{dom}(N)$  is an elementary submodel of  $\text{H}(\theta)$  of cardinality  $\kappa$  with  $\kappa + 1 \subseteq \text{dom}(N)$ ,  $E^N = \in \upharpoonright (\text{dom}(N) \times \text{dom}(N))$ ,  $c_\gamma^N = \gamma$  for all  $\gamma \leq \delta$  and all new function symbols are interpreted as the corresponding Skolem functions. By elementarity, we know that the set  $\text{dom}(N) \cap {}^\eta \delta$  has cardinality  $\kappa$ . Since  $\kappa$  is a weak Löwenheim–Skolem–Tarski number for  $\mathcal{L}$ , we can find an  $\mathcal{L}$ -elementary submodel  $M$  of  $N$  with  $|\text{dom}(M)| < \kappa$ . Then  $\bar{M}$  is an elementary submodel of  $\text{H}(\theta)$  with  $\delta + 1 \subseteq \bar{M}$ . Moreover, we know that there is a function  $s \in N \cap {}^\eta \delta$  that is not an element of  $M$ . But, the set  $\mathcal{L}_{\rho,\omega}(\tau)$  contains a sentence  $\phi_s$  that states that there is a function from  $c_\eta$  to  $c_\delta$  that sends  $c_\gamma$  to  $c_{s(\gamma)}$  for all  $\gamma < \eta$ . It then follows that  $N \models_{\mathcal{L}} \phi_s$  holds and  $M \models_{\mathcal{L}} \phi_s$  fails, contradicting the fact that  $M$  is an  $\mathcal{L}$ -elementary submodel of  $N$ .  $\square$

An application of Lemma 6.4 now shows that the interval  $(\rho, \kappa]$  contains a  $C^{(n)}$ -weakly shrewd cardinal  $\mu$  with the property that  $\delta^\eta < \mu$  holds for all  $\delta < \mu$  with  $\delta^\eta < \kappa$ . By the above claim, we can conclude that  $\delta^\eta < \mu$  holds for all  $\delta < \mu$ . This shows that (2) holds in this case.

Now, assume that (2) holds. Pick an abstract logic  $\mathcal{L}$  and a natural number  $n > 0$  such that  $\mathcal{L}$  is definable by a  $\Sigma_n$ -formula with parameters. We can now apply (6) of Lemma 6.1 to see that every sufficiently large  $C^{(n)}$ -weakly shrewd cardinal  $\kappa$  with the property that  $\nu^{<\omega} \leq \kappa$  holds for all  $\nu < \kappa$  is a weak Löwenheim–Skolem–Tarski number for  $\mathcal{L}$ . Since our assumption ensures that there is a proper class of such cardinals, we can conclude that (1) holds in this case.  $\square$

The above result leaves open the question of whether the existence of weak Löwenheim–Skolem–Tarski numbers for all abstract logics is provably equivalent to the assumption that Ord is essentially faint (see Question 7.1 in the next section). Below, it is shown that a separation of these principles would require substantial large cardinal assumptions. Recall that the *Singular Cardinal Hypothesis* SCH states that  $\kappa^{\text{cof}(\kappa)} = \kappa^+$  holds for every singular cardinal  $\kappa$  with  $2^{\text{cof}(\kappa)} < \kappa$ . In the following, we say that the SCH *eventually holds* if this implication holds for eventually all singular cardinals. Remember that a classical result of Solovay in [20] shows that the existence of a strongly compact cardinal implies that the SCH eventually holds.

**Corollary 6.7.** *The following schemes are equivalent over ZFC + “SCH eventually holds”:*

- (1) *Every abstract logic has a weak Löwenheim–Skolem–Tarski number.*
- (2) *Ord is essentially faint.*

*Proof.* In one direction, a combination of Theorem 3.5 with Theorem 6.6 directly shows that (1) implies (2). In the other direction, note that, if the SCH eventually holds, then the proof of [10, Theorem 5.22] shows that for every cardinal  $\mu$ , all sufficiently large weakly inaccessible cardinals  $\kappa$  have the property that  $\nu^{<\mu} < \kappa$  holds for all  $\nu < \kappa$ . In particular, it follows that all sufficiently large  $C^{(n)}$ -weakly shrewd cardinals satisfy the assumptions of (6) in Lemma 6.1 and therefore these cardinals are weak Löwenheim–Skolem–Tarski numbers for abstract logic definable by  $\Sigma_n$ -formulas. This shows that (2) implies (1).  $\square$

## 7. OPEN QUESTIONS

We end this paper by discussing two questions raised by the above results.

First, while the existence of strict Löwenheim–Skolem–Tarski numbers for all abstract logics was shown to be equivalent to the assumption that Ord is essentially faint in Theorem 5.5, it is unclear whether these principles are also equivalent to the existence of weak Löwenheim–Skolem–Tarski numbers for all abstract logics. More specifically, Theorem 6.6 leaves open the possibility that the equivalence stated in Theorem 3.5 can be strengthened so that the existence of  $C^{(n)}$ -weakly shrewd cardinals satisfying the cardinal arithmetic assumptions listed in Theorem 6.6 can be derived from the principle “Ord is essentially faint”.

**Question 7.1.** *Assume that Ord is essentially faint. Given a natural number  $n > 0$  and a cardinal  $\eta$ , is there is a  $C^{(n)}$ -weakly shrewd cardinal  $\kappa$  with  $\delta^n < \kappa$  for all  $\delta < \kappa$ ?*

Note that, as shown in the proof of Corollary 6.7, a negative answer to Question 7.1 implies the consistency of large cardinal assumptions in the region of measurable cardinals of high Mitchell order, because it implies that the SCH fails at a proper class of singular cardinals and such a failure is known to imply the existence of large cardinals in canonical inner models (see, for example, [18]). However, it is not clear if the given fragment of the SCH outright follows from the assumption that Ord is essentially faint.

Second, we ask whether the analysis carried out to separate the principle “Ord is essentially subtle” from the principle “Ord is essentially faint” can be extended to obtain characterizations of the consistency of stronger subtlety properties of the class of ordinals through the existence and non-existence of set-theoretic sentences with certain provable consequences. More specifically, Theorem 1.10 shows that the consistency of the class-version of the large cardinal property of being a regular limit of subtle cardinals is equivalent to the existence of a consistent sentence implying that Ord is essentially faint. In addition, Theorem 1.13 provides a characterization of the consistency of the class-version of subtle limits of subtle cardinals through the non-existence of a sentence axiomatizing

the principle “Ord is essentially subtle” over the principle “Ord is essentially faint”. These results raise the question whether the consistency of the class-version of the next stronger large cardinal property, i.e., the property of being a stationary limit of subtle cardinals, can be characterized in a similar manner.

**Question 7.2.** *Are there analogs of Theorems 1.10 and 1.13 that characterize the consistency of the theory*

ZFC + “The class of subtle cardinals is stationary in Ord”?

#### REFERENCES

1. Joan Bagaria,  $C^{(n)}$ -cardinals, Arch. Math. Logic **51** (2012), no. 3-4, 213–240.
2. Joan Bagaria and Philipp Lücke, *Patterns of Structural Reflection in the large-cardinal hierarchy*, Trans. Amer. Math. Soc. **377** (2024), no. 5, 3397–3447.
3. Joan Bagaria and Jouko Väänänen, *On the symbiosis between model-theoretic and set-theoretic properties of large cardinals*, J. Symb. Log. **81** (2016), no. 2, 584–604.
4. Will Boney, *Model theoretic characterizations of large cardinals*, Israel J. Math. **236** (2020), no. 1, 133–181.
5. Will Boney, Stamatis Dimopoulos, Victoria Gitman, and Menachem Magidor, *Model theoretic characterizations of large cardinals revisited*, Trans. Amer. Math. Soc. **377** (2024), no. 10, 6827–6861.
6. Chen C. Chang and H. Jerome Keisler, *Model theory*, third ed., Studies in Logic and the Foundations of Mathematics, vol. 73, North-Holland Publishing Co., Amsterdam, 1990.
7. Keith J. Devlin, *Constructibility*, Perspectives in Mathematical Logic, Springer-Verlag, Berlin, 1984.
8. Sakaé Fuchino and Hiroshi Sakai, *Weakly extendible cardinals and compactness of extended logics*, preprint, 2022.
9. Peter Holy, Philipp Lücke, and Sandra Müller, *Outward compactness*, preprint, 2024.
10. Thomas Jech, *Set theory*, Springer Monographs in Mathematics, Springer-Verlag, Berlin, 2003, The third millennium edition, revised and expanded.
11. Ronald B. Jensen and Kenneth Kunen, *Some combinatorial properties of  $L$  and  $V$* , handwritten notes, 1969.
12. Philipp Lücke, *Strong unfoldability, shrewdness and combinatorial consequences*, Proc. Amer. Math. Soc. **150** (2022), no. 9, 4005–4020.
13. ———, *Structural reflection, shrewd cardinals and the size of the continuum*, J. Math. Log. **22** (2022), no. 2, Paper No. 2250007, 43.
14. Menachem Magidor, *On the role of supercompact and extendible cardinals in logic*, Israel J. Math. **10** (1971), 147–157.
15. Menachem Magidor and Jouko Väänänen, *On Löwenheim-Skolem-Tarski numbers for extensions of first order logic*, J. Math. Log. **11** (2011), no. 1, 87–113.
16. Johann A. Makowsky, *Vopěnka’s principle and compact logics*, J. Symbolic Logic **50** (1985), no. 1, 42–48.
17. Pierre Matet, *Some aspects of  $n$ -subtlety*, Z. Math. Logik Grundlag. Math. **34** (1988), no. 2, 189–192.
18. William J. Mitchell, *On the singular cardinal hypothesis*, Trans. Amer. Math. Soc. **329** (1992), no. 2, 507–530.
19. Michael Rathjen, *Recent advances in ordinal analysis:  $\Pi_2^1$ -CA and related systems*, Bull. Symbolic Logic **1** (1995), no. 4, 468–485.
20. Robert M. Solovay, *Strongly compact cardinals and the GCH*, Proceedings of the Tarski Symposium, Proc. Sympos. Pure Math., vol. Vol. XXV, Published for the Association for Symbolic Logic by the American Mathematical Society, Providence, RI, 1974, pp. 365–372.
21. Robert M. Solovay, William N. Reinhardt, and Akihiro Kanamori, *Strong axioms of infinity and elementary embeddings*, Ann. Math. Logic **13** (1978), no. 1, 73–116.
22. Andrés Villaveces, *Chains of end elementary extensions of models of set theory*, J. Symbolic Logic **63** (1998), no. 3, 1116–1136.

PHILIPP LÜCKE. FACHBEREICH MATHEMATIK, UNIVERSITÄT HAMBURG, BUNDESSTRASSE 55, HAMBURG, 20146, GERMANY

*Email address:* philipp.luecke@uni-hamburg.de