

Analysis III for engineering study programs

Ingenuin Gasser

Departments of Mathematics
Universität Hamburg

Technische Universität Hamburg–Harburg
Wintersemester 2021/22

based on slides of Prof. Jens Struckmeier from Wintersemester 2020/21

Content of the course Analysis III.

- 1 Partial derivatives, differential operators.
- 2 Vector fields, total differential, directional derivative.
- 3 Mean value theorems, Taylor's theorem.
- 4 Extrem values, implicit function theorem.
- 5 Implicit representation of curves and surfaces.
- 6 Extrem values under equality constraints.
- 7 Newton-method, non-linear equations and the least squares method.
- 8 Multiple integrals, Fubini's theorem, transformation theorem.
- 9 Potentials, Green's theorem, Gauß's theorem.
- 10 Green's formulas, Stokes's theorem.

Chapter 1. Multi variable differential calculus

1.1 Partial derivatives

Let

$f(x_1, \dots, x_n)$ a scalar function depending n variables

Example: The constitutive law of an ideal gas $pV = RT$.

Each of the 3 quantities p (pressure), V (volume) and T (temperature) can be expressed as a function of the others (R is the gas constant)

$$p = p(V, T) = \frac{RT}{V}$$

$$V = V(p, T) = \frac{RT}{p}$$

$$T = T(p, V) = \frac{pV}{R}$$

1.1. Partial derivatives

Definition: Let $D \subset \mathbb{R}^n$ be open, $f : D \rightarrow \mathbb{R}$, $x^0 \in D$.

- f is called **partially differentiable** in x^0 with respect to x_i if the limit

$$\begin{aligned}\frac{\partial f}{\partial x_i}(x^0) &:= \lim_{t \rightarrow 0} \frac{f(x^0 + te_i) - f(x^0)}{t} \\ &= \lim_{t \rightarrow 0} \frac{f(x_1^0, \dots, x_i^0 + t, \dots, x_n^0) - f(x_1^0, \dots, x_i^0, \dots, x_n^0)}{t}\end{aligned}$$

exists. e_i denotes the i -th unit vector. The limit is called **partial derivative** of f with respect to x_i at x^0 .

- If at every point x^0 the partial derivatives with respect to every variable $x_i, i = 1, \dots, n$ exist and if the partial derivatives are **continuous functions** then we call f **continuous partial differentiable** or a \mathcal{C}^1 -function.

Examples.

- Consider the function

$$f(x_1, x_2) = x_1^2 + x_2^2$$

At any point $x^0 \in \mathbb{R}^2$ there exist both partial derivatives and both partial derivatives are continuous:

$$\frac{\partial f}{\partial x_1}(x^0) = 2x_1, \quad \frac{\partial f}{\partial x_2}(x^0) = 2x_2$$

Thus f is a \mathcal{C}^1 -function.

- The function

$$f(x_1, x_2) = x_1 + |x_2|$$

at $x^0 = (0, 0)^T$ is partial differentiable with respect to x_1 , but the partial derivative with respect to x_2 does **not** exist!

An engineering example.

The acoustic pressure of a one dimensional acoustic wave is given by

$$p(x, t) = A \sin(\alpha x - \omega t)$$

The partial derivative

$$\frac{\partial p}{\partial x} = \alpha A \cos(\alpha x - \omega t)$$

describes at a given time t the **spacial** rate of change of the pressure.

The partial derivative

$$\frac{\partial p}{\partial t} = -\omega A \cos(\alpha x - \omega t)$$

describes for a fixed position x the **temporal** rate of change of the acoustic pressure.

Rules for differentiation

- Let f, g be differentiable with respect to x_i and $\alpha, \beta \in \mathbb{R}$, then we have the rules

$$\frac{\partial}{\partial x_i} (\alpha f(x) + \beta g(x)) = \alpha \frac{\partial f}{\partial x_i}(x) + \beta \frac{\partial g}{\partial x_i}(x)$$

$$\frac{\partial}{\partial x_i} (f(x) \cdot g(x)) = \frac{\partial f}{\partial x_i}(x) \cdot g(x) + f(x) \cdot \frac{\partial g}{\partial x_i}(x)$$

$$\frac{\partial}{\partial x_i} \left(\frac{f(x)}{g(x)} \right) = \frac{\frac{\partial f}{\partial x_i}(x) \cdot g(x) - f(x) \cdot \frac{\partial g}{\partial x_i}(x)}{g(x)^2} \quad \text{for } g(x) \neq 0$$

- An alternative notation for the partial derivatives of f with respect to x_i at x^0 is given by

$$D_i f(x^0) \quad \text{oder} \quad f_{x_i}(x^0)$$

Gradient and nabla-operator.

Definition: Let $D \subset \mathbb{R}^n$ be an open set and $f : D \rightarrow \mathbb{R}$ partial differentiable.

- We denote the **row vector**

$$\text{grad } f(x^0) := \left(\frac{\partial f}{\partial x_1}(x^0), \dots, \frac{\partial f}{\partial x_n}(x^0) \right)$$

as **gradient** of f at x^0 .

- We denote the symbolic vector

$$\nabla := \left(\frac{\partial}{\partial x_1}, \dots, \frac{\partial}{\partial x_n} \right)^T$$

as **nabla-operator**.

- Thus we obtain the **column vector**

$$\nabla f(x^0) := \left(\frac{\partial f}{\partial x_1}(x^0), \dots, \frac{\partial f}{\partial x_n}(x^0) \right)^T$$

More rules on differentiation.

Let f and g be partial differentiable. Then the following **rules on differentiation** hold true:

$$\text{grad}(\alpha f + \beta g) = \alpha \cdot \text{grad} f + \beta \cdot \text{grad} g$$

$$\text{grad}(f \cdot g) = g \cdot \text{grad} f + f \cdot \text{grad} g$$

$$\text{grad} \left(\frac{f}{g} \right) = \frac{1}{g^2} (g \cdot \text{grad} f - f \cdot \text{grad} g), \quad g \neq 0$$

Examples:

- Let $f(x, y) = e^x \cdot \sin y$. Then:

$$\text{grad} f(x, y) = (e^x \cdot \sin y, e^x \cdot \cos y) = e^x (\sin y, \cos y)$$

- For $r(x) := \|x\|_2 = \sqrt{x_1^2 + \dots + x_n^2}$ we have

$$\text{grad} r(x) = \frac{x}{r(x)} = \frac{x}{\|x\|_2} \quad \text{für } x \neq 0,$$

where $x = (x_1, \dots, x_n)$ denotes a row vector.

Partial differentiability does not imply continuity.

Observation: A partial differentiable function (with respect to all coordinates) is not necessarily a **continuous** function.

Example: Consider the function $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ defined as

$$f(x, y) := \begin{cases} \frac{x \cdot y}{(x^2 + y^2)^2} & : \text{ for } (x, y) \neq 0 \\ 0 & : \text{ for } (x, y) = 0 \end{cases}$$

The function is partial differentiable on the **entire** \mathbb{R}^2 and we have

$$f_x(0, 0) = f_y(0, 0) = 0$$

$$\frac{\partial f}{\partial x}(x, y) = \frac{y}{(x^2 + y^2)^2} - 4 \frac{x^2 y}{(x^2 + y^2)^3}, \quad (x, y) \neq (0, 0)$$

$$\frac{\partial f}{\partial y}(x, y) = \frac{x}{(x^2 + y^2)^2} - 4 \frac{xy^2}{(x^2 + y^2)^3}, \quad (x, y) \neq (0, 0)$$

Example (continuation).

We calculate the partial derivatives at the origin $(0, 0)$:

$$\frac{\partial f}{\partial x}(0, 0) = \lim_{t \rightarrow 0} \frac{f(t, 0) - f(0, 0)}{t} = \frac{t \cdot 0}{(t^2 + 0^2)^2} - 0 = 0$$

$$\frac{\partial f}{\partial y}(0, 0) = \lim_{t \rightarrow 0} \frac{f(0, t) - f(0, 0)}{t} = \frac{0 \cdot t}{(0^2 + t^2)^2} - 0 = 0$$

But: At $(0, 0)$ the function is **not** continuous since

$$\lim_{n \rightarrow \infty} f\left(\frac{1}{n}, \frac{1}{n}\right) = \frac{\frac{1}{n} \cdot \frac{1}{n}}{\left(\frac{1}{n} \cdot \frac{1}{n} + \frac{1}{n} \cdot \frac{1}{n}\right)^2} = \frac{\frac{1}{n^2}}{\frac{4}{n^4}} = \frac{n^2}{4} \rightarrow \infty$$

and thus we have

$$\lim_{(x,y) \rightarrow (0,0)} f(x, y) \neq f(0, 0) = 0$$

Boundedness of the derivatives implies continuity.

To guarantee the continuity of a partial differentiable function we need additional conditions on f .

Theorem: Let $D \subset \mathbb{R}^n$ be an open set. Let $f : D \rightarrow \mathbb{R}$ be partial differentiable in a neighborhood of $x^0 \in D$ and let the partial derivatives $\frac{\partial f}{\partial x_i}$, $i = 1, \dots, n$, be **bounded**. Then f is **continuous** in x^0 .

Attention: In the previous example the partial derivatives are **not** bounded in a neighborhood of $(0,0)$ since

$$\frac{\partial f}{\partial x}(x, y) = \frac{y}{(x^2 + y^2)^2} - 4 \frac{x^2 y}{(x^2 + y^2)^3} \quad \text{für } (x, y) \neq (0, 0)$$

Proof of the theorem.

For $\|x - x^0\|_\infty < \varepsilon$, $\varepsilon > 0$ sufficiently small we write:

$$\begin{aligned} f(x) - f(x^0) &= (f(x_1, \dots, x_{n-1}, x_n) - f(x_1, \dots, x_{n-1}, x_n^0)) \\ &+ (f(x_1, \dots, x_{n-1}, x_n^0) - f(x_1, \dots, x_{n-2}, x_{n-1}^0, x_n^0)) \\ &\vdots \\ &+ (f(x_1, x_2^0, \dots, x_n^0) - f(x_1^0, \dots, x_n^0)) \end{aligned}$$

For any difference on the right hand side we consider f as a function in one single variable:

$$g(x_n) - g(x_n^0) := f(x_1, \dots, x_{n-1}, x_n) - f(x_1, \dots, x_{n-1}, x_n^0)$$

Since f is partial differentiable g is differentiable and we can apply the mean value theorem on g :

$$g(x_n) - g(x_n^0) = g'(\xi_n)(x_n - x_n^0)$$

for an appropriate ξ_n between x_n and x_n^0 .

Proof of the theorem (continuation).

Applying the **mean value theorem** to every term in the right hand side we obtain

$$\begin{aligned} f(x) - f(x^0) &= \frac{\partial f}{\partial x_n}(x_1, \dots, x_{n-1}, \xi_n) \cdot (x_n - x_n^0) \\ &+ \frac{\partial f}{\partial x_{n-1}}(x_1, \dots, x_{n-2}, \xi_{n-1}, x_n^0) \cdot (x_{n-1} - x_{n-1}^0) \\ &\vdots \\ &+ \frac{\partial f}{\partial x_1}(\xi_1, x_2^0, \dots, x_n^0) \cdot (x_1 - x_1^0) \end{aligned}$$

Using the boundedness of the partial derivatives

$$|f(x) - f(x^0)| \leq C_1|x_1 - x_1^0| + \dots + C_n|x_n - x_n^0|$$

for $\|x - x^0\|_\infty < \varepsilon$, we obtain the **continuity** of f at x^0 since

$$f(x) \rightarrow f(x^0) \quad \text{für } \|x - x^0\|_\infty \rightarrow 0$$

Higher order derivatives.

Definition: Let f be a scalar function and partial differentiable on an open set $D \subset \mathbb{R}^n$. If the partial derivatives are differentiable we obtain (by differentiating) the **partial derivatives of second order** of f with

$$\frac{\partial^2 f}{\partial x_j \partial x_i} := \frac{\partial}{\partial x_j} \left(\frac{\partial f}{\partial x_i} \right)$$

Example: Second order partial derivatives of a function $f(x, y)$:

$$\frac{\partial^2 f}{\partial x^2} = \frac{\partial}{\partial x} \left(\frac{\partial f}{\partial x} \right), \quad \frac{\partial^2 f}{\partial y \partial x} = \frac{\partial}{\partial y} \left(\frac{\partial f}{\partial x} \right), \quad \frac{\partial^2 f}{\partial x \partial y}, \quad \frac{\partial^2 f}{\partial y^2}$$

Let $i_1, \dots, i_k \in \{1, \dots, n\}$. Then we define recursively

$$\frac{\partial^k f}{\partial x_{i_k} \partial x_{i_{k-1}} \dots \partial x_{i_1}} := \frac{\partial}{\partial x_{i_k}} \left(\frac{\partial^{k-1} f}{\partial x_{i_{k-1}} \partial x_{i_{k-2}} \dots \partial x_{i_1}} \right)$$

Higher order derivatives.

Definition: The function f is called k -times partial differentiable, if all derivatives of order k ,

$$\frac{\partial^k f}{\partial x_{i_k} \partial x_{i_{k-1}} \dots \partial x_{i_1}} \quad \text{for all } i_1, \dots, i_k \in \{1, \dots, n\},$$

exist on D .

Alternative notation:

$$\frac{\partial^k f}{\partial x_{i_k} \partial x_{i_{k-1}} \dots \partial x_{i_1}} = D_{i_k} D_{i_{k-1}} \dots D_{i_1} f = f_{x_{i_1} \dots x_{i_k}}$$

If all the derivatives of k -th order are continuous the function f is called k -times continuous partial differentiable or called a C^k -function on D . Continuous functions f are called C^0 -functions.

Example: For the function $f(x_1, \dots, x_n) = \prod_{i=1}^n x_i^i$ we have $\frac{\partial^n f}{\partial x_n \dots \partial x_1} = ?$

Partial derivatives are not arbitrarily exchangeable.

ATTENTION: The order how to execute partial derivatives is in general **not** arbitrarily exchangeable!

Example: For the function

$$f(x, y) := \begin{cases} xy \frac{x^2 - y^2}{x^2 + y^2} & : \text{ for } (x, y) \neq (0, 0) \\ 0 & : \text{ for } (x, y) = (0, 0) \end{cases}$$

we calculate

$$f_{xy}(0, 0) = \frac{\partial}{\partial y} \left(\frac{\partial f}{\partial x}(0, 0) \right) = -1$$

$$f_{yx}(0, 0) = \frac{\partial}{\partial x} \left(\frac{\partial f}{\partial y}(0, 0) \right) = +1$$

i.e. $f_{xy}(0, 0) \neq f_{yx}(0, 0)$.

Theorem of Schwarz on exchangeability.

Satz: Let $D \subset \mathbb{R}^n$ be open and let $f : D \rightarrow \mathbb{R}$ be a \mathcal{C}^2 -function. Then it holds

$$\frac{\partial^2 f}{\partial x_j \partial x_i}(x_1, \dots, x_n) = \frac{\partial^2 f}{\partial x_i \partial x_j}(x_1, \dots, x_n)$$

for all $i, j \in \{1, \dots, n\}$.

Idea of the proof:

Apply the mean value theorem twice.

Conclusion:

If f is a C^k -function, then we can exchange the differentiation in order to calculate partial derivatives up to order k **arbitrarily!**

Example for the exchangeability of partial derivatives.

Calculate the partial derivative of third order f_{xyz} for the function

$$f(x, y, z) = y^2 z \sin(x^3) + (\cosh y + 17e^{x^2})z^2$$

The order of execution is exchangeable since $f \in \mathcal{C}^3$.

- Differentiate first with respect to z :

$$\frac{\partial f}{\partial z} = y^2 \sin(x^3) + 2z(\cosh y + 17e^{x^2})$$

- Differentiate then f_z with respect to x (then $\cosh y$ disappears):

$$\begin{aligned} f_{zx} &= \frac{\partial}{\partial x} \left(y^2 \sin(x^3) + 2z(\cosh y + 17e^{x^2}) \right) \\ &= 3x^2 y^2 \cos(x^3) + 68xze^{x^2} \end{aligned}$$

- For the partial derivative of f_{zx} with respect to y we obtain

$$f_{xyz} = 6x^2 y \cos(x^3)$$

The Laplace operator.

The **Laplace-operator** or **Laplacian** is defined as

$$\Delta := \sum_{i=1}^n \frac{\partial^2}{\partial x_i^2}$$

For a scalar function $u(x) = u(x_1, \dots, x_n)$ we have

$$\Delta u = \sum_{i=1}^n \frac{\partial^2 u}{\partial x_i^2} = u_{x_1 x_1} + \dots + u_{x_n x_n}$$

Examples of important partial differential equations of second order (i.e. equations containing partial derivatives up to order two):

$$\Delta u - \frac{1}{c^2} u_{tt} = 0 \quad (\text{wave equation})$$

$$\Delta u - \frac{1}{k} u_t = 0 \quad (\text{heat equation})$$

$$\Delta u = 0 \quad (\text{Laplace-equation or equation for the potential})$$

Vector valued functions.

Definition: Let $D \subset \mathbb{R}^n$ be open and let $f : D \rightarrow \mathbb{R}^m$ be a vector valued function.

The function f is called **partial differentiable** on $x^0 \in D$, if for all $i = 1, \dots, n$ the limits

$$\frac{\partial f}{\partial x_i}(x^0) = \lim_{t \rightarrow 0} \frac{f(x^0 + te_i) - f(x^0)}{t}$$

exist. The calculation is done componentwise

$$\frac{\partial f}{\partial x_i}(x^0) = \begin{pmatrix} \frac{\partial f_1}{\partial x_i} \\ \vdots \\ \frac{\partial f_m}{\partial x_i} \end{pmatrix} \quad \text{for } i = 1, \dots, n$$